

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/






•



.



· · · · · · · · · · · · · · · · · · ·	·		
	•		

1878.]

CITY DOCUMENT.

[No. 24.

TWENTY-THIRD, ANNUAL REPORT

CPON THE

BIRTHS, MARRIAGES, AND DEATHS

IN THE

CITY OF PROVIDENCE,

FOR THE YEAR

1877,

By EDWIN M. SNOW, M. D., superintendent of health and city registrar.



PROVIDENCE:
J. A. & R. A. REID, PRINTERS.
1878.



TWENTY-THIRD ANNUAL REPORT

UPON THE .

BIRTHS, MARRIAGES, AND DEATHS

IN THE

CITY OF PROVIDENCE,

FOR THE YEAR

1877,


By EDWIN M. SNOW, M. D.,
SUPERINTENDENT OF HEALTH AND CITY REGISTRAR.



PROVIDENCE:

J. A. & R. A. REID, PRINTERS.

1878.



THE CITY OF PROVIDENCE.

JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved September 9, 1878.]

RESOLVED, That the Twenty-third annual report of the City Registrar, for the year ending December 31, 1877, be received, and that six hundred copies thereof, and of the communication accompanying the same, be printed in pamphlet form for the use of the City Council, under the direction of the joint committee on printing.

A true copy, witness,

SAMUEL W. BROWN, CITY CLERK.

INTRODUCTION.

CITY REGISTRAR'S OFFICE,
PROVIDENCE, July 30, 1878.

To the Honorable the City Council:

The Twenty-Third Annual Report upon the Births, Marriages, and Deaths in the City of Providence is herewith presented. It contains the statistics for the year ending December 31, 1877. During the year 1877, there were 2,543 births in Providence, 1,906 persons were married, and 1,938 persons died. This is a decrease of 88 in the number of children born, an increase of 8 in the persons married, and an increase of 73 in the number of deaths as compared with the previous year.

The census of the city in January of the present year, showed that there had been no increase of population since the State census of 1875. We therefore call the population 100,675 for the middle of the year 1877, the same as in 1875. The ratio of births, marriages, and deaths to population, in 1877, was therefore as follows:—

One birth in 39.58; or 25.28 in each 1,000.

One person married in 52.81; or 18.93 in each 1,000.

One death in 51.94; or 19.25 in each 1,000.

During the whole period of twenty-two years, from 1855 to 1876 inclusive, the ratio of births, marriages, and deaths to population in Providence was as follows:

One birth in 34.63; or 28.87 in each 1,000.

One person married in 39.53; or 25.29 in each 1,000.

One death in 50.70; or 19.72 in each 1,000.

During these years there has been a considerable decrease in the proportion of births and of marriages to population; and some decrease in the proportion of deaths.

A complete alphabetical index of the names of all the persons whose births, marriages, or deaths are recorded in Providence from the year 1636 to 1850 inclusive has been prepared, and is now printing. Besides the names of the persons, it contains the dates of the events recorded, and the volume and page of the city records where the names are found. This will be of great value for reference, and will do much for the preservation of the record books.

Respectfully,

EDWIN M. SNOW, M. D.,
Superintendent of Health and City Registrar.

CONTENTS.

Births.	PAGE. 1-14
Wards, months, sex, and color, (Table I.)	1
Births in connection with locality, sex, and season, (Table II) Proportion of the sexes born at different seasons	3
Number and proportion of the sexes ; 24 years Color : — Births, marriages, and deaths in 22 years. Parentage of children born in each ward. (Table III).	4-5
Parentage of children born in each ward. (Table III)	5
Number and percentage of American and of foreign children, 22 years.	ti-7
Parentage: — Mixture of different nations, (Table IV) Number of 1st, 2d, etc., children in 1877, and 1855–1876, (Table V)	8
Average number of children to each mother, 23 years, (Table VI)	10-11
Plurality births; statistics for 1877	12
Plurality births; 23 years' summary	
Marriages at different seasons, 23 years, (Table VIII)	15-16
Birthplaces of parties married in 1877, (Table 1X)	16
Bride and groom born in same country, etc	17
Proportions of American, foreign, and mixed marriages; 23 years	18-19
Average age of persons married. Marriages of colored persons. Number of times married, (Table XI) Denominational. Nativity and residence.	19 20
Denominational. Nativity and residence	21
Death to population in 1877. Proportion in each class. Deaths by sex, condition, color, nativity, parentage, etc., (Table XII) Deaths by season: number and proportions, recapitulation.	23
Deaths by sex, condition, color, nativity, parentage, etc., (Table XII)	95 98
Sex, condition, color, nativity, parentage, and locality of deaths, 1877	26-28
Aggregate and average age by parentage, 1877, (Table XIII)	28
Difference between average age of American and foreign decedents	
Percentages at different ages by parentage, (Table XVI)	30
Percentages under 5 years of age by parentage, 22 years, (Table)	33
Deaths by age and parentage in each month and quarter, 1877, (Table	3
XVIII)	35-37
Causes of death; sex, age, parentage, and percentages, (Table XIX) Causes of death in connection with season, (Table XX)	42-43
Notes on the Causes of Death in 1877	44-67
Cholera infantum; decedents by sex and parentage, 22 years	47-48
Diarrhea and dysentery; recapitulation for 38 years	51-52
Diphtheria; statistics for 1877	58 54
Diphtheria in Providence, 20 years, (Table) Diphtheria; season and mortality, compared with scarlatina and croup,	OF
(Tables)	57
Meningitis, cerebro-spinal; deaths at each age; 14 years. Scarlatina; statistics for 1877.	60-61
Scarlating in Providence 23 years (Table)	42.4
Small-pox; recapitulatory table, 22 years. Still-born children in 1877.	65
Still-born children, 1856-1877	68-69
GENERAL SUMMARY, (Table XXI)	. 70
Births and deaths by parentage. Recapitulatory table	
CITY ORDINANCE RELATING TO REGISTRATION	70

CITY REGISTRAR'S REPORT.

1877.

BIRTHS.

THERE were 2,543 births in the city of Providence, during the year 1877. The following table shows the number in each ward, in each month and quarter of the year; also the number of each sex, and the number of colored children, born in the whole city in each month and quarter of the year:—

TABLE I. BIRTES, 1877. Wards, Months, Sex, and Color.

PROVIDENCE.		WARDS.								w	Whole City.			Colored children included in preceding.		
1877.	ı.	п.	m.	ıv.	v.	V1.	VII.	VIII	ıx,	x.	Males.	Females.	Monthly Totals.	M.	F,	Totals,
January February March	24 25 36	2 6 5	14 21 18	5 9 6	18 17 23	7 10 8	12 17 15	18 22 23	31 35 36	35 33 41	101	94	195		3 6 4	8 15 10
1st Quarter	85	13	53	20	58	25	44	63	102	109	316	256	572	20	13	33
April May June	25 28 24	4 3 6	20 15 26	7 12	9 13 12	7 4 14	10 14 23	15 26 16	28 36 26	46 36 54	81 100 99	90 82 114	171 182 213	5 7 8	313	8 8 11
2d Quarter	77	13	61	26	34	25	47	57	90	136	280	286	566	20	7	27
July August September	28 33 39	6 3 7	26 18 18	13 10 7	20 24 19	14 11 11	21 17 12	22 32 21	39 48 51	33 40 32	117 117 99	105 119 118	222 236 217	3 5 4	3 5 9	6 10 6
3d Quarter	100	16	62	30	63	36	50	75	138	105	333	342	675	12	10	22
October November December	31 37 38	9 10 4	17 20 21	6 15 7	14 16 21	12 13 15	20 23 16	25 28 27	57 42 31	54 45 56	126 124 123	119 125 113	245 249 236	2 2	9 3 2	13 5 4
4th Quarter	106	23	58	28	51	40	59	80	130	155	373	357	730	8	14	22
Whole year	368	65	234	104	206	126	200	275	460	505	1,302	1,241	2,543	60	44	104

The number of births continues to decrease in Providence. The number was 88 less (57 males, 31 females) in 1877 than in 1876; and was 237 less than in 1874. There was an increase of 82 in the first, second, fifth, sixth, and ninth wards collectively, and a decrease of 170 in the other wards.

We estimate the population at 100,675, in the middle of the year 1877, the same number as in 1875. The ratio of births to population, in 1877, was therefore one in 39.58, or 25.25 in each 1,000. This rate is lower than at any previous period, except in the last three years of the war of the rebellion.

Season. The largest monthly total of births, in 1877, was 249 in November; the smallest monthly total was 166 in January. The largest quarterly total was 730 in the fourth quarter of the year; the smallest, 566, in the second quarter. There were 1,138 births in the first half of the year 1877, and 1,405 in the last half, a difference of 267. This excess is very much greater than has been usual in Providence.

The number of births in each quarter of the year in 1877, compared with the aggregate for twenty-one years, 1856-1876 inclusive, was as follows:—

Twenty-one years	Jan. March.	April. June,	First 6 mos.	July. Sept.	Oct. Dec.	Second 6 mos.	Whole Year.
1877			. 1,138	The second second		. 1,405	
Twenty-two years	10.094	10.286	.20.380	10.191	10.702	.20.893	41.273

In a period of twenty-three years, from 1855 to 1877 inclusive, the monthly totals of the births in Providence were as follows: the daily average and the total of births for each month, in the aggregate for the whole period are also given:—

Births in	January.	February.	March.	April.	May.	June.
23 years, 1855-1877	3,369	3,292	3,790	3,292	3,576	3,781
Daily average	4.73	5.06	5.32	4.77	5.02	5.48
Births in	July.	August.	Sept.	Oct.	Nov.	Dec.
23 years, 1855-1877	3,521	3,586	3,499	3,667	3,644	3,856
Daily average	4.94	5.03	5.07	5.14	5.28	5.41
Whole number of b	irths in 23 y	years, 42,87	3. Daily	average.	5.10.	

There were five months in which the daily average exceeded the general average for the period: March, June, October, November, and December; the largest daily average was in June; the least, in January; the largest monthly total, in December; the smallest, in February and April.

Sex and Locality. Table II gives the number of births, and the number of each sex in each ward of the city, in each quarter and in the whole year 1877:—

	1			2d Quarter. 3d Quarter.			441	h Qu	erter.	Whole Year.					
WARDS.	M.	F.	Total.	M.	F.	Total.	M.	F.	Total.	M.	F.	Total.	¥.	F.	Total.
I	46 8	39 5	85 18 53 20	36 11 29	41 2 32	77 18 61	49	51 9	16	50 15 29	8	28	181 41	187 24 128	65
ΪΫ Ψ ΨΪ	25 12 29	28 8 29 11	20 58 25	14 16	12	26 84	28 14 26	34 16 37	63	12 27 22	16 24	28 51 40	1111 522 98 65	52 108 61	104
VII	14 26 38	18 25	44 63	10 23 37	24 20	25 47 57 90	19 21 39	17 29 36 58	86 50 75	20 44	18 39 36	59 80	90 158 256	110 117 204	200 275
X	58 58		102	63	73	136	80 50	55 	138 105	75 79	55 76	180 155	250	255	480 505

TABLE II. BIRTHS, 1877. Wards, Sex, and Season.

Proportion of the Sexes. The whole number of children born in the city in 1877 was 2,543; males, 1,302; females, 1,241; this gives 104.9 males to each 100 females; or 51.20 males, and 48.80 females in each 100 children born.

The number and proportion of each sex, in each quarter of the year 1877, were as follows:—

	Males.	Fe	males	L	Males.	F	emale	16.	Males.	1	Female	6.			
JanMarch	316	and	256,	or	123.4	to	100,	or	55.24	and	44.76	in	each	100 1	oorn.
April-June	280	and	286,	or	97.9	to	100,	or	49.47	and	50.53	in	each	100	born.
July-Sept	333	and	342,	or	97.4	to	100,	or	49.33	and	50.67	in	each	100	born.
OctDec	. 373	and	357,	or	104.5	to	100,	or	51.10	and	48.90	in	each	100 I	born.

Whole Year...1,302 and 1,241, or 104.9 to 100, or 51.20 and 48.80, in each 100 born.

The same statistics are given below for the period of twentytwo years, 1856-1877 inclusive; the number of children born during the period being 41,273 (males, 21,195; females, 20,078):—

```
Males. Females. Males. Females. Males. Females. Jan.-March...5,214 and 4,880, or 106.8 to 100, or 51.65 and 48.35 in each 100 born. April-June...5,314 and 4,972, or 106.9 to 100, or 51.66 and 48.34 in each 100 born. July-Sept....5,155 and 5,036, or 102.4 to 100, or 50.58 and 49.42 in each 100 born. Oct.-Dec.....5,512 and 5,190, or 106.2 to 100, or 51.50 and 48.50 in each 100 born.
```

²² yrs. total..21,195 and 20,078, or 105.6 to 100, or 51.35 and 48.65 in each 100 born.

In the twenty-four years, 1854-1877 inclusive, there were 44,242 children born in Providence; the number and proportion of each sex in each year, were as follows:—

```
Females, Males, Females, Males,
1854.....711 and
                       658; or 108,0 to 100; or 51.91 and 48.09 in each 100 born.
1855......807 and
                       793; or 101.7 to 100; or 50.41 and 49.56 in each 100 born.
1856......891 and
                       784; or 113.6 to 100; or 53.19 and 46.81 in each 100 born.
1857......872 and
                       816; or 106.8 to 100; or 51.66 and 48.34 in each 100 born.
1858.....890 and
                       834; or 106.7 to 100; or 51.62 and 48.38 in each 100 born.
1859......825 and
                       768; or 107.4 to 100; or 51.79 and 48.21 in each 100 born.
1860.....809 and
                       839; or 96.4 to 100; or 49.09 and 50.91 in each 100 born.
738; or 107.2 to 100; or 51.73 and 48.27 in each 100 born.
1862.....791 and
1863.....689 and
                       618; or 111.4 to 100; or 52.71 and 47.29 in each 100 born.
1864......663 and
                       681; or 97.3 to 100; or 49.33 and 50.67 in each 100 born.
                       607; or 113.8 to 100; or 53.24 and 46.76 in each 100 born.
1865......691 and
1866......849 and
                       783; or 108.4 to 100; or 52.02 and 47.98 in each 100 born.
1867......830 and
                       794; or 104.5 to 100; or 51.11 and 48.89 in each 100 born.
1868.....944 and
                       922; or 102.4 to 100; or 50.59 and 49.41 in each 100 born.
1869......945 and
                       874; or 108.1 to 100; or 51.95 and 48.05 in each 100 born.
1870.........967 and 922; or 104.8 to 100; or 51.19 and 48.81 in each 100 born.
1871...........956 and 1,004; or 95.6 to 100; or 48.78 and 51.22 in each 100 born.
1872.......1,079 and 1,127; or 95.7 to 100; or 48.91 and 51.09 in each 100 born.
1873.......1,109 and 1,019; or 108.8 to 100; or 52.11 and 47.89 in each 100 born.
1874......1,408 and 1,372; or 102.6 to 100; or 50.65 and 49.35 in each 100 born.
1875.......1,390 and 7,274; or 109.1 to 100; or 52.18 and 47.82 in each 100 born.
1876..........1,359 and 1,272; or 106.8 to 100; or 51.65 and 48.35 in each 100 born.
1877.......1,302 and 1,241; or 104.9 to 100; or 51.20 and 48.80 in each 100 born.
```

24 years.....22,713 and 21,529; or 105.5 to 100; or 51.34 and 48.66 in each 100 born.

Color. The number of colored children born in Providence in 1877 was 104; this was four less than in 1876, and three more than in 1875; 60 were males, and 44, females, a proportion 136.4 males to each 100 females; or 57.69 males and 42.31 females in each 100 children born; the number of colored male children is generally larger than that of females; but this proportion is much larger than usual. In the 22 years, 1856-1877 inclusive, the number of colored children born in the city was 1,595 (males, 849; females, 746); giving a proportion of 113.8 males to each 100 females; or 53.23 males and 46.77 females in each 100 children born. In 1877, the births were as follows in the different wards: first ward, 14; second, 12; third, 10; fourth 2; sixth, 19; seventh, 6; eighth, 12; ninth, 29; fifth and tenth, none. As to season, the numbers in the different months were: 15 in February; 13 in October; 11 in June; 10 each in March and August;

8 each in January, April, and May; 6 each in July and September; 5 in November; 4 in December.

The statistics of births, marriages, and deaths among the colored population of Providence, for the 22 years, 1856–1877 inclusive, are as follows:—

Twenty-one years, 1856-18761,491	births728 marriages1,576 deaths.
1877 104	births 46 marriages 105 deaths.
Twenty-two years' total1,595	births774 marriages1,681 deaths.

In 1877, the proportion of births to deaths was 100 to 100.9; for the twenty-two years, 1856-1877, the proportion was 100 to 105.4.

PARENTAGE.

The facts in regard to this subject are of considerable importance and interest, as showing how many different nations are represented in our city, and how these different nationalities are mingling with each other and with the native population.

Table III shows how many children, according to parentage, were born in each ward; also the number and percentage of each class in the whole city. Where the parents were of different foreign countries, the parentage is given according to the birth-place of the father:—

TABLE III. BIRTHS, 1877. Wards and Parentage.

	WARDS.									City.	929	
PARENTAGE.	L	II.	III.	IV.	v.	VI.	VII.	AIII	ıx.	X.	Whole	In each 100 born there were
American	140	49	84	54	72	81	96	139	229	110	1,054	41.45
Irish. English and Scotch. German. British American. Portuguese. Other Foreign.	101 21 7 16		73 7 1 4 28 4	27 3 1 5	9 9	5 1 	3 1	52 17 2 13	117 36 9 2 3 7	197 63 16 16 1 3	714 177 50 65 28	28.08 6.96 1.96 2.56 1.10 1.30
Total Foreign	30		13	6	14	34 7	68 12 24			296 48 51	1,067 184 238	41.96 7.23 9.36
Ward totals	368			_		126	200					100.00

The "other foreign" in this table were as follows: of Swedish parentage, 9; Italian, 8; Polish, 5; Danish, French, and Russian, 3 each; Mexican and Norwegian, 1 each.

The following statement shows the number and percentage of children of American, foreign, and mixed parentage, born in Providence in each of the twenty-two years, 1856–1877 inclusive; and also for the whole period:—

Year.	American. Number, Per cent.	Foreign. Number, Per cent.	Amer. father, For, mother. Number. Per cent.	For. father, Amer. mother. Number. Per ct.
1856	MON 10.00	857 51.17.	553.22	. 583.52
1857	. 65038.51	937 55.51.	513.02	. 502.96
1858	. 71241.30	886 51.39.	583.36	. 683.95
1859	. 59937.60	., 887 55.69.	593.70	. 483.01
1860	. 66140.11	857 52.00.	694.19	. 613.70
1861	. 65437.91	916 53.10.	864.99	. 694.00
1862	. 58938.52	799 52,26.	744.84	. 674.38
1863	. 46735.73	711 54.40.	624.74	. 675.13
1864	52038.69	699 52.00.	604.47	. 654.84
1865	. 47636.67	684 52.70.	705.39	. 685.24
1866	. 65640.20	821 50.30.	704.29	. 855.21
1867	. 64539.72	791 48.70.	804.93	.1086.65
1868	77941.75	868 46.51.	1035.52	.1166.22
1869	82245,19	776 42.16.	1045.72	.1176.43
1870	80042.35	862 45.64.	904.76	.137 7.25
1871	87944.85	828 42.24.	1015.15	.1527.76
1872	90741.12	1,002 45.42.	1285.80	.1697.66
1873	86940,84	941 44.22.	1346.29	.1848.65
1874	1,06838.42	1,326 47.70.	1495.36	.237 8.52
1875	99937.50	1,256 47.15.	1816.79	.2288.56
1876	1,00738.27	1,230 46.75.	162 6.16	.2328.82
1877	1,05441.45	1,067 41.96.	1847.23	.238 9.36
22 years' totals	16,518 40.02	20,001 48.46	2,130 5.16	2,624 6.36

The percentages in the different classes in 1877 are quite different from those of the preceding year; the percentage of "American" is larger than it has been since 1871; while the percentage of "foreign" is the smallest that has ever been recorded, and is far below the average for the whole period. On the other hand, in both classes of mixed parentage, the percentage for 1877 is the largest on record, and considerably above the average for the whole period. As usual, there is a larger percentage of children with foreign father and American mother, than of the opposite class.

The following table shows the percentage of children, born in Providence, of parents of each nation, for each of the twenty-two years, 1856-1877:—

	American. Per cent.	Irish. Per cent.	English, 8. a W. Per cent.	German. Per cent.	British America. Per cent.	Portuguese. Per cent.	Other for'n. Per cent.
1856	42.09	41.79	4.84	2.27			2.27
1857	38.51	44.32	5.92	3.73			1.54
1858	41.30	42.98	4.82	2.26			1.33
1859	37.60	45.58	5.34	3.26			1.51
1860	40.11	40.96	6.55	2.85		· · · · - · · · · ·	1.64
1861	37.91	42.61	6.03	2.49			1.97
1862	38.52	41.53	6.02	3.07			1.64
1863	35.73	43.08	6.35	2.98			1.99
1864	38.69	41.36	6.55	2.75			1.34
1865	36.67	42.99	5.93	1.85			1.93
1866	40.20	39.52	6.25	2.21			2.32
1867	39.72	37.81	5.66	3.02			2.21
1868	41.75	36.01	6.48	1.88	1.07		1.07
1869	45.19	34.14	4.01	1.81	1.32		1.38
1870	42.35	34.68	6.19	1.96	1.91		
1871	44.85	32.25	4.54	1.63	2.09		1.73
1872	41.12	32.46	5.67	2.77	2.31		2.31
1873	40.84	30.50	7.19	1.88	2.49		2.16
1874	38.42	32.63	8.13	2.34	2.19	1.22	1.19
1875	37.50	31.53	7.69	2.85	2.67	98	1.43
1876	38.27	31.05	7.75	2.59	2.89	1.22	1.25
1877	41.45	28.08	6.96	1.96	2.56	1.10	1.30

The British American were included in the "other foreign" until the year 1868; and the Portuguese until 1874.

The parentage of the children born during the year 1877 is more exactly shown in Table IV, which gives the birthplaces of both parents:—

TABLE IV. BIRTHS, 1877. Parentage.

	BIRTHPLACES OF MOTHERS.									
EIRTHPLACES OF FATHERS.	United States.	Ireland.	England, Scotland, and Wales.	Germany.	British America.	Portugal and Western Islanda.	Other Countries.	Whole number according to the father's birthplace.		
United States Ireland. England, Scotland and Wales. Germany British America. Portugal and Western Islands. Other Countries.	1,054 140 45 18 23 2 10	109 666 35 8 7 3 5	44 30 124 8 	2 36 	29 18 12 	 25	2 6 1 25	1,238 854 222 68 88 30 43		
Whole number, according to the mother's birthplace	1.292	833	208	39	108	25	38	2,543		

This table, like the preceding one, shows that the number of children of American parentage in 1877, was larger than usual. In addition to the 1,054 whose parents were both natives of the United States, there were 184 who had American fathers and foreign mothers, and 238 who had foreign fathers and American mothers; making in all 1,238 with American fathers, and 1,292 with American mothers. Of the children whose parents were natives of the same foreign country, the proportion was less than usual, 925, or 36.38 per cent., being of this class; of those whose parents were both foreign, but of different countries, there were 142, or 5.58 per cent. of all; of purely American parentage, 1,054, or 41.45 per cent.; of those whose parents were one American and one foreign, 422, or 16.59 per cent.

NUMBER OF THE CHILD.

The following table shows how many of the 2,543 children born in Providence in 1877 were the first, second, third, etc., children of their mothers; also the same facts for the 40,330 children born during the twenty-two years preceding, 1856-1876 inclusive:—

TABLE V. BIRTHS,	1877.	Number	of	the	Child.
------------------	-------	--------	----	-----	--------

NUMB	EE	0	F CI	HILD.	1877.	22 years, 1855-1876.	NUM	BER	OF	CHI	LD.	1877.	22 years, 1855-1876.
First chi					557	10,177	Twelfth						153
Decount	**		4.6	**	548	8,351	Thirteen			**	16	6	58
Third	44	**	- 46	46	443	6,377	Fourteen			44	**	1	27
Fourth	**	16	44	**	314	4,765	Fifteent!	1 11		**	44	1	13
Fifth		44	*6	46	236	3,480	Sixteent/	2 60	44	16	44		9
Sixth	16	44	48.	44	156	2,521	Seventeer	nth "	46	11	64	1	7
	15	*	44	**	117	1.777	Eighteen	th	**	46	66		
	11	**	40	44	60	1.125	Nineteen		4.6	66	44		4
	46	44	44	46	40	751	1	****					-
	44	24	40	44.	36	481	Whole n	umbe	ar c	f ch	ildren.	2,543	40,330
Eleventh		66.		**	13	262				- 011		2,010	20,000

Of the 2,543 children born in Providence in the year 1877, 557, or 21.90 per cent., were the first children of their mothers. Of the 40,330 children born during the twenty-two years preceding, 10,177, or 25.23 per cent., were the first children their mothers had borne.

There were 27 twin births in Providence in 1877, so that the number of mothers was 2,516. These 2,516 mothers had borne in all 8,705 children, an average of 3.45 children to each mother.

The statistics upon this subject are given in Table VI, for each of the twenty-three years, 1855-1877 inclusive.

This table shows how many children these mothers had already borne, and the average number for each; but it does not show the average number of children the same mothers may have in a life-time.

This may be ascertained approximately by dividing the number of children born in a long series of years, by the number of mothers bearing their first children during that period. Thus, the whole number of children born in Providence during the twenty-three years, 1855–1877 inclusive, was 42,873, and the number of mothers bearing their first children during that time was 10,734. This would give 3.99 as the average number of children each mother would bear in a life-time.

Table VI. Births, 1855 to 1877.

Number of child to each mother.

Years.	No. of	No. of Children	Aggregate No. of children.	Average	
				3.28 children to	each mother.
				3.35 children to	
				3.44 children to	
				3.33 children to	
			•	3.58 children to	
	•	•	•	3.59 children to	
			•	3.55 children to	
	•		•	3.52 children to	
	•	•	•	3.64 children to	
			•	3.68 children to	
	-	•	•	3.60 children to	
				3.56 children to	
		•	•	3.50 children to	
	•	•	,	3.44 children to	
			•	3.30 children to	
	•	•	•	3.31 children to	
			•	3.31 children to	
	•		•	3.38 children to	
				3.20 children to	
			•	3.27 children to	
				3.41 children to	
				3.42 children to	
				3.46 children to	
23 years	42,334	42,873	144,955	3.42 children to	each mother.

Table VII gives the number of the child in connection with the age of the mother, and shows how many first, second, etc., children were born in 1877, to mothers at the respective ages given; also the aggregate and average number of children the mothers at each age had borne:—

Table VII. Births, 1877. Number of child, with age of mother.

16.	AGE OF THE					NU	MBE	R OI	C	нп	D.							mothers age.	st7.	dren they borne.	number of to mothers ch age.
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	MOTHERS IN YEARS.	1st,	2d.	3d.	4th.	3th.	6th.	7th.	8th.	9th.	10th.	11th.	12th.	13th.	14th.	15th,	17th.	Number of mothers at each age.	Children th	Total child	Average nu children to at each
46	17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 44.	111 222 399 533 477 644 466 500 288 200 183 39 66 77 22 23 31 11	4 9 17 33 *40 151 54 39 *42 25 24 19 18 8 11 11 11 22 11	6 4 13 20 26 48 33 40 40 *34 40 *34 18 11 8 *7 6 6 †55	4 10 21 20 24 32 27 29 24 *25 15 14 *16 4 3 2 2 11	27 6 21 24 15 *19 *20 *23 13 14 9 66 11 12 21 11 11	5 5 7 11 16 16 16 18 11 10 10 7 9 9 4 5 2 2 11 1	22 33 38 22 5 100 121 100 88 100 9 177 9 8 8 5 5	1 2 4 5 5 7 7 7 4 5 5 5 5 3 2 1 1	222345534421		1112211	332221					66 12 26 68 68 68 68 68 68 68 68 68 68 68 68 68	66 12 26 36 37 37 28 38 39 108 135 153 153 153 153 153 153 153 153 153	13 30 49 491 128 256 294 431 510 542 543 558 363 449 449 441 449 449 441 449 449	1.000 1.151 1.481 1.471 1.000 1.152 1.000

^{*} One of these mothers bore twins.

The reference marks in this table indicate the age of the mother and the number of the first child in the cases of plurality birth. Thus, at the age of 38 years, three mothers bore twins; in one instance, they were the third and fourth children; and in two instances, they were the seventh and eighth children.

The oldest women bearing children in Providence in 1877, were

[†]Two of these mothers bore twins.

two women of 47 years, each of whom had borne seven children before. There were 111 women in Providence, over 40 years old, who bore children in 1877. They had been the mothers of 795 children in all, or an average of 7.16 children to each mother. In the fourteen years and six months from July 1st, 1863, to December 31st, 1877, there were 1,435 women in Providence, of the age of 40 years or more, who bore children. They had borne in all 11,039 children, or an average of 7.69 each.

This record of the age of the mothers has been kept for fourteen years and six months, from July 1st, 1863, to December 31st, 1877, during which time 28,953 children have been borne by mothers whose age was stated. The following table shows the number of children borne by mothers of each age during this period:—

Age of Mother.	Children.	Age of Mother.	Children.	Age of Mother.	Children.
14	2	27	1,684	40	672
15	18	28	2,020	41	191
16	46	29	1,457	42	225
17	129	30	2,327	43	128
18	325	31	986	44	84
19	529	32	1,321	45	89
20	857	33	1,095	46	23
21	992	34	929	47	23
22	1,421	35	1,455	48	7
23	1,579	36	858	49	3
24	1,747	37	706	50	4
25	•	38	777		
26	•	39	483	All, known ag	es28.953

The mothers of these children were divided as to age as follows:—

Under twenty years	1,049
Twenty, and under twenty-five	6,596
Twenty-five, and under thirty	8,922
Thirty, and under thirty-five	6,658
Thirty-five, and under forty	4,279
Forty, and under forty-five	1,300
Forty-five and over	149
m-a-1	00.082

From this statement it may be seen that of the 28,953 children born during this period, 16,567, or 57.22 per cent. of all, were borne by mothers less than thirty years of age; 10,937, or 37.78 per cent., by mothers between thirty and forty; and 1,449, or 5.00 per cent., by mothers of forty years of age and over.

PLURALITY BIRTHS.

The cases of plurality births in Providence in 1877 numbered 27, the same number that occurred in 1876; all were twin births.

The following particulars are given concerning them :-

Cases. Whole number, 27; children born, 54.

Locality. First ward, 7; second, none; third, 3; fourth, 1; fifth, 3; sixth, 2; seventh, 1; eighth, 1; ninth, 7; tenth, 2.

Season. In January, there was one case; in February, 5; in March, 1; in April, 1; in May, 2; in June, 1; in July, none; in August, 2; in September, 2; in October, 5; in November, 3; in December, 4.

Sex. In nine cases both children were males; in nine, both were females; and in nine, there was one male and one female child. Thus the children were equally divided, as to sex: males, 27; females, 27.

Color. The parents of the children were white in every case but one.

Number of the child and age of the mother. In five cases, the twins were the first and second children of mothers, two of whom were 21 years old; one, 19; one 29, and one, 42; in five cases, they were the second and third children of mothers, two of whom were 23 years old; one, 22; one, 27, and one, 28; in six cases, they were the third and fourth children of mothers, 27, 29, 32, 38, and two of forty years; in two cases, they were the fourth and fifth children of mothers 32 and 35 years old; in three cases, the fifth and sixth children of mothers of 30, 31, and 36 years; in three cases, the seventh and eighth children of mothers, one of whom was 36 years old, and two were 38; in one case, the eighth and ninth children of a mother 36 years old; and in two cases, the ninth and tenth children of mothers 37 and 45 years old. In sixteen of the twenty-seven cases, the mothers were thirty years old or over.

Parentage. In twelve cases, both parents were American; in nine, both were Irish; in one, both were Portuguese; in two cases the father was Irish, and the mother, American; in two, the father English, and the mother, Irish; and in one, the father English, and the mother, British American. There were, therefore, fourteen American mothers; eleven Irish; one Portuguese, and one British American.

TWENTY-THREE YEARS' SUMMARY OF PLURALITY BIRTHS.

In the twenty-three years, 1855-1877 inclusive, there were 530 cases of plurality births in Providence; 521 cases of twins; and 9, of triplets; whole number of children, 1,069.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children, and one female; making 9 male and 18 female children in the nine cases.

Of the 521 cases of twin births, in 164 cases, both children were males; in 179 cases, both were females; in 178 cases, there was one male and one female child; thus of the 1,042 children, 506 were males, and 536 females. Of the 1,069 children in all the cases of plurality births, there were 515 males, and 554 females.

Color. The parents of all the triplets were white. Of the 521 cases of twins, in 503 cases the parents were white, and in 18 cases, colored.

Parentage. Of the triplets, in one case, the parents were both American; in five cases, both were Irish; in one case, both German; in one, the father was English and the mother Irish; and in one, the father Scotch and the mother Irish. There was but one American mother in the nine cases.

The following statistics of the 521 cases of twin births, give the nativity of both parents, the first letter representing the father's birthplace and the second the mother's.

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British American; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish:—

Parentage.	No.	Parentage.	No.	Parentage.	No.	Parentage.	No.
A. and A	174	I. and A	13	G. and I	1	I. and S	3
I. and I	204	E. and A	12	B. A. and I.	1	E. and S	1
						I. and W	
S. and S	4	B. A. and A.	6	It. and I	1	A. and B. A.	4
G. and G	13	G. and A	1	P. and I	1	I. and B. A.	1
B. A. and B.	A. 9	Sw. and A	1	A. and E	$\dots ar{3}$	E. and B A	1
						S. and B. A.	
P. and P	5	E. and I	11	S. and E	1		
		S. and I					521

From this statement it may be seen that in 209 of these cases of twin births, the mothers were natives of the United States; in 234 cases, the mothers were natives of Ireland; in 33 cases, of

England, and so on. The whole number of foreign mothers was 312. In the 530 cases of plurality births in the twenty-three years, the number of American mothers was 210; foreign mothers, 320.

Proportion of plurality births to whole number. The following numbers show how the year 1877 compares with the twenty-two years previous, 1855-1876 inclusive, as to the total number of births, and the proportion of plurality births to the whole number:—

Single births.				Whole No. Children
22 years, 1855–187639,315			•	•
18772,489	27	••••	2,516	2,543
23 years' total41,804	521	9	42,334	42,873
Cases of plurality bir	th s : –	_		
	Twins			Triplets.
1855–1876	.1 in 80	.6		1 in 4,424.2
1877	.1 in 93	3.2		•••
1855-1877	.1 in 81	3		1 in 4,703.7
Children of plurality	births	:		
	Twins			Triplets.
1855-1876	1 in 40	.8		1 in 1,493.7
1877	.1 in 47	.1	· · · · · · · · · · · · · · · · · · ·	•••
		_		

Proportion of plurality births by parentage. In the twenty-three years, 1855-1877 inclusive, there were 42,334 births. The following statement shows the number of births of American, foreign, and mixed parentage; and the number and proportion of plurality births in each class:—

	Whole No. Births.	Plurality Cases. 1 in Per cent.
American parents	16,981	175, or 97.03, or 1.03
Foreign parents		
Mixed, American and foreign	4,808	53, or 90.72, or 1.10
Total	42,334	530, or 79.88, or 1.25

Of the whole number of plurality births in the twenty-three years, the 175 cases of American parentage were 33.02 per cent.; the 302 cases of foreign parentage were 56.98 per cent.; and the 53 cases of mixed parentage were 10.00 per cent.

MARRIAGES.

The number of marriages in Providence in the year 1877, was 953.

Table VIII shows the number in each month, each quarter, and the whole year 1877, and in each of the six years preceding, 1871-1876 inclusive; also the aggregate and average number for the corresponding seasons, in the period of twenty-three years, 1855-1877 inclusive:—

TABLE VIII. MARRIAGES. Season.

=======================================				===	===	 .	·		
MONTHS.	1877.	1876.	1875.	1874.	1873.	1872.	1871.	23 years. 1855-1877.	23 years average
January February March	68 65 38	79 93 44	75 79 71	93 87 56	105 111 49	97 83 53	75 75 47	1,623 1,453 937	71 63 41
First quarter	171	216	225	236	265	233	197	4,013	175
April May June	83 85 83	70 80 94	84 82 100	95 113 101	87 91 107	93 85 94	72 78 94	1,473 1,618 1,658	64 70 72
Second quarter.	251	244	266	309	285	272	244	4,749	206
July	85 76 87	56 62 93	101 73 111	88 80 129	105 63 108	91 71 83	74 65 73	1,425 1,219 1,710	62 53 74
Third quarter	248	211	285	297	276	245	212	4,354	189
October November December	99 102 82	95 109 74	107 114 82	127 114 93	118 125 81	124 121 89	99 120 71	1,919 2,047 1,518	83 89 66
Fourth quarter.	283	278	303	334	324	334	290	5,484	238
Yearly totals	953	949	1,079	1,176	1,150	1,084	943	18,600	808

The number of marriages in 1877 was four more than in 1876, but still much smaller than in the four years previous to that. Having estimated the population of the city in the middle of the year 1877, at 100,675, we find the ratio of persons married to the population to be one in 52.82, or 18.93 persons to each thousand

inhabitants. The ratio of persons married to children born, was 75 to 100; of persons married, to deaths, 98 to 100.

Season. According to the general rule, the majority of the marriages were in the last six months of the year; the largest quarterly number being in the fourth quarter. The largest number in one month was 102 in November; the smallest, 38, in March. The number in the first quarter is always the smallest, but for this year it was even smaller than usual. From the aggregate number of marriages as given in Table VIII for the 23 years, 1855–1877, we find that of each 100 marriages, about 25 are in the winter months, December, January, and February; 22 in the spring, 23 in the summer, and 30 in the autumn.

The number and percentage of marriages in each quarter of the year 1877, and of the aggregate of marriages in the 22 years previous, 1855-1876, were as follows:—

	1877		1858 Marriages.	5 1876.
	Marriages.	Per cent.	Marriages.	Per cent.
January-March	171	17.94	3,842	21.77
April-June		26.34	4,498	25.49
July-September	248	26.02	4,106	23.27
October-December	283	29.70	5,201	29.47
Total		100.00	17.647	100.00

Nativity. The following table gives the birthplaces of both parties, in each of the 953 marriages in the year 1877: --

TABLE IX. MARRIAGES, 1877. Nativity of the parties.

	BIRTHPLACE OF BRIDES.								
Birthplace of Grooms.	United States.	Ireland.	England, Scot- land, and Wales.	Germany.	British America.	Portugal and Western Islands.	Other countries.	Whole number of Grooms.	
United States	545 39 26 5 19 1	28 149 7 2	21 3 27 1 1	1 12 	24 3 4 6		3 1 6	621 194 65 17 26 12	
Whole number of brides	644	186	54	13	37	9	10	953	

Bride and Groom born in the same Country. In 754 of these marriages, or 79.12 per cent. of the whole number, both parties were natives of the same country. In 545, or 57.19 per cent., the bride and groom were both born in the United States; in 149, or 15.63 per cent., both were natives of Ireland; in 27, both were natives of England, Scotland, or Wales; in 12, both of Germany; in 6, both of British America; in 9, both of Portugal or the Western Islands; in 6, both parties were of other foreign countries than those named in the table.

Bride and Groom born in different Countries. In 199 marriages, or 20.88 per cent. of all, the bride and groom were natives of different countries; 39 men born in Ireland married women who were natives of the United States; and 28 men born in the United States married women born in Ireland. Of the 621 grooms born in the United States, 76, or 12.24 per cent., married women of different foreign countries; of the 194 grooms born in Ireland, 45, or 23.20 per cent., married natives of other countries. Of the brides, 26 natives of the United States married men born in England, Scotland, or Wales; 24 natives of British America married men born in the United States, etc. Of the 644 brides, who were natives of the United States, 99, or 15.37 per cent., married men of other countries; of the 37 born in British America, 31, or 83.78 per cent., married men born in other countries.

Of the 18 grooms born in countries not named in the table, 7 were natives of France; 4, of Sweden; 3, of Italy; and one each of Cuba, Russia, Poland, and St. Helena. Of the 10 brides of other countries, 2 each were natives of France, Poland, and Sweden; and one each of Hungary, Italy, Switzerland, and the East Indies.

Of the 953 grooms, 621, or 65.16 per cent., were born in the United States; this number is six more than the number of this class in 1877; of the 953 brides, 644, or 67.58 per cent., were natives of the United States; 4 more than in 1876. The number of grooms, born in foreign countries, was 332; two less than in 1876; the number of brides 309, which is just the same as in the preceding year.

The number and percentage of American, foreign, and mixed marriages in Providence, are shown in the following table for each of the 23 years, 1855–1877 inclusive. By mixed marriages are understood those where one of the parties was of American, and one of foreign birth:—

	Amer	ican.	For	eign.	Mix	
Year.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent
1855	308	48.73	265	41.93	59	9.3
1856	352	54.74	212	32.97	79	12.2
1857	384	62.54	175	28.50	55	8.9
1858	283	50.54	214	38.21	63	11.2
1859	333	54.50	224	36.66	54	8.84
1860	336	54.19	225	36.29	59	9.55
1861	304	55.58	182	33.27	61	11.13
1862	288	52.65	179	32.72	80	14.6
1863	342	56.07	192	31.47	76	12.40
1864	427	57.78	211	28.55	101	13.6
1865	417	58.82	202	28.49	90	12.6
1866	471	58.00	244	30.05	97	11.9
1867	485	56.73	269	31.46	101	11.8
1868	530	60.78	215	24.66	127	14.5
1869	530	56.99	286	30.75	114	12.2
1870	552	57.56	262	27.32	145	15.1
1871	523	55.46	278	29.48	142	15.0
1872	591	54.52	305:	28.14	188	17.3
873	652	56.69	317	27.57	181	15.7
874		55.10		29.42		15.4
875		54.31				16.0
876	200000000000000000000000000000000000000	57.64		25.39	SECTION AND ADDRESS.	16.9
1877		COCCERCE CO.		24.45		

The proportion of American marriages is slightly less than in 1876, but greater than for several years previous to that; the proportion of foreign marriages is the smallest recorded in the whole period, and the proportion of mixed marriages considerably larger than in any previous year. The apparent discrepancy in the number of marriages, as given here, and the number for the same period of years, as given in Table VIII, is owing to omissions in the early years of the records, and is explained in the report for 1875.

Age of persons married. Table X, on the next page, shows the number and percentage of brides and grooms in each division of ages.

Of the 953 grooms included in the table, 317, or 33.26 per cent., were less than twenty-five years old; of the brides, 556, or 58.34 per cent. The aggregate and average age of those married in Providence in 1877, were as follows:—

953 malesAggregate age	27,904 years Average as	ze, 29,28
953 femalesAggregate age		

TABLE X. MARRIAGES, 1877. Ages of the Parties.

		Ac	es of	тнв В	RID E S.			number ooms.	Percentage of Grooms.
Ages of the Grooms	Under 20 years.	30 to 25.	25 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	Whole numb of Grooms	
Under 20 years			32 100 69 10 4	2 23 65 29 11 1 1	1 7 12 11 6 1 1	5 3 1	2	21 296 320 208 61 31 12 3	2.20 31.06 33.58 21.83 6.40 3.25 1.26 .31
Whole number of Brides	158	398	215	133	38	9	2	953	
Percentage of Brides	16.58	41.76	22.56	13.96	3.99	.94	.21		100.00

The following table shows the aggregate and average age of the persons married in Providence, in each of the twenty-two years from 1856 to 1877 inclusive:—

	Males.	Females.
Years,	No. Aggregate age. Average age.	No. Aggregate age. Average ag
1856	639 18,104 years. 28.33 years	637 15,559 years. 24.42 years
1857	607 16,753 years27.59 years	606 14,427 years23.84 years
1858	560 15,850 years 28.29 years	560 13,702 years24.47 years
1859	611 16,989 years27.81 years	611 14,616 years23.92 years
1860	620 17,135 years27.60 years	620 14,667 years 23.66 years
1861	547 15,448 years28.24 years	547 13,316 years22.33 year
1862	547 15,773 years28.83 years	547 13,565 years24.80 years
1863	610 17,208 years28.21 years	610 14,792 years 24.24 years
1864	739 21,361 years28.90 years	739 18,075 years24.45 years
1865	709 20,539 years28.96 years	709 17,605 years 24.83 years
1866	812 22,801 years28.08 years	812 20,036 years . 24.67 year
1867	855 24,443 years 28.58 years	855 21,078 years24.65 year
1868	872 24,546 years28.15 years	872 21,440 years24.59 year
1869	930. 26,548 years . 28.55 years	930 23,135 years24.88 year
1870	959 27,343 years28.51 years	959 23,773 years24.79 year
1871	943 26,743 years28.37 years	943 23,290 years24.70 year
1872		1,084 26,835 years 24.76 year
		1.150 28,701 years 24.96 year
		1,176 29,510 years 25.09 year
	1,079 30,848 years28.59 years	
	949 27,496 years28.97 years	
	953 27,904 years29.28 years	. •

In 1856, the ages of 17 males and 19 females, and in 1857, the ages of 7 males and 8 females, were not given. This will account for the apparent discrepancy in the first lines of this table.

Marriages of colored persons. The number of marriages of colored persons in this city in 1877, was 46; 8 more than in 1876. The aggregate and average age of the parties in these marriages were as follows:—

46 males	Aggregate age, 1	337 yearsAverage	age, 29.06
		,143 yearsAverage	

The number of marriages of colored persons, during the twenty-two years, 1856-1877 inclusive, was 774; the aggregate and average age, as follows:—

774 males Aggregate age, 22,719 years Averag	e age, 29.35
774 females Aggregate age, 20,370 years Averag	

Number of the marriage. The following table shows how many of those married in Providence in 1877, were married for the first, second, third, or fourth time:—

TABLE XI. MARRIAGES, 1877. Number of times married.

	BRIDES.	Number of the	Marriage.	Whole
GROOMS. Number of the Marriage.	First.	Second.	Third.	number of Grooms.
First	703	63	1	767
Second	88	80	4	172
Third	5	1 6	1	12
Fourth	1		1	2
Whole number of Brides	797	149	7	953

In 703 instances, or 73.77 per cent. of all, it was the first marriage of both parties; in 80 instances, it was the second marriage of both. One marriage was the fourth of the groom, and the first of the bride.

Denominational. The number of marriages performed in Providence in 1877, by the clergymen of the different religious denominations, was as follows:—

	· · · · · · · · · · · · · · · · · · ·
Roman Catholic305	Bethel Pastor 10
Calvinistic Baptist	Lutheran
Free Will Baptist 30	Advent Christian 6
Methodist	Evangelical Advent
Episcopalian	New Jerusalem 3
Congregational 58	Latter Day Saints 2
Unitarian 42	Jewish 2
Universalist 47	_
Presbyterian 10	Total953
Christian 19	

Nativity and Residence. There were 953 marriages in Providence in 1877; in only 61 were the bride and groom both born in the city; these 122 persons were 6.40 per cent. of the whole number of persons married. In 147 more marriages, the groom only was born in Providence; and in 101 others, the bride only. In all there were 370 of the persons married, who were natives of the city; this is 19.41 per cent. of the whole number. In 104 marriages, or 10.91 per cent. of all, both parties were non-residents.

Marriage and Education. The number of persons married in Providence in 1877, was 1,906; they were divided, as to sex and nativity, as follows:—

Born in United	States	Males,	621	Females,	644	.Total,	1,265
Born in foreign c	ountries	Males.	332	Females,	309	.Total,	641

In each class, the number and proportion who signed the marriage certificate with a mark, were as follows:—

1877.	married.	Signed with a mark.	100.
Males, born in United States	621		4.35
Females, born in United States	644		5.75
		_	
Total, born in United States	1,265	64	5.06
Males, born in foreign countries	332	84	25.30
Females, born in foreign countries	309	105	
		100	~~
Total, born in foreign countries	641	189	29.49

Of the 64 born in the United States, who signed with a mark, 33 (18 males and 15 females) were colored persons, and 27 more (8 males and 19 females) were of foreign parentage. This leaves only 4 persons (one male and 3 females), of white American parentage, who signed the marriage certificate with a mark.

The following are the statistics on this subject for the eighteen years, 1859–1876 inclusive:—

18591876.	Whole number married.	Signed with a mark.	In each 100.
Males, born in United States	9,536	480	5.03
Females, born in United States	9,720	676	6.95
Total, born in United States	19,256	1,156	6.00
Males, born in foreign countries	. 5,656	1,950	34.48
Females, born in foreign countries	5,472	2,799	51.15
			_
Total, born in foreign countries	11,128	4,749	42.68

These statistics vary slightly from year to year; thus, in 1875, the number of persons married was 2,158; the proportion who signed the certificate with a mark, 18.44 in each hundred; in 1876, the number married, 1,898; the proportion who signed with a mark, 13.07; in 1877, the number married, 1,906; the proportion who signed with a mark, 13.27 in each hundred.

The statistics for 1877 show a slight increase over those for 1876, both in numbers and proportions, of those who signed with a mark in both the classes mentioned; but of the persons of American nativity a much smaller proportion were of white American parentage; the number in 1877 being just one fourth of the number in the preceding year.

DEATHS.

There were 1,938 deaths in Providence during the year 1877; this was 73 more than in 1876, and 23 more than in 1875. The number of males in 1877 was 40 more than in 1876; of females, 33 more; of whites, 78 more; of colored, 5 less; of natives of the United States, 69 more; of natives of foreign countries, 4 more; of American parentage, 33 more; of foreign parentage, 40 more.

The average death-rate in Providence for the twenty-three years, 1855–1877 inclusive, was one in 50.82 of the population. In 1877, it was as follows among the different classes of the people:—

	Population, 1875.		Deaths to To population each 1.000 one death in population
Whole number	100,675		
Males	48,701	922	52.8218.9
Females			
Whites	•		
Colored	•		
Born in the United States	71,438	1,511	47.2821.1
Born in Ireland	18,458	292	63.2115.8
Born in England, Scotland, and Wale	s 6,110	83	73.6113.5
Born in Germany	1,246	18	69.2214.4
Born in other foreign countries	3,423	34	100.67 9.9
Total of foreign born	29,237	427	68.4714.6
American parentage	47,316	908	52.1119.1
Irish parentage	37,440	719	52.0719.2
English, Scotch, and Welsh parentage	9,636	177	54.4418.3
German parentage	2,516	44	57.1817.4
Other foreign parentage	. 3,767	90	41.8623.8
Total of foreign parentage	53,359	1,030	51.8019.3

The following table gives some general facts concerning the deaths in 1877: —

TABLE XII. DEATHS, 1877.

Sex, condition, color, nativity, parentage, locality, season.

				<u> </u>	. —				_:=	=_			
1877.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Whole year.
Whole Number of Deaths	153	117	141	157	160	123	189	180	188	173	175	182	1,938
Ser. — Males Females	68 85		71 70	75 82	77 83		109 80			1 75 98		70 112	922 1,016
Condition.— Married	17			45 90 14 8	85 20	6	54 120 10 5	15	116 11 6	115 10 7	101	21 5	565 1,135 164 68 6
Color. — Whites	144 9	113 4	133 8	144 13	150 10	116 7	178 11	161 19	184 4		169 6	175 7	1,833 105
Nativity. — United States	123 19 8 	14 3 2 2	101 28 9 2 1	26		16 9	150 31 4 3 1	24 7 2	29 4 2 1	24 9 1 1	22 7 4 4	34 6 2 6	1,511 292 83 18 22 3 9
Parentage. — American Irish English and Scotch German: British American Portuguese Other foreign.	79 50 15 1 6 1	58 39 11 3 3	60 52 17 6 3	71 68 11 2 1 2 2	1	1 3 1	84 65 19 7 10 3	90 60 18 6 4 2	78 84 12 4 2 3 5	80 70 11 5 2	90 57 16 5 6	9	908 719 177 44 52 16 22
Locality, Wards.— I	16 7 14 9 17 12 19 12 20 20	21 10 12 10 5 6 11 17 17	20 4 15 6 16 7 6 18 18 22	25 2 7 10 16 12 18 14 18 33 2	25 11 10 10 7 13 6 17 18 35 8	6 10 4 12 8 10 9	27 4 33 11 16 11 13 15 27 24 8	19 9 14 9 14 12 19 14 25 34	32 5 19 8 16 16 18 17 19 30 8	25 3 26 9 20 10 14 9 23 30 4	23 16 11 12 9 18 18 24 34 6		243
Totals	153	117	141	157	160	123	189	180	188	173	175	182	1,938

Season. As usual the largest number of deaths was in the third quarter of the year; the smallest number, though very gen-

erally in the second quarter, was in 1877, as in the preceding year, in the first quarter. The number of deaths in each quarter of the year 1877, and of each of the five years preceding, is given below; also the quarterly aggregate and average numbers for the twenty-two years, 1855-1876 inclusive:—

	1877.	1876.	1875.	1874.	1873.	1872.	22 years 1855-1876. Aggr'te. Av'ge.
January-March		383					
April-June							
July-September	557	573	531	556.	504	518	8,276376
October-December	530	490	495	486.	381	391	6,648302
					_		
Whole year	1,938.	.1,865	.1,915.	.1,987.	1,719.	1,603.	27,5371,252

The number and percentage in each quarter of each year, 1855-1877, were as follows:—

	Janus	ry-March.		ril-June.		y-Sept.		-Dec.
_	eaths.			Per cent.				
1855	213	21.49	208	20.99	339	34.21	231	23.31
1856	.308	28.92	244	22.91	296	27.80	217	20.37
1857	214	23.14	173	18.70	286	30.92	252	27.24
1858	.279	27.43	278	27.34	257	25.27	203	19.90
1859	.213	23.69	205	22.80	270	30.04	211	23.4
1860	243	24.27	209	20.88	297	29.67	252	25.18
				22.45				
				21.01				•
				20:43				
				21.31				
				19.82				
				21.04				
				21.04				
				21.98				
				19.90				
				21.46				
				20.98				
1872	330	20.59	364	22.71	518	32.31	391	24.3
1873	427	24.84	407	23.68	504	29.32	381	22.1
1874	444	22.35	501	25.21	556	27.98	486	24.4
1875	488	25.48	401	20.94	531	27.73	495	25.8
1876	383	20.54	419	22.47	573	30.72	490	26.2
						28.74		

The number and percentage of deaths in each quarter of the year 1877, and in the aggregate for the twenty-two years, 1855—1876 inclusive, were as follows:— \

		1877.	23 years	, 1855–1876.
	Deaths.	Per cent.	Deaths.	Per cent.
January-March	. 411	21.21	6,567	23.85
April-June	. 440	22.70	6,046	21.96
July-September	557	28.74	8,276	30.05
October-December	. 530	27.35	6,648	24.14
	1,938	100.00	27,537	100.00

It is a well established fact that the highest percentage of mortality is in the third quarter of the year; and almost as general a rule that the lowest percentage is in the second quarter, though there are more exceptions to the latter, occasioned by the prevalence of epidemic diseases at certain seasons of some years. In 1856, scarlatina, croup, and pneumonia prevailed in the first and second quarters of the year, making the percentages at those seasons larger than usual. In 1858, scarlatina and measles prevailed, and caused the same deviation from the general rule. The large percentage in the second quarter of 1874, was caused by many deaths from scarlet fever, and from consumption, pneumonia, and congestion of the lungs. In the year 1877, the percentages in the different quarters of the year follow the general rule, that of the third quarter being the largest; it is, however, not so large as that of 1876; while in each of the other quarters the percentage is more than in the preceding year. The percentage of the last quarter in 1877 is larger than in any previous year, except 1862, owing to the large number of deaths from diphtheria.

Sex and Parentage. The decedents in 1877 were divided, as to sex and parentage, as follows:—

American parentage	410 males	498 females	Total, 908
Foreign parentage	512 males	518 females	Total, 1.030
Total	922 males	1,016 females	Total, 1,938

The number of female decedents was 94 more than of males; of this excess of females, 88 were of American parentage, and 6 were of foreign parentage. It has been quite a general rule that among the latter class, the males have exceeded the females; but in 1876 and in 1877, the reverse was true.

The decedents of both classes during the twenty-three years, 1855-1877 inclusive, were divided, as to sex, as follows:—

Condition. A comparison of the decedents in the various classes of civil condition with those of 1876, shows the following differences: married, 76 more than in the preceding year; single, 40 more; widows, 33 less; widowers, 7 less; divorced, 3 less.

Color. The number of deaths among the whites in 1877, was 78 more than in 1876; among the colored population, 5 less.

The number of births among the colored in 1877, was 104, or 4 less than in the year previous; proportion of births to deaths in 1877, 100 to 100.9.

In the twenty-three years, 1855-1877, there have been, among our colored population, 1,639 births and 1,735 deaths; a proportion of 105.8 deaths to each 100 births.

Nativity and Parentage. Table XII shows a great difference between the number of decedents registered according to nativity, and those registered according to parentage. It is an important distinction in these statistics, as very many of those who are born in the United States, are yet, as children of foreign parents, subject to the same social and hygienic conditions and influences as the class more strictly called foreign, because born in other countries. The number of decedents of American birth was 1,511, or 77.97 per cent. of the whole number; but the number of American parentage was only 908, or 46.85 per cent. of all; the number of foreign nativity was 427, or 22.03 per cent.; the number of foreign parentage, 1,030, or 53.15 per cent. Of the 9 born in other countries than those named in the table, 3 were born in France; 2, in Italy; 2, in Sweden; 1, in Greece, and 1, in the West Indies.

Of the 22 of "other foreign" parentage, 6 were of French parentage; 5, of Italian; 4, of Swedish; 2 each, of Swiss and Polish; and 1 each of Greek, African, and West Indian parentage.

Locality. Table XII gives the number of deaths in each ward, and in public institutions, in each month and in the whole year. The number in the first ward was 62 more than in 1876; in the second ward, 6 less; in the third, 11 less; in the fourth, 23 more; in the fifth, 3 more; in the sixth, 7 more; in the seventh, 28 more; in the eighth, 10 less; in the ninth, 68 less; in the tenth, 52 more; in public institutions, 7 less; summing up these numbers, we find that in the first, fourth, fifth, sixth, seventh, and tenth wards, collectively, the number of deaths was 175 more than in 1876; in the second, third, eighth, and ninth wards, and

in public institutions, collectively, the number was 102 less than in the preceding year.

The number of decedents in each ward, according to parentage, was as follows:—

Wards.	American.	Foreign.	Wards.	American.	Poreign
I	107	180	VII	84	77
II	55	12	VIII	118	54
ш	77	118	IX	133	110
IV	50	62	X	85	248
V	74	95	Public Instituti	ons 39	39
VI	86	35		-	
Who	le city			908	1,030

Average Age. In Table XIII are given the number of deaths, and the aggregate and average age of the decedents, according to parentage, for each month of the year 1877:—

Table XIII. Deaths, 1877.

Aggregate and Average Age, and Parentage.

	Amer	ican Pare	ntage.	Forei	gn Parer	itage.	Total American and Foreign.			
. 1877.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	
January February March April May June July August September October November December	79 58 60 71 68 68 84 90 78 80 90 82	3,319 2,065 2,112 2,681 2,428 2,120 1,994 2,355 2,242 1,907 2,710 2,420	42.01 35.60 35.20 37.76 35.71 31.18 23.74 26.17 28.74 23.84 30.11 29.51	74 59 81 86 92 55 105 90 110 93 85 100	1,867 1,529 2,145 2,191 2,477 1,476 2,078 2,274 2,235 2,084 2,234 2,820	25.23 25.91 26.48 25.48 26.92 26.84 19.79 25.27 20.32 22.41 26.28 28.20	153 117 141 157 160 123 189 180 188 173 175 182	5,186 3,594 4,257 4,872 4,905 3,596 4,072 4,629 4,477 3,991 4,944 5,240	33.89 30.72 30.19 31.03 30.65 29.23 21.54 25.72 23.81 23.07 28.25 28.79	
Whole year.	908	28,353	31.22	1,030	25,410	24.67	1,938	53,763	27.74	

The average age of all decedents in 1877 was 27.74 years; this was 0.67 of a year less than in 1876. The average age of American decedents in 1877 was 31.22 years, which was 1.95 years less than in 1876; the average age of foreign decedents, in 1877, 24.67 years, or 0.47 of a year more than in the preceding year. There were five months of the year 1877 in which the average age of

decedents of American parentage was greater than in the corresponding months of 1876, namely, January, February, April, June, and September; in the other seven months of the year, the average age was less than in 1876. As to the decedents of foreign parentage, there were also five months in 1877 when the average age was greater than in 1876; they were May, July, August, November, and December; in the remaining seven months of the year, the average age was less than in the preceding year.

Table XIV is a recapitulation, and gives these same statistics for the period of twenty-two years, 1856–1877 inclusive:—

TABLE XIV. DEATHS, TWENTY-TWO YEARS, 1856-1877.

Recapitulation. Aggregate and Average Age by Parentage.

	Ameri	can Pare	ntage.	Foreig	gn Paren	tage.	Total, American and Foreign.			
TWENTY- TWO YEARS, 1856-1877.	Number of Deaths.	Ag gregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	
January	1,239	42,653	34.43 34.34	1,059	26,620		2,298	69,273	30.14	
February March	1,199	39,976 42,207	35.20	1,023	22,614 27,135	24.18	2,187 2,280	62,590 69,342	28.62 30.41	
April		39,917	34.00	1,102	28,250		2,276	68,167		
May		40,414	36.53	1,008	25,748 22,731		2,115	66,192		
June July		31,988 32,739	25.88	1,513	27,723	23.85 18.32	1,887	54,719 60,462		
August	1,429	38,832	27.17	1,628	29,447		3,057	68,279	22.33	
September	1,266	37,053	29.27	1,393	26,218		2,659	63,271	23.80	
October	1,125	37,305	33.16	1,203	27,674	23.00	2,328	64,979		
November	1,150	38,421	33.41	1,118	28,354	25.36	2,268	66,775	29.44	
December	1,215	44,438	36.57	1,136	28,650	25.22	2,351	73,088	31.09	
Total	14,267	465,973	32.66	14,217	321,164	22.59	28,484	787,137	27.63	

A careful examination of these two tables will show many differences between the average age of American decedents, and that of foreign decedents in the different months. In the statistics for 1877 (Table XIII) we find the highest average age of decedents of American parentage is 42.01 years, in January; the lowest average age, 23.74 years, in July. Among the decedents of foreign parentage, the highest average age is 28.20 years in December; the lowest, 19.79 years in July. In every month the average age of the foreign decedents is less than that of the American, the greatest difference being in January, when it is

16.78 years. The least difference between the two classes is in August, when the average age of the foreign decedents is 0.90 of one year less than that of the American. The difference between them in the whole year is 6.55 years.

In the statistics for a longer period, as given in Table XIV, the differences are equally marked; here also the average age of the decedents of foreign parentage is uniformly less than that of the Americans. The highest average age of American decedents is 36.57 years in December; the lowest, 25.88 years in July. Of foreign decedents, the highest average age is 25.64 years in April; the lowest, 18.09 years in August. Thus the highest average age of the foreign decedents is less than the lowest average among those of American parentage. In the long period the greatest difference between the two classes is in February, where it is 12.23 years; the least difference, 7.56 years in July.

Taking the same months in the single year and in the long period of years, we can see many variations. These may be best shown by putting the figures in the form of a table, it being remembered that in every case, the average age of the decedents of foreign parentage was less than that of American decedents:—

Difference in years, between the average age of American and foreign decedents.

	_ _	1877.		22 years, 1856-1877.				
Months.	Averag	re Age.		Averag				
	American.	Foreign.	Difference.	American.	Foreign.	Difference.		
January		25.23	16.78	34.43	25.14	9.29		
February	35.20	25.91 26.48	9.69 8.72	34.34 35.20	22.11 24 18	12.23 11.02		
April	35.71	25.48 26.92	12.28 8 79	34.00 36.53	25 64 25 54	8.36 10.99		
June	31.18	26.84 19.79	5.34 3.95	34.25 25.88	23 85 18.32	10.40		
August September	26 17	25.27 20.32	0.90 8.42	27.17 29.27	18.09 18.82	9.08 10.45		
October	23 84	22.41 26.28	1.43 3.83	33.16 33.41	23 00 25 36	10.16 8.06		
December	29.51	28.20	1 31	36.57	25.22	11.35		
Whole year	31.22	24.67	6.55	32.66	22 59	10.07		

As a rule, the difference is greater in the long period than in the single year; January and April being the only exceptions. In the month of August, the difference in 1877 was but the fraction of a year, but in the period of years, it was 9.08 years. A

still greater contrast is in the month of December, when in 1877 the difference between the two classes was 1.31 years, while in the long period it was 11.35 years.

The largest monthly aggregate of deaths was in August, and the smallest in June, among both classes of decedents. The percentages in the different quarters for the entire period of twenty-two years were as follows:—

American parentage	1st quarter.	2d quarter.	3d quarter.	4th quarter.
Foreign parentage				

Table XV gives the number of decedents, and the aggregate and average age, according to parentage, in each of the twenty-two years, 1856–1877, and for the entire period:—

Table XV. Deates, 1856-1877.

Recapitulation. Aggregate and average age by parentage.

1	Americ	an Parei	ntage.	Foreig	n Paren	tage.		America Oreign.	n and
YEARS.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregato Years.	Average Years.	Deaths.	Aggregate Years,	Average Years.
1856	541	14,554	26.90	524	8.258	15.75	1,065	22,812	21.42
1857	523	16,314	31.19	402	8,007	19.91	925	24,321	26.29
1858	512		31.07	505	8,478	16.78	1.017	24,388	23.98
1859	479	14,993	31.30	420	8,640	20.57	899	23,633	26.28
1860	512	16,660	32.54	489	9,751	19.94	1,001	26,411	26.38
1861	538	16,327	30.34	513	9,605	18.72	1,051	25,932	24.67
1862	460	14,589	31.71	454	10,994		914	25,583	
1863	597	19,398	32.49	617	13,757	22.29	1,214	33,155	27.31
1864	649	21,732	33.48	632	11,365	17.98	1,281	33,097	25.83
1865	631	19,628	31.10	580	11,991	20.67	1,211	31,619	
1866	513	18,906	36.85	523	12,762	24.40	1,036	31,668	
1867	528	19,258	36.47	432	12,149	28.12	960	31,407	32,72
1868	596	21,779	36.54	514	12,539	24.39	1,110	34,318	30,92
1869	627	20,490	32.68	629	14,776	23.49	1,256	35,266	28.08
1870	642	21,078	32.83	621	14,393		1,263	35,471	28.08
1871	643	22,725	35.34	611	17,005		1,254	39,730	31.68
1872	832	27,572	33.14	771	19,160		1,603	46,732	29.15
1873	862	28,675	33.27	857	19,061		1,719	47,736	27.77
1874	885	27,588	31.17	1,102	24,802		1,987	52,390	
1875	914	30,419	33.28	1,001	24,302		1,915	54,721	28.57
1876	875	29,025	33.17	990	23,959		1,865	52,984	
1877	908	28,353	31.22	1,030	25,410	24.67	1,938	53,763	27.74
22 y'rs' total,	14.267	465,973	32.66	14.217	321,164	22.59	28,484	787,137	27.63

The difference between the two classes ranges from 6.55 years in 1877, to 15.50 years in 1864. The difference in the whole period is 10.07 years.

Number and percentage of decedents at different ages. In Table XVI are given the number and percentage of decedents, according to parentage, in each division of ages, in 1877; and the same statistics for the whole number of decedents during the twenty-two years, 1856–1877:—

TABLE XVI. DEATHS, 1877.

Percentage at different ages, according to parentage.

		rican itage. 77.		itage.	Total and can an eign.	d For-		ears, 0 1877.
Ages.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Under 1 year	162		203	19.71	365	18.83	5,639	19.79
1 to 2	54	5.95	87	8.44		7.28	2,571	9.03
2 to 5	83	9.14	135	13.11	218	11.25	2,556	8.97
Total under 5	299	32.93	425	41.26	724	37.36	10,766	37.79
5 to 10	69	7.60	62	6.02	131	6.76	1,363	4.78
10 to 15	19	2.09	20	1.94	39		577	2.03
15 to 20	17	1.87	34		51	2.63	976	3.43
20 to 30	80	8.81	109	10.58	189		2,910	10.22
30 to 40	70	7.71	86	8.35	156	8.05	2,624	9.21
40 to 50	70	7.71	78	7.57	148	7.64	2,218	7.79
50 to 60	74	8.15	62	6.02	136	7.02	2,036	7.15
60 to 70	84	9.25	70	6.80	154	7.95	1,992	6.99
70 to 80	87	9.58	62	6.02	149	7.69	1,853	6.51
80 to 90	35	3.86	17	1.65	52	2.68	969	3.40
90 and over	4	.44	5	.49	9	.46	200	.70
Total	908	100.00	1,030	100.00	1,938	100.00	28,484	100.00

The percentage of deaths of children under 5 years of age, in 1877, was less in both classes of decedents, than in 1876; of those of American parentage, 1.36 per cent. less; of foreign parentage, 1.87 per cent. less; of the whole number, 1.62 per cent. less.

The following statement is a recapitulation of these statistics for each of the twenty-two years, 1856-1877 inclusive:—

	American. Under five years.		Total American and Foreign Under five years.
1856	37.15 per cent	•	
	32.31 per cent		
	35.35 per cent	-	•
1859	28.60 per cent	48.34 per cent	37.82 per cent.
1860	32.81 per cent	48.87 per cent	40.66 per cent
1861	35.87 per cent	52.63 per cent	44.05 per cent.
1862	31.08 per cent	42.07 per cent	
1863	28.47 per cent	46.84 per cent	37.81 per cent.
1864	28.81 per cent	51.11 per cent	39.82 per cent.
1865	29.95 per cent	46.37 per cent	37.82 per cent.
866	23.78 per cent	40.53 per cent	32.24 per cent.
867	24.05 per cent	36.35 per cent	29.59 per cent.
868	27.35 per cent	43.19 per cent	34.69 per cent,
869	30.78 per cent	43.24 per cent	37.02 per cent.
1870	31.00 per cent	42.67 per cent	36.72 per cent.
871	28.62 per cent	36.01 per cent.	32.22 per cent.
872	30.89 per cent	40.00 per cent	35.27 per cent.
1873	30.39 per cent	43.05 per cent	
1874	34.02 per cent	43.55 per cent	39.31 per cent.
1875	32.06 per cent	40.96 per cent	36.66 per cent.
1876	34.29 per cent	43.13 per cent.	38.98 per cent.
1877	32.93 per cent	41.26 per cent	37.36 per cent.

In the thirty-eight years, 1840-1877 inclusive, there have been, in Providence, 41,629 decedents, of whom 16,501 were less than 5 years old; the number and percentage at each age under 5 years, were as follows:—

Whole number of decedents	41,629
Under one year	8,144, or 19.56 per cent. of all.
One, and under two	4,354, or 10.46 per cent. of all.
Two, and under five	4,003, or 9.62 per cent. of all.
Total, under five	16,501, or 39.64 per cent. of all.

Table XVII is a recapitulation, and shows the number and percentage of decedents at different ages, according to parentage, in the period of twenty-two years, 1856-1877:—

TABLE XVII. DEATHS, 1856-1877.

Number and Percentage at different Ages according to Parentage.

	age, 22 ye	an Parent- ears, 1856 to 877.		Parentage, 1856–1877.	Foreign F	can and arentage 1856-1877.
AGE.	Number of Deaths.	Percentage in each divi- sion of age.	Number of Deaths.	Percentage in each d'yi- sion of age.	Number of Deaths.	Percentage in each divi- sion of age,
Under one year. 1 and under 2 2 and under 5	2,470 921 1,047	17.31 6.46 7.34	3,169 1,650 1,509	22.29 11.61 10.61	5,639 2,571 2,556	19.79 9.03 8.97
Total under 5	4,438	31.11	6,328	44.51	10,776	37.79
5 and under 10.	695	4.87	668	4.70	1,363	4.78
10 and under 15.	314	2.20	263	1.85	577	2.03
15 and under 20.	534	3.74	442	3.11	976	3.43
20 and under 30.	1,355	9.50	1,555	10.94	2,910	10.22
30 and under 40.	1,269	8.89	1,355	9.53	2,624	9.21
40 and under 50.	1,122	7.86	1,096	7.71	2,218	7.79
50 and under 60.	1,179	8.26	857	6.03	2,036	7.15
30 and under 70.	1,214	8.51	778	5.47	1,992	6.99
70 and under 80.	1,279	8.97	574	4.04	1,853	6.51
80 and under 90. 90 and over	734 134	5.15	235 66	1.65	969 200	3.40
Totals	14,267	100.00	14,217	100.00	28,484	100.00

From Tables XIII and XIV, we saw that the average age of decedents of foreign parentage was considerably less than that of American decedents. We should expect, therefore, to find among the former class, a larger percentage of deaths of young children, and the above table shows this to be the case. Of decedents under five years of age, the percentage of American parentage is 31.11; while of foreign parentage, 44.51 per cent. are less than five years old. Between the ages of five and forty, there are slight differences in the percentages; sometimes one class being in excess, and sometimes the other; of decedents of forty years and over, the larger percentages are always of American parentage, 39.69 per cent. of this class living beyond the age of forty years, and only 25.36 per cent. of those of foreign parentage.

By comparing Tables XVI and XVII, it will be seen that the proportion of decedents under five years of age in 1877, was slightly below the average; namely, 0.43 per cent. The proportion of decedents of this class, of American parentage, was 1.82 per cent. above the average; the proportion of foreign parentage,

3.25 below. Of those who lived beyond the age of fifty years, the percentage of American parentage in 1877 was 31.28, or 0.55 per cent. below the average; the percentage of foreign parentage, was 20.98, or 3.33 per cent. above the average. This accords with facts to be obtained from Tables XIII and XIV, namely, that the average age of all decedents of American parentage in 1877, was 1.44 years below the average age of that class for the period of the twenty-two years, 1856–1877; and the average age of all foreign decedents in 1877 was 2.08 years above the average for the same period.

Deaths by age and parentage in each month and quarter of the year. Table XVIII shows the number of deaths at each age, and of all ages, in each month and quarter of the year 1877; the statistics being given according to parentage. From it may be seen at what season of the year occurred the greatest mortality in the different classes. The number of deaths in 1877 was 1,938; of American parentage, 908; of foreign parentage, 1,030; distributed through the year as follows:—

	1st quarter.	2d quarter.	3d quarter.	4th quarter.
American parentage	21.70	22.80	27.75	27.75
Foreign parentage	20.78	22.62	29.61	26.99

As has already been mentioned, the greatest mortality is in the third quarter of the year; from the above statement it will be seen that it is greater at that season among those of foreign than of American parentage; the same was true of a longer period of time, as shown in the percentages given on page 31 in the remarks upon Table XIV. In 1877 it was quite remarkable that of the decedents of American parentage, the percentage in the fourth quarter was exactly the same as in the third. In the third quarter, the number of deaths in this class was 252; of this number, 74, or 29.37 per cent., were less than one year old; this is 3.30 per cent. less of the same age than in 1876. Of the 162 children of American parentage under one year old who died in 1877, the 74 who died in the third quarter were 45.68 per cent., which is 4.62 per cent. less than in the preceding year. It is always the case that there is great mortality among very young children at this season of the year; in 1877 there was also a larger percentage of deaths than usual of children in the fourth quarter of the year; the number of decedents of American parentage in the fourth quarter was 252; of these, 120, or 47.62 per cent., were children under ten years of age; in 1876, the percentage of this class at the same season was 38.62; in 1875, 27.73. This excess in 1877 was caused by the extraordinarily large mortality from diphtheria in the last quarter of the year.

Among the decedents of foreign parentage, the number of deaths in the third quarter of the year was 305; of these, 93, or 30.49 per cent., were less than one year old; this is 3.05 per cent. less than in 1876. Of the 203 children of foreign parentage, under one year old, who died in 1877, 93, or 45.81 per cent., died in the third quarter; 2.17 per cent. less than in 1876. From these figures it may be seen that the mortality among children at this season in 1877, in both classes, American and foreign, was less than in the year 1876.

Causes of death; sex, age, parentage, and percentage from each cause. Table XIX gives the number of deaths in 1877 from each cause, according to sex and parentage, and in each division of ages; also the percentage from each cause of all from known causes.

Season and disease. Table XX, though also giving the number of deaths from each cause, is quite different in its details from Table XIX. It gives the number from each cause in each month and quarter, and in the whole year 1877, and is important as showing how the different diseases prevail at different seasons.

Table XVIII. Deaths by age and parentage in each month, 1877.

	1000			1	ivi	SION	AL	PER	IOD	S OF	Lu	P.H.				00
•	DIVISIONS OF THE YEAR.	Under 1.	1 to 2.	2 to 5.	5 to 10.	10 to 15.	15 to 20.	30 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.	80 to 90.	90 to 100.	ALL AGES.
	January February March	8 8 12	6 1	8 6 5	2 1	3	1 2	4 3 8	6 3	10 3 4	7 8 6	9 9 3	16 5 10	2 1 3	1	. 58
	1st quarter	28	7	19	3	5	7	15	15	17	21	21	31	6	2	197
L'ARENTAGE.	April. May. June.	9 8	3 1 4	5 6 7	3 10 5	2 2	1	12 3 8	5 7 6	6 5 4	7 7 6	7 8 6	7 7 7	6 3 2		71 68 68
X X	2d quarter	26	8	18	18	4	3	23	18	15	20	21	21	11	1	207
AMERICAN I'A	July. August. September	27 30 17	10 3 4	3	8	2	3	9 7 6	7 5 8	5 9 6	5 5	12 8	7 4 5	2 2 3		84 90 78
Z Z	3d quarter	74	17	12	18	3	4	22	20	20	15	24	16	7		252
3	October November December	11 11 12	976	14	8	3 1 3	3	5 9 6	4 5 8	4 7 7	6 10 2	6 6	6 8 5	1 4 6	'n	80 90 82
	4th quarter	34	22	. 34	30	7	3	20	17	18	18	18	19	11	1	252
	Whole year	162	54	83	69	19	17	80	70	70	74	84	87	35	4	908
	January February March	17 11 12	4 4 10	8 7 8	4 2 5	1 1 1	5 2 1	9 9	4 4 11	6 8 7	3 3 4	8 6 8	2 1 4	3 1 1		74 59 81
	1st quarter	40	18	23	11	3	8	27	19	21	10	22	7	5		214
LABENIAGE.	April	19 10 7	5 4	8 19 5	6 5 8	3 2 2	3	11 11 5	5 7 7	4 7 3	9 10 1	2 6 7	9 6 4	1 	1	80 92 55
EN	2d quarter	36	14	32	19	7	8	27	19	14	20	15	19	2	1	233
	July August September	41 27 25	7 9 15	4 7 15	4 4 7	3 3 2	5 3 2	9 3 9	10 6 9	7 5 10	4 3 3	5 7 6	5 11 5	1 1 2	i	105 90 110
FOREIGN	3d quarter	93	31	26	15	-8	10	21	25	22	10	18	21	4	1	305
4	October November December	18 6 10	7 7 10	17 19 18	5 7 5		3 2 3	9 11 14	9 5 9	4 10 7	11 6 5	3 6 6	4 5 6	6	1 1 1	93 85 100
	4th quarter	34	24	54	17	2	8	34	23	21	22	15	15	6	3	278
	Whole year	203	87	135	62	20	34	109	86	78	62	70	62	17	5	1,030

TABLE XIX. DEATER, 1877. Disease, sex, age, parentage, and percentage from each cause.

	æ	SEX.					DIV	STOR.	At. P1	DIVISIONAL PERIODS OF LIFE	0 8	Lin					PAR	PARENTAGE		KBOWn Whole
CAUSES OF DEATH.	Malca	Femalos.	Total.	Your. Your. Dan I	under 2.	under 5. 5 and	Of 19bau 10 and	la sinder 15. Sand 16. 20.	20 and	So and 30 and 10.	40 and under 50.	50 and under 60.	On no on under 70.	70 and under 80. 80 and	under 90. bun 08	OVOF.	Americani.	Poreign.	Total.	Ratio of death each cause to number from l causes.
Abecesses.	€ ₹		-4-82	- 9	<u>: </u>	 -:-		: :		1 61	1		- "		<u> : </u>	<u> </u> ; ;	6116	ei g	4 %	មន្ត
	∞ ×	-	E K	-	-ca	c1 -		<u> </u>	<u>:</u>				; <u></u>			::	N) F	∞ ₹	2	9
Falls	==	87-		<u> </u>	<u>:</u> : :	<u>.</u>	, <u>:</u> -	<u>: </u>		- 61 - 61	. :	C1 +	61	 -		::	- 1- ,	, 0 ,	<u> </u>	863
Railway	110	1:	1 20 -	: : -	<u>:</u>	:-:-	-	<u>:</u>		: -	- C1	- :		: :		::	= c17	- €	1 10 ·	58
Anemis	- • •	-:}	- - ;		<u>: :</u>	:- : -	\vdots	<u>: :</u>	<u></u>	:"	CI	::		<u>:</u> -		::		:60	- 4	84
	3-	3 ⁻⁶¹	3 m			<u>:-:</u>	<u>: :</u> : :	<u>: :</u>	-	°1 :	ے :	ĉ. - -	55 21	<u>:</u> 2	<u>- ا</u>		8	<u>ಹ ಜ</u>	8 0	2.39
Bladder, Diseases of	<u> </u>	:::	9 6			-:	<u>:</u>					61	ຕັ	:-	:-	:	ကင	က	· · · ·	£.5
Brain, Congestion of	4 5	ńĖ	ي ص <u>د</u>	4.0	 	-:-	:	:		- 4	:	8	_ e		-	::	101	. eo ×	1 C. 2	7
	នគ	8 8	۱Ę4	121	: :	. : 2°	<u>:</u>	<u> </u>		- FG	1014	C1 -	C1 :			::	1815	ह	12:	35.
Cancer (various)	œ	2 2	228		,	,				100	1016	100) (~ K	- -	 -	::	3 2 5	3 <u>1</u> E	223	98
	4	===	55	-:-		-:-		<u>:</u>	-	-		₩ 4	4 4	ci-			11- 5	- œ c	12.3	Ė
_		œ €1	œ c						-	· •• -	~_	'	,	-	+	::	7-6	1	100.0	345
Chicken Dow	:	<u></u>	-		-		<u>:</u>	:		 		<u> </u>		: :	÷	::	1 -+ -	m	11	8,8
·52	24	귏애	ដ្ឋាទ	· 2	:: -a	-		! ! !			: : : -	<u> </u>		=======================================		:::	- - -	6. 20	<u> 81</u> .0	3. E. E.

itatio of destha from cach cause to whole number from known number Disease, sex, age, parentage, and percentage from each cause. Total. PARKNTAGE. Foreign. American. .1970 **bas** 06 80 and under 90. 70 and .08 Tobau 60 and 10. 100. DIVISIONAL PERIODS OF LIFE. 50 and 05. 40 and .05 Tobau -8 30 and under 40. 20 and 100 nader 30. Is and .02. 20. lo and Gl Tebuu 6 and under 10. S and ander 5. **DEATHS**, 1877. l sad Lander 2. Under 1 Year. Total. BEX. Females. CONTINUED. Maloa Dropsy
Dysenkery
Enteritis
Epilopsy
Erysipolas Chores. Debility. Premature Birth.... Diphtheria Hip Joint, Disease of Hooping Cough Hydrocephalus Hydrophobia Consumption Diabetes..... Gaugrene, Senile Glanders Heart, Discases of Fever, Malarial................. Convulsions Chronic.... CAUSES OF DEATH. Hypertrophy of TABLE XIX. Typhold. Hemorrhage

TABLE XIX. CONTINUED. DEATHS, 1877. Disease, sex, age, parenlage, and percentage from each cause.

	-	SEX.	i				ā	IVISI	DIVISIONAL PERIODS OF LIFE.	PER	HODS	30	LIVE.				1	PARE	PARENTAGE.	1	o whole
JAUSES OF DEATH.	Males.	Females.	Total	Underl	I and I and under 2.	2 and under 5.	5 and under 10.	10 and	15 and under 20.	20 and under 30.	30 mder 40.	40 and	50 and under 60.	60 and under 70.	nuger 80,	80 and moder 90.	90 and over	American	Foreign.	Total.	Ratio of deat each cause to number from causes,
nsanity ntemperance Addisor's Delirium Tremens Addisor's Diseases of Addisor's Disease of Addisor's Disease of Addisor's Disease of Addisor's Diseases of Antary Aren, Diseases of Antary Arennights, Cerebro-Spinal Arrder and Homicide Reglect Place Place Addisor's Disease of Arennights, Cerebro-Spinal Arrder and Homicide Reglect Arrange Arran	2000HF-184-1100 H-101 : 420 : 80H-1:	22 : 122 : 1 : 22 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 -	のでいるないないといっているないののは、	:::3:::::::::::::::::::::::::::::::::::		C1 :H1 :	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		я н мн	פי היי היי היי היי מחה מחה	04 : HHO : 00 H : : : : : : : : : : : : : : : :	нн н н ю пннан н	01	HO : 0 10 10H 4 . HOH 10HH	тенне по на не	H 184 0101		4	85484 THE	 @###################################	444444444444444444444444444444444444444

Table XIX. Continued. Deates, 1877. Diwase, set, age, parentage, and percentage from each cause.

•		Sex	×					DIVISIONAL PERIODS OF LIFE	IOKA	L Pr	RIOD	8 OF	Lim		 	İ	!	PAR	PARENTAGE	, #	morita Solw gwoga
6	CAUSES OF DEATH.	Males. Females.	Total	I 19baU	Year.	bas 2	5 and 5 and 10, under 10,	lo and of and and and of the state of the st	Is and .0% Tebru	bna 02 .06 19bnu	50 and 06 .04 Tebau	40 and 50.	50 and under 60.	60 and under 70.	70 and 70 and 100.	80 and under 90.	90 and over.	Атменсвл.	Foreign.	Total.	Ratio of death cach cause to mumber from
	Quinay Rheumatism Rickets Releats Searlatus Serofula Skin, Disease of Spleen, Disease of Spleen, Disease of Stricture Stricture Stricture Sucide Burgical Operations Syphilis Tabes Meenterica. Teething Thush. Tuberculosis Tuberculosis Ulloer Ulloer			<u>wr-ww-wwd-woor-wada-o</u>	: : : : : : : : : : : : : : : : : : :										9			88 : 64 : HEQHERRUGHER : 4	:	ar-18 2 2 2 2 2 2 2 2 2	
•	Total	922 101	1938	, ,	365 141	1 218	131	66	21	8	156	148	136	151	₹ 6	ន	6	8	1,030	1,938	100.00

1

Table XX. Deaths, 1877. Season and Disease.

Whole Year.	第283年度12日本 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
Fourth Quarter,	HENNESS - ETTER & LANCESS & ALWELL COLS
December.	© © © 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
November,	2000 - 100 -
October.	000
Third Quarter.	818-42 HoF40282251-38-201202
September.	מיניניין בא מיניין בא מיניין בא בא מיניין בא בא בא מיניין בא
August,	04000-8450 OH440 F-044 H000 10
July.	400-000-0000000000000000000000000000000
Second Quarter,	80000404 :800000000000000000000000000000
June,	1000000000 Houne House 4 10 104
May.	DONA 4400 SOMETH HIS OFF TAXON SO
April.	040040 : Son-Wall - 1040 : - 10
First Quarter.	454555 : 61000 t : 10040 20040 5004 : 12
March.	M
February.	DOOD-DH : 704- 2 12000 : 0 : 12
January.	© © © © © © © © © © © © © © © © © © ©
CAUSES OF DEATH.	Accidents (all kinds) Apoplexy Brain, Diseases of Brain, Diseases of Brain, Diseases of Brain, Diseases Cancer (all kinds) Child-birth and Puerperal Diseases Cholera Infantum Cholera Infantum Cholera Infantum Cholera Infantum Cholera Acute and Chronic Diphtheria Diphtheria Diphtheria Dropsy Dropsy Entertitis Erysipelas Erysipelas Erysipelas Erysipelas Entertitis Entertit

Table XX. Continued. Deaths, 1877. Season and Discuse.

	<u> </u>	: !	-	,= 	_	_	_	-		•	•		_	-	! -	. – :	•
Сливы ог Велти.	January.	Рергиалт.	жиср.	First Quarter.	April.	May.	June.	Second Quarter.	July.	August.	September	Third Quarter.	October.	November	December.	Fourth Quarter.	Whole Year.
Paralysis Partonitis	1		:-	-6	₩-	=	4.2	66	8	8	:-	9-		40	8	oc ec	% 5
Pneumonia and Congestion of Lungs	_	13.	Ţ.	8	2	00	41	13	တ	7	9	10	'9	11-	2	22	88
ScarlatinaSuicide	eo -	:6	:-	හ 4	69	:-	6	30 ec		<u>:</u> -	:	67	:	4:	01 -	90	22
Teething	177	' : ' : '	· :	1770		· :			-		• : ·	00,	-	161	1 1	₹,	c
Other causes as given	183	141	17	48	72	18	10	49	19	21	19	26	:81	:00	78	22	216
Total	153	117	141	411	167	160	123	440	189	188	188	557	173	178	182	230	1,938

NOTES ON THE CAUSES OF DEATH,

DURING THE YEAR 1877.

These notes are to call attention to some of the facts given in Tables XIX and XX; they also give some statistics not found there, and compare the same with the statistics for the twenty-two years, 1856–1877 inclusive.

The percentage of deaths from the various causes is always reckoned upon the number of deaths from known causes.

The number of deaths in Providence in 1877 was 1,938, of which only 6, or 0.31 per cent., were from unknown causes. The number of which the cause was stated, and on which all the percentages are reckoned, was 1,932. During the twenty-two years, 1856–1877, there were 28,484 deaths, of which 508, or 1.78 per cent., were from unknown causes, and 27,976 from known causes. During the thirty-eight years, 1840–1877, the number of deaths was 41,629; 1,510, or 3.63 per cent., from unknown causes, and 40,119 from known causes.

In Table XX is given the number of deaths from each cause in each month and quarter of the year, and by it may be noted the comparative prevalence of different diseases at different seasons. Thus, of deaths from diseases of the brain, the largest number was in the first quarter of the year; of those from bronchitis, croup, and diphtheria, the largest number was in the last quarter; from pneumonia and congestion of the lungs, in the first quarter, and from consumption in the second quarter. All diarrhoeal diseases are of course chiefly in the third quarter of the year; and the deaths from typhoid fever are almost entirely in the third and fourth quarters, 81.40 per cent. of them occurring in the last six months of the year.

Abscesses. The number of deaths reported from abscesses, in 1877, was 4, or 0.21 per cent. of all from known causes; this

was just half as many as in 1876. Those specified are one psoas abscess, and one each of the pelvis and the perinæum. The decedents were 3 males, and one female; 2 of American and 2 of foreign parentage. The statistics for the 22 years, 1856–1877, are as follows:—

Number of deaths from all known causes, 27,976. From absceses: males, 44; females, 37; total, 81, or 0.29 per cent. of all. American parentage, 36; foreign parentage, 45.

Accidents. The number of deaths from accidents of all kinds, in 1877, was 56; three more than in 1876, and 2.90 per cent. of all from known causes. Of these, there were specified in the tables, 13 from burns and scalds; 5 from drowning; 13 from falls; 2 from poisoning; and 5 railway accidents; 18 more were from various causes, not named in the tables; some of these from various causes were as follows: two died of asphyxia, caused in one case by a gas stove; the other, not specified; three were run over by wagon or dray; one was kicked by a horse; one killed while coasting; one by a blow on the head, from a crank, and five from injuries received at or before birth. Of those from burns and scalds, two were from the breaking or explosion of a kerosene lamp; two from the upsetting of a tea-pot, and of two the record is the "clothes caught fire." Of the falls, 3 were "down stairs"; 2, "into the hold of a vessel"; 2, "fall of staging"; 2, "falling of a bank of earth." Of those by poisoning, one was by morphine, and one by spirits of ammonia. Of the five deaths by railway accidents, two were run over by the cars, and three were hit by the cars, or by a railroad bridge.

The statistics of sex and parentage are as follows:—

Males, 42; females, 14; total, 56, or 2.90 per cent. of all. American parentage, 21; foreign parentage, 35.

For the twenty-two years, 1856-1877: —

Males, 617; females, 198; total, 815, or 2.91 per cent. of all. American parentage, 324; foreign parentage, 491.

Aneurism. Four males, one of American, and 3 of foreign parentage, died from aneurism in 1877; in two cases, it was of the aorta, and in one, aneurism of the left carotid; one, not specified; the four were 0.21 per cent. of all deaths from known causes.

For the twenty-two years, 1856-1877, the statistics of Aneurism are:—

Males, 16; females, 3; total, 19, or 0.07 per cent. of all. American parentage, 8; foreign parentage, 11.

Apoplexy. This was returned as the cause of 50 deaths in 1877; 6 more than in 1876, and 2.59 per cent. of the whole number from known causes. Males, 25; females, 25; American parentage, 32; foreign, 18. (Vide "Paralysis.")

The statistics in regard to apoplexy for the twenty-two years, 1856-1877, are as follows:—

Males, 276; females, 302; total, 578, or 2.07 per cent. of all. American parentage, 416; foreign parentage, 162.

Brain, Diseases of. There were 82 deaths in 1877 from various diseases of the brain; 9 were specified as "congestion," and 51 as "inflammation"; many of the others were "softening of the brain." The particulars are:—

Males, 44; females, 38; total, 82, or 4.25 per cent. of all. American parentage, 45; foreign parentage, 37.

For the twenty-two years, 1856-1877:-

Males, 601; females, 496; total, 1,097, or 3.92 per cent. of all.

American parentage, 586; foreign parentage, 511.

Bronchitis. The number of deaths from bronchitis in 1877 was 44, or 2.27 per cent. of all from known causes. Males, 21; females, 23; American parentage, 25; foreign, 19.

The statistics for the twenty-two years, 1856-1877, are as follows:—

Males, 142; females, 180; total, 322, or 1.15 per cent. of all.

American parentage, 145; foreign parentage, 177.

Cancers. The number of deaths from cancers in 1877, was 66; 13 more than in 1876, and 3.41 per cent. of all from known causes. As usual, by far the greater number of decedents were females, namely, 54 females and 12 males; of American parentage, 37; of foreign parentage, 29. In the tables there are specified, 18 from cancer of the breast; 15 from cancer of the stomach, and 12 from cancer of the uterus; besides these there

were 6 from cancer of the liver; 3, from cancer of the rectum; 2, of epithelial cancer, and one death each from cancer of the lip, jaw, mouth, throat, bladder, bowels, and right axilla; and three were not specified.

The statistics of deaths from cancer for twenty-two years, 1856-1877, are:—

Males, 163; females, 484; total, 647, or 2.31 per cent. of all. American parentage, 435; foreign parentage, 212.

Childbirth. There were 17 deaths in 1877 from childbirth; 0.87 per cent. of all from known causes, and 0.68 per cent. of the 2,516 of mothers bearing children in 1877. Of these seventeen deaths, 2 were from puerperal convulsions, and 7 from puerperal fever. The decedents were 7 of American, and 10 of foreign parentage.

In the twenty-two years, 1856–1877, the number of deaths from childbirth were 338, or 1.21 per cent. of all from known causes, and 0.83 per cent. of the 40,746 mothers who bore children during that period. The number of American mothers was 18,937; of these, 143, or 0.75 per cent., have died; the number of mothers of foreign parentage was 21,809, of whom 195, or 0.89 per cent., have died.

Cholera Infantum. The number of deaths from this cause in 1877 was 122; the same number as in 1876, and 6.31 per cent. of all from known causes. There were 97 decedents under one year old; 21 between one and two, and four between two and five. Males, 70; females, 52; of American parentage, 43; of foreign parentage, 79.

The statistics are as follows, for each of the twenty-two years, 1856-1877 inclusive, and for the whole period:—

	Males.	EX. Females.	PARES American.	TAGR. Foreign.	Whole Number.	Percentage
1856			16			
			26			
			13			
			13			
			26			
1861	29	32	25	36	61	6.02
1862	25	19	17		44	4.98
1863	36	30	19	47	66	5.56
1864	29	29	18	40	58	4.63
1865	20	30	13	37	50	4.21
1866	30		15	32	47	4.64
1867	24		23	26	49	5.19
1868	38	32	22	48	70	6.44
1869	35	30	27	38	65	5.24
1870	48	45	37	56	93	7.44
1871	30	32	26	36	62	5.00
			58			
			39			
			44			
			50			
			43			
1877	70	52	43	79	122	6.31
vears' total	s866	813	613	1.066	1,679	6.00

Consumption. There were 294 deaths from consumption in 1877; 15.22 per cent. of all from known causes. This was 10 more deaths than in 1876, but 0.12 per cent. less. But it is the first time since the beginning of these reports that the deaths from consumption have not exceeded, and generally, largely exceeded, those from any other cause; the numbers for 1877 show just one more death from diphtheria than from consumption. The decedents from consumption were divided as follows as to sex and parentage:—

American parentagemales, Foreign parentagemales,	•		
	-		
Totalmales,	127 females,	167total,	294

For the twenty-two years, 1856-1877, the statistics are as follows:—

Whole number, 4,770, or 17.05 per cent. of	all.		
American parentagemales,	1,002females,	1,339total, 2,3	41
Foreign parentage males,	1,159females,	1,270total, 2,4	129
			_
Totalmales,	2,161females,	2,609total, 4,7	770
Males, 45.30 per cent	females	, 54.70 per cent.	
American parentage, 49.08 per cent	.foreign parentage	, 51.92 per cent.	
			_

Estimating the population of the city in the middle of the year 1877 at 100,675, we find the death-rate from consumption to be one in 342 of the population, or 2.92 deaths to each thousand persons; in 1876, the death-rate was one in 357, or 2.80 in each thousand. The average annual rate for the fifteen years, 1840–1854 inclusive, was one in 209.9, and for the twenty-three years, 1855–1877, one in 300.2.

By periods of 5 years, from 1840 to 1874 inclusive, and for the whole period of 35 years, the deaths from consumption in Providence were as follows:—

	Deaths from consumption.	Per cent. of all from known cause
1840-1844	745	
1845-1849	761	
1850-1854	958	
18 55-1859	930	
1860-1864	997	
1865-1869		
1870-1874		15.1
		·
1840-1874, 35 year	s	

The deaths in each of the last eight years have been :-

	Number.	Per cent
1870		
1871		
1872		
1873		
1874		
••••	294	

The number and percentage of the decedents from consumption, in each division of age, were as follows for the year 1877, and for the period of 22 years, 1856 to 1877 inclusive:—

PERIODS,	Under 1 year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60,	60 and under 70.	70 and under 80.	80 and under 90.	TOTAL
1877. Number	4	1	2	4	5	25	106	49	40	32	19	7		294
Per cent	1.36	.34	.68	1.36	1.70	8.50	38.06	16.67	13.61	10.88	6.46	2.38		100.00
1856-1877. Number	62	56	53	49	90	436	1,459	1,000	679	450	310	115	n	4,770
Per cent	1.30	1.17	1.11	1.03	1.89	9.14	30.59	20.96	14.24	9.43	6.50	2.41	.23	100.00

It will be seen that the largest percentages are between the ages of 20 and 50 years, 66.34 per cent. of the decedents being within these limits; just 50.00 per cent. lived beyond the age of 30 years. The average age of decedents from all causes, in 1877, was 27.74 years. Of the decedents from consumption in the twenty-two years, 1856–1877, 53.77 per cent. lived beyond 30. years of age; the average age of decedents from all causes in the same time, was 27.63 years.

Convulsions. The number of deaths from this cause in 1877 was 38; 14 less than in 1876, and 1.97 per cent. of all from known causes. Males, 18; females, 20; American parentage, 16; foreign parentage, 22.

The following are the statistics for the twenty-two years, 1856-1877:—

Males, 405; females, 328; total, 733, or 2.62 per cent. of all. American parentage, 306; foreign parentage, 427.

Croup, was the cause of 40 deaths in 1877; 2.07 per cent. of all from known causes. This was 25 less deaths, and 1.44 less per cent., than in 1876. The decedents were all less than ten years old; and just 50 per cent. were between the ages of 2 and 5 years. Males, 18; females, 22; American parentage, 14; foreign, 26.

For the twenty-two years, 1856-1877, the statistics are: -

Males, 332; females, 326; total, 658, or 2.35 per cent. of all. American parentage, 267; foreign parentage, 391.

Debility. This was reported as the cause of 26 deaths in 1877, including 10 from premature birth; 1.35 per cent. of all from

known causes. All but two of the decedents were less than one year old. Number of males, 16; females, 10; of American parentage, 16; of foreign parentage, 10.

The statistics for the twenty-two years, 1856-1877, are:

Males, 267; females, 236; total, 503, or 1.80 per cent. of all. American parentage, 201; foreign parentage, 302.

Diarrhæa and Dysentery. The number of deaths from these diseases in 1877, was as follows: acute diarrhæa, 29; chronic diarrhæa, 5; dysentery, 21; total, 55, or 2.85 per cent. of all from known causes. The number of deaths was seven less than in 1876; the per cent., 0.50 less. The number of males in 1877, 21; of females, 34; of American parentage, 27; of foreign parentage, 28. It is quite unusual to find the two classes of decedents so nearly equal; those of foreign parentage being generally considerably in excess of those of American parentage.

In this connection may be given the following statistics concerning diseases of a similar nature: Deaths from Cholera Infantum, 122; from Cholera Morbus, 6; from Enteritis, 21; these combined with the numbers given above for Diarrhæa and Dysentery make the total of deaths from diarrhæal diseases as follows:

Males, 106; females, 98; total, 204, or 10.56 per cent. of all. American parentage, 80; foreign parentage, 124.

The number of deaths is 2 less than in 1876, and the per cent. 0.57 less. The statistics for these five diseases for the twenty-two years, 1856–1877, are as follows:—

Whole number of deaths, 3,365; annual average, 153.

Males, 1,688; females, 1,677; total, 3,365, or 12.03 per cent. of all.

American parentage, 1,359; foreign parentage, 2,006.

It will be seen how large a majority of these decedents are of foreign parentage, the proportion being 148 of this class to each 100 of American parentage; or 59.61 of foreign and 40.39 of American parentage in each 100 decedents. There are one or two facts which serve to explain this; the mortality from these diseases is always largely among young children; from Table XIX it may be seen that 155 of the decedents in 1877, or nearly 76 per cent., were children under two years of age. The number of children of foreign parentage is considerably larger than the number of American parentage. In the twenty-two years,

1856-1877 inclusive, the number of children born was 41,273; of these, 22,625, or 54.82 per cent., were of foreign parentage; and 18,648, or 45.18 per cent., of American parentage; or 121 foreign to each 100 American. In 1877 the decedents from diarrheal diseases were 155 of foreign to each 100 of American parentage; of children born, the proportion was 105 foreign to each 100 American. It is moreover frequently the case that the children of foreign parentage live under less favorable conditions than those of the other class, and through the ignorance or neglect of the parents are more liable to be attacked by these diseases.

The following table shows the number of deaths from each of these causes, with the number from the whole class, and their percentage of all from known causes, reported in Providence during the period from 1840 to 1854 inclusive; during each year, from 1855 to 1877 inclusive, and in the aggregate during the whole period of 38 years, 1840 to 1877 inclusive:—

	Cholera	Cholera D	fameliona 1	Desantore	Pataritia (All of all from
1840-1854				796		
1855						
1856	57	5	. 29	35	. 6	. 13211.17
1857	50			29		
1858	49	2	. 27	12	. 8	. 98 9.63
1859	33	1	. 28	13	. 4	. 79 8.78
1860	68	2	. 21	18	. 9	. 11811.78
1861	61	5	. 22	32	. 6	. 12611.99
1862	44	1	31	22	. 9	. 10711.70
1863	66	5	. 35	77	. 9	. 19215.81
1864	58	4	. 45	40	13	. 16012.49
1865	50	7		62		
1866	47	17				
1867	49	2	. 15	43	. 10	. 11912.39
1868	70					. 12010.81
1869	65	6	24	23	. 6	. 124 9.87
1870	93	3		16		
1871	62	6				. 12610.15
1872	151	10	52			The state of the s
1873	92	4	37	7	. 25	
1874	134	3	50	17		
1875		6		0.0000000000000000000000000000000000000		,
1876	122	4				
77577333	122			21		
201111						

The number of deaths from known causes in the thirty-eight years, 1840–1877, was 40,119.

Deaths from Asiatic cholera are not included in these statistics. They have been as follows: in 1849, 158; in 1854, 159; in 1866, 14; total, 331.

Diphtheria. In the report of 1876, it was stated that this disease prevailed, in that year, to an extent never before known in Providence, and caused 111 deaths, or 6.00 per cent. of all from known causes; males, 53; females, 58; American parentage, 43; foreign, 68. But in 1877 these numbers were much more than doubled, the statistics being as follows:—

Males, 146; females, 149; total, 295, or 15.27 per cent. of all. American parentage, 129; foreign parentage, 166.

These deaths were distributed through the year, as follows: 1st quarter, 15.93 per cent.; 2d, 23.73 per cent.; 3d, 20.34 per cent.; 4th, 40.00 per cent. In the different wards of the city the deaths were:—

I62	VII29
II 8	VIII11
III34	IX38
IV20	X56
V24	
VI13	Total295

As in previous years, the mortality was largely among children, 276 of the decedents, or 93.56 per cent., being less than ten years old. The first cases of diphtheria reported in Providence, were in 1858; for the twenty years, 1858–1877, the statistics are as follows:—

Decedents from all known causes in 20 years, 26,048.
From diphtheria, 740; annual average, 37.
Males, 361; females, 379; total, 740, or 2.84 per cent. of all.
American parentage, 368; foreign parentage, 372.

The 295 deaths from diphtheria in 1877 are eight times the annual average, and comprise 39.86 per cent. of the number from this disease in the whole period of twenty years. In the following table, are given the statistics as to sex, parentage, season, and locality, for each of the twenty years, 1858–1877, and for the entire period:—

DEATHS FROM DIPHTHERIA IN PROVIDENCE, TWENTY YEARS.

Total.	28.8	361	368	858422845558	H8H488188
1877.	205 15.27	146	120	######################################	8828282828
1876.	111	588	38	は 一本ででののななない	00184000F80
1875.	14 0.74	900	104	1 00 HOS HOSTH	H0303 03H 0003H
1874.	1.00	99	97	0505H 0570H H0005H	- : : : : : : : : : : : : : : : : : : :
1873.	88.	140	158	004H 4 HH H	20 HO CO CO CO
1872	12 E	110	8-	10 4 10 05 HO 05 HO 05 HO	00-03-4-03-03-4-00
1871.	1.69	230	7-	HH03HH034 : :00403	4 4
1870.	10	1010	40	H H H 63 63 60	m 1-
1869.	13	0010	Ø.	000 HH H01HH	00HHHH01 4
1868.	0.46	10	100	111111111111111111111111111111111111111	01 00
1867.	0.53	60.03	14	04	H
1866.	11.08	50.00	910		
1865.	1.35	27 *	10	H 01 H H 03 4 00 03	H-000101+00 :::
1864.	3.27	813	828	00 40 5-60 40 H 05 05 00 05 00 40	F-F-5305001000
1863.	3.54	32.2	80	₩ 10 10 10 10 10 10 10 10 10 10 10 10 10	# 10 10 10 10 10 10 10 10 10 10 10 10 10
1861.	1.82	134	129	00 H H H H H H M M M	
1881	85.17 71.19	220	510		G-8 884
1880.	8.60	##	169	01-H 0100-H0000	
1859.	1.98	8000	19	00 HH 400	HH- 65.05
1858.	0.50	-4	411	OS MAH	# H
Бичинки.	Whole number Percentage	Males. Females. Papentage	American Foreign	January Rebrandy Rebrandy March April April Alay June June September September October Docember	Ward II

The subject of the connection of season with the mortality from diphtheria is exceedingly interesting and important of itself, and still more so when the mortality from this disease is compared with that of scarlatina and croup. The following table shows the number and the percentage of deaths in Providence, from diphtheria, scarlatina, and croup, in each month and quarter of the year. The figures for scarlatina and croup include the aggregate mortality from these diseases during the twenty-three years, from 1855 to 1877 inclusive. The figures for diphtheria are for twenty years, from 1858 to 1877 inclusive.

		HERIA. -1877.		ATINA. -1877.	CROUP. 1855-1877.	
Montes.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
January	50	6.76	192	12.87	88	
February	45		170		79	11.52
March	56	7.57	148	9.92	75	10.93
1st Quarter	151	20.41	510	34.18	242	35.28
April	42	5.68	141	9.45	54	7.87
May	51	6.89	145	9.72	40	
June	47	6.35	136	9.12	25	3.64
2d Quarter	140	18.92	422	28.29	119	17.34
July	30	4.05	80	5.36	26	3.79
August	24		60		10	
September	72	9.73	67	4.49	40	5.83
3d Quarter	126	17.02	207	13.87	76	11.08
October	120	16.22	77	5.16	67	9.77
November	113	15.27	117	7.84	84	12.24
December	90		159	10.66	98	14.29
4th Quarter	323	43.65	353	23.66	249	36.30
Totals	740	100.00	1,492	100.00	686	100.00

Dividing the year into periods of four months, the numbers and proportions are as follows:—

Gr ugova	Dipнī	HERIA.	SCARL	ATINA.	CROUP.		
Seasons.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	
1st 4 months 2d 4 months 3d 4 months		26.09 20.53 53.38	651 421 420	43.63 28.22 28.15	296 101 289	43.15 14.72 42.13	

Dividing the year into two periods, the number and proportions are:—

Walter	DIPHT	CHERLA.	SCARL	ATINA.	Choup.		
SEASONS.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent	
1st 6 months 2d 6 months	291 449	39.33 60.67	932 560	62.47	361 325	52.62 47.38	

The most cursory examination of the preceding figures shows very great differences in the three diseases named, in regard to the season of mortality.

With regard to *Diphtheria*, we find a very large excess of mortality in the fourth quarter of the year, the percentage in that quarter being more than double that of any other quarter.

Scarlatina shows the greatest mortality in the first quarter, with a mortality nearly as great in the second quarter.

Croup has the greatest mortality in the fourth quarter, though very slightly different from that of the first quarter.

The least monthly mortality from all three diseases, is in August; the greatest monthly mortality is in October, from diphtheria; in January, from scarlatina; in December, from croup.

Dividing the year into periods of four months each, the differences in the diseases are still more marked. More than half of all the deaths from diphtheria are in the last four months of the year. Nearly forty-four per cent. of all the deaths from scarlatina are in the first four months of the year. With regard to croup, the percentages of deaths in the first four and in the last four months are nearly equal. With the year divided into two periods of six months each, we find 60.67 per cent. of all the deaths from diphtheria in the last half of the year; and 62.47 per cent. of all the deaths from scarlatina in the first half of the year; while the deaths from croup are nearly equally divided, with a slight excess in the first half of the year.

It would seem that all three diseases are more especially prevalent in cold weather; diphtheria in the autumn, scarlatina in the latter part of winter and in the spring, while the prevalence of croup seems to depend, very uniformly, upon the degree of cold, the disease increasing quite regularly from August to December, and decreasing from December to August.

Dropsy was reported as the cause of five deaths in 1877; 0.26 per cent. of all from known causes; males, 2; females, 3; of American parentage, 2; of foreign parentage, 3.

Fever, Typhoid. The number of deaths reported from this cause in 1877, was 43; 10 more than in 1876, and 2.22 per cent. of the whole number from known causes. Number of males, 20; females, 23; American parentage, 22; foreign, 21. In the following statistics for the twenty-two years, 1856–1877, besides the deaths from typhoid, are included the few that have been reported, in some years, from "fever," "bilious fever," and about 30 from typhus fever:—

Whole number, 935; annual average, 42.

Males, 489; females, 446; total, 935, or 3.34 per cent. of all.

American parentage, 517; foreign parentage, 418.

Heart, Diseases of. The number of deaths from diseases of the heart, in 1877, was 93; males, 47; females, 46; American parentage, 53; foreign parentage, 40; total, 93, or 4.81 per cent. of all from known causes. The statistics for twenty-two years, 1856–1877, show 1,284 deaths from diseases of the heart, or 4.59 per cent. of all; annual average, 58. The following are the details for each year:—

	Males.	Females.	PARE American	NTAGE. Foreign.	Whole Number.	Percentage
1856						2.03
1857	13	18		9	31	3.45
1858	13	23		8	36	3.61
1859	13	11		7	24	2.76
1860	15	16	21	10	31	3.23
1861	23	18	29	12	41	4.05
1862	26	23		16	49	5.54
1863	23	19		12	42	3.54
1864	26	32	. 44	14	58	4.63
1865	18	29		14	47	3.95
1866	19	21		13	40	3.95
1867	26	23		14	49	5.19
1868	25	27	34	18	52	4.79
1869	33	29		27	62	5.00
1870	28	31		26	59	4.72
1871	40	37	46	31	77	6.22
1872	43	49	63	29	92	5.82
1873	46	54		41	100	5.88
1874	46	58	62	42	104	5.29
1875	43	45	47	41	88	4.65
1876	41	47		37	88	4.75
1877	47	46		40	93	4.81
22 years' total.	<u></u> 615	669	818	466	1,284	4.59

Hemorrhage. There were 9 deaths reported from this cause in 1877. One case was called "internal" hemorrhage; 5, umbilical; 2, hemorrhage of the bowels, and one, of the lungs. Number of males, 6; females, 3; American parentage, 6; foreign, 3; total, 9, or 0.47 per cent. of all from known causes.

Hooping Cough. There were 16 deaths from this cause in 1877; less than half the number in the preceding year. Just one half of the decedents were less than one year old; and nearly half of the deaths (43.75 per cent.) were in the third quarter of the year. Males, 5; females, 11; American parentage, 4; foreign, 12; total, 16, or 0.83 per cent. of all from known causes.

The statistics for the twenty-two years, 1856-1877 inclusive, are: -

Whole number, 328; annual average, 15.

Males, 139; females, 189; total, 328, or 1.17 per cent. of all.

American parentage, 129; foreign parentage, 199.

Hydrocephalus. The number of deaths from this cause in 1877, was 21; 1.09 per cent. of all from known causes. In 1876, the number was 45, or 2.43 per cent. The decedents in 1877 were 8 males; 13 females; 12 of American, and 9 of foreign parentage. The statistics for the twenty-two years, 1856–1877, are:—

Whole number, 659; annual average, 30.

Males, 362; females, 297; total, 659, or 2.36 per cent, of all.

American parentage, 333; foreign parentage, 326.

Insanity. The number of deaths from insanity in 1877, was 6; just half as many as in 1876. Number of males, 3; females, 3; American parentage, 4; foreign parentage, 2; total, 6, or 0.31 per cent. of all from known causes. For the twenty-two years, 1856-1877, the statistics are:—

Whole number, 219; annual average, 10.

Males, 109; females, 110; total, 219, or 0.78 per cent. of all.

American parentage, 159; foreign parentage, 60.

The decedents are very equally divided, as to sex; but very unequally as to parentage; 72.60 per cent. being of American parentage, and only 27.40 per cent. of foreign.

This fact, however, does not indicate the real proportions of insanity in the American and foreign population of the city, as the deaths reported mostly occur in the Butler Hospital for the Insane, in which, for obvious reasons, the insane population of American parentage is more fully represented than that of foreign parentage.

Intemperance. There were 8 deaths from this cause in 1877, of which 3 were from delirium tremens. Number of males, 5; females, 3; American parentage, 1; foreign, 7; total, 8, or 0.42 per cent. of all from known causes.

The following are the statistics for the twenty-one years, 1857-1877:—

Whole number from known causes, 26,948.
From intemperance, 213; annual average, 10.
Males, 155; females, 58; total, 213, or 0.79 per cent of all.
American parentage, 61; foreign parentage, 152.

Of the 213 decedents, 72; 77 per cent. are males; and 71.36 per cent. are of foreign parentage.

Kidneys, Diseases of, caused 49 deaths in 1877, 34 of which were from Bright's disease, and 3 from Addison's disease; males, 31; females, 18; American parentage, 24; foreign parentage, 25; total, 49, or 2.54 per cent. of all from known causes. The statistics for the twenty-two years, 1856–1877, are:

Whole number, 390; annual average, 18.

Males, 230; females, 160; total, 390, or 1.39 per cent. of all.

American parentage, 231; foreign parentage, 159.

Liver, Diseases of. The number of deaths from this cause in 1877, was 22, or 1.14 per cent. of all from known causes; males, 11; females, 11; American parentage, 12; foreign parentage, 10. For the twenty-two years, 1856-1877, the numbers are:

Whole number, 319; annual average, 14.

Males, 159; females, 160; total, 319, or 1.14 per cent. of all.

American parentage, 156; foreign parentage, 163.

Malformations. The number of deaths reported from this cause in 1877, was 16; just twice as many as in 1876, and 0.83 per cent. of all from known causes. Number of males, 8; females, 8; American parentage, 10; foreign, 6. There were 8 cases of malformation of the heart; 5, of the spine; 2, of the rectum; and one was not specified.

The following are the statistics of malformations for the twentytwo years, 1856-1877: —

> Males, 95; females, 57; total, 152, or 0.54 per cent. of all. American parentage, 104; foreign parentage, 48.

Marasmus was reported as the cause of 27 deaths in 1877; 6 less than in the preceding year, and 1.40 per cent. of all from known causes; males, 14; females, 13; American parentage, 10; foreign parentage, 17.

The statistics for the twenty-two years, 1856-1877, are as follows: -

Whole number, 549; annual average, 25.

Males, 291; females, 258; total, 549, or 1.96 per cent. of all.

American parentage, 282; foreign parentage, 267.

Measles. There were two deaths in Providence from measles in 1877; the first that have occurred since 1874; this disease is very irregular in the degree of its prevalence, as may be seen from the following table, which gives the number of deaths in each year since 1856:—

1856 1	1862 1	1868 4	18744
1857 2	186316	1869 6	18750
185840	1864 7	187016	18760
1859 0	1865 7	1871 1	18772
1860 0	1866 9	1872 7	-
1861 6	1867 0	187328	Total, 22 vrs 157

The two deaths in 1877 were 0.10 per cent. of all from known causes; one male, and one female; one of American, and one of foreign parentage. The statistics for the 22 years, 1856–1877, are:

Males, 75; females, 82; total, 157, or 0.56 per cent. of all.
American parentage, 44; foreign parentage, 113.

Meningitis, Cerebro-Spinal. The number of deaths from this cause in 1877, was 7; 3 more than in 1876, and 0.36 per cent. of all from known causes; males, 5; females, 2; American parentage, 4; foreign, 3. The first deaths from this cause, in Providence, were reported in 1864; the following list gives the number in each year since that time:—

1864 2	1869 0	187410
1865 2	1870 0	1875 6
1866 1	187112	1876 4
1867 0	187210	1877 7
1868 1	187339	Total, 14 years94

Whole number from known causes, 1864-1877, 20,139; from cerebro-spinal meningitis, 94.

Males, 51; females, 43; total, 94, or 0.47 per cent. of all. American parentage, 45; foreign parentage, 49.

The number and percentage of deaths in each division of age, and according to sex and parentage, were as follows, for the 14 years:—

AGM.	Under 1 Year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	Males.	Females.	American	Foreign.	Total.
No.	20	10	10	83	6	13	13	œ	9	1	1	51	84	40	67	8.
Pan cr.	5.82	5.32	10.64	24.47	9.08	13.83	13.83	8.51	8.38	1.06	1.06	24.26	45.74	19.18	52.13	100.00

Of these 94 decedents, 52, or 55.33 per cent., were under fifteen years of age.

Murder and Homicide. Only two deaths are returned under this heading, in 1877; 0.10 per cent. of all from known causes; one male, and one female; both of foreign parentage.

Neglect. One death was reported from neglect, in 1877; an infant, under one year old; a female of American parentage; 0.05 per cent. of all from known causes.

Old Age, was given as the cause of 63 deaths, in 1877; 9 less than in 1876, and 5 more than in 1875. Number of males, 24; females, 39; American parentage, 27; foreign, 36; total, 63, or 3.26 per cent. of all from known causes. Only 8 of the decedents lived beyond the age of 90 years; a smaller number than for some years previous. The following are the statistics for the twenty-two years, 1856–1877:—

Whole number, 1,129; annual average, 51.

Males, 371; females, 758; total, 1,129, or 4.04 per cent. of all.

American parentage, 697; foreign parentage, 432.

Of the whole number, 1,129, 67.14 per cent. are females; and 61.74 per cent. are of American parentage.

How small a proportion of those who live to advanced age are reported under this head may be seen from the following numbers.

During the year 1877, the number of persons who died at the age of 70 years or over, was 210; 126 of American, and 84 of foreign parentage; this was 10.87 per cent. of all the decedents from known causes.

In the twenty-two years, from 1856 to 1877 inclusive, the number of decedents of the age of 70 years and over, was 3,022, or 10.80 per cent. of all from known causes; of these, 2,147 were of American, and 875 of foreign parentage.

Paralysis. This is not, strictly speaking, a distinct disease, but, more properly, a symptom, of which the cause might often be specified more exactly than it is. But it was reported, in 1877, as the cause of 24 deaths; the same number as in 1876. Males, 13; females, 11; American parentage, 14; foreign parentage, 10; total, 24, or 1.24 per cent. of all from known causes. The statistics for the twenty-two years, 1856–1877, are:—

Whole number, 354; annual average, 16.
Males, 183; females, 171; total, 354, or 1.27 per cent. of all.
American parentage, 268; foreign parentage, 86.

If to these deaths are added those from apoplexy, we have the following statistics for each of the twenty-two years, 1856-1877:

Years.	Apoplexy.	Paralysis.	Total.	Years.	Apoplexy.	Paralysis.	Total.
1856	21	2	23	1868	22	24	46
1857	18	9	27	1869	32	16	48
1858	15	3	18	1870	28	24	52
1859	16	6	22	1871	33	28	61
1860	17	11	28	1872	31	20	51
1861	14	11	25	1873	36	21	57
1862	19	5	24	1874	36	23	59
1863	20	8	28	1875	41	35	76
1864	13	16	29	1876	44	24	68
1865	20	18	38	1877	50	24	74
1866	21	8	29				_
1867	31	18	49	To	tal, 22 years.		932

The statistics of both causes, as to sex and parentage, for this period, are: —

Whole number, 932; annual average, 42.

Males, 459; females, 473; total, 932, or 3.33 per cent. of all.

American parentage, 684; foreign parentage, 248.

It will be seen how large a proportion of these decedents are of American parentage; 73.39 per cent. being of this class, and only 26.61 per cent. of foreign.

Peritonitis. The number of deaths from this cause, in 1877, was 10; or 8 less than in 1876, 0.52 per cent. of all from known causes. Number of males, 3; of females, 7; of American parentage, 6; of foreign parentage, 4.

The statistics for the twenty-two years, 1856-1877, are: -

Whole number, 184; annual average, 8.

Males, 80; females, 104; total, 184, or 0.66 per cent. of all.

American parentage, 104; foreign parentage, 80.

Pneumonia. There were 88 deaths from this cause, in 1877; and 10 from congestion of the lungs; total, 98. In 1876, the deaths from these two causes, numbered 163; and in 1875, 197. The statistics as to sex and parentage in 1877, are as follows:—

Males, 44; females, 54; total, 98, or 5.07 per cent. of all. American parentage, 47; foreign parentage, 51.

The statistics for the twenty-two years, 1856-1877:

Whole number, 2,017; annual average, 92.

Males, 1,053; females, 964; total, 2,017, or 7.21 per cent. of all.

American parentage, 1,024; foreign parentage, 993.

Scarlatina. The number of deaths from this cause in 1877, was 12. Only twice in the twenty-three years, 1855-1877 inclusive, has a smaller number been reported; in 1866, the number was 3; and in 1867, it was 10. In 1877, the number of males was 4; females, 8; American parentage, 3; foreign, 9; total, 12, or 0.62 per cent. of all from known causes. For the twenty-two years, 1856-1877, the statistics are:—

Whole nnmber, 1,450; annual average, 66.

Males, 666; females, 754; total, 1,450, or 5.18 per cent. of all.

American parentage, 616; foreign parentage, 834.

The following table gives the particulars, as to sex, parentage, season, and locality, for each of the twenty-three years, 1855-1877, and for the entire period:—

SCARLATINA IN PROVIDENCE, TWENTY-THREE YEARS.

Total.	1,492	718	857	854448888FFF8	28855888888888
1877.	128	+00	00	co ; ot →	
1876.	1.87	19	75	400 40 1 1	4 :
1875.	4.91	68	50 40 40 40 40 40 40 40 40 40 40 40 40 40	はなるなるではなるないよい	Brodus は本でのは
1874.	268	55	191	862224255	408520222
1873.	25.7	82	25	数なってはいこうものの	第22日のの日日の
1872	1.20	×1	81	:00 05 00 to : : : : : : : : : : : : : : : : : :	10 H H H H H H
1871.	1.69	128	0.01	ØF-4-405-	H054H0005050000
1870.	188	198	281	Book ou him	H H40F0F
.1869.	138	1911	28.88	%221×441354	ಷ್ಕೆ ಪ್ರಾಪ್ತ ಪ್ರಾಪ್ತ ಪ್ರಾಪ್ತ ಪ್ರಾಪ್ತ ಪ್ರವಸ್ತೆ ಪ್ರವಸ್ತೆ ಪ್ರವಸ್ತೆ ಪ್ರಾಪ್ತ ಪ್ರವಸ್ತೆ ಪ್ರವಸ್ತೆ ಪ್ರವಸ್ತೆ ಪ್ರವಸ್ತೆ ಪ
1868.	009.	82	98	14の14の224442	数 :0-10+2500
1867.	100	-100	2-00	H92 HH HH02	+01- H
1866.	0.20	03	02 H		H 64 1 1
1865.	9.08	25.55	68	888日日の日の十一日二	**************************************
1864.	141	855	82		######################################
1863.	85.78	253	11	H00400H H : 10	∞∞∞00000 : : :
1868.	17	900	133		00 05 05 05 00 05
1861.	888	191	22	4	3000-000-
1860.	17.	101	19	8 H 988H H H	401- 01-00
1850.	45	22	127	***************	ಎ ವಬಾರ್ರವ ಕ
1858.	52.88	28	88	***********	504年288日
1857	3.58	13	18	жи жи	₹0001-40H : : !
1856.	14,00	775	120	######################################	81528333
1855	4.37	818	288		он б шижь ! ! !
BCARLATINA.	hole number	Males. Females.	ARENTAGE: American. Foreign.	Alanuary Pebruary March May May May May May May May August September October December	Ward II

Scrofula. The number of deaths from scrofula, in 1877, was 8; males, 6; females, 2; American parentage, 4; foreign parentage, 4; total, 8, or 0.41 per cent. of the whole number from known causes.

Small-Pox. There have been no deaths in Providence from this disease, since 1875. For the twenty-two years, 1856-1877, the general statistics are as follows:—

Whole number, 79, or 0.28 per cent. of all.

Males, 58; females, 21; American parentage, 48; foreign parentage, 31.

The particulars for each year are as follows:	:
---	---

		SEX.	Pare	TAGE.	Whole		
	Males.	Females.	American.	Foreign.		Percentage	
1856	7	2		4	9	0.87	
1857		• • • • • • • • • •			• • • • • • • • •		
1858						• • • • • •	
1859	2				5	0.5	
1860	4	1		2	5 .	0.5	
1861	. .				• • • • • • • • •		
1862	2	2	1	3	4	0.4	
1863					• • • • • • • • •		
1864	7		3	4	7	0.56	
1865	9	2	8	3	11	0.9	
1866					• • • • • • • • •		
1867	1		1		1	0.1	
1868	1		1	. 	1	0.0	
1869	3		3		3	0.2	
1870		2		2	2	0.1	
1871			1	2	3	0.2	
1872							
1873	12	2	8	6	14	0.8	
1874				-			
1875							
1876				· · · · · - · · · · ·			
1877							
			—	· · · · · · · · · · · · · · · · · · ·		· · · · · · · · · · · · · · · · · · ·	
22 years' tota	1 KQ	91	48	21	70	0.96	

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Spine, Disease of. The number of deaths from this cause in 1877, was 3, or 0.16 per cent. of all from known causes; 1 male, and 2 females; 1, of American, and 2, of foreign parentage.

Spleen, Disease of. Two deaths were reported from this cause in 1877; the first since 1873. There was one male decedent, and one female; one of American, and one of foreign parentage.

Stomach, Disease of. There were 20 deaths reported from this cause in 1877; 5 more than in 1876; and 1.03 per cent. of all from known causes; number of males, 6; females, 14; of American parentage, 9; foreign, 11.

Suicide. The number of suicides in 1877 was 12; this was 2 more than in 1876, and 0.62 per cent. of all from known causes; number of males, 8; females, 4; of American parentage, 7; of foreign parentage, 5. In 4 cases, it was by hanging; in 2 cases each, by laudanum and by pistol-shot; in one case each, by strychnine, by bichromate of potash, by drowning, and by jumping from the cars.

The following are the statistics for the twenty-two years, 1856-1877:—

Males, 94; females, 38; total, 132, or 0.47 per cent. of all. American parentage, 77; foreign parentage, 55.

Teething. The number of deaths reported from this cause in 1877, was 7; the same number as in 1876, and 0.36 per cent. of all from known causes; all were females, 3 of American, and 4 of foreign parentage.

Tuberculosis. This was returned as the cause of 20 deaths in 1877; 5 more than in 1876, and 1.03 per cent. of all from known causes; males, 7; females, 13; American parentage, 17; foreign parentage, 3.

If all tubercular diseases, such as consumption, hydrocephalus, and others, were united under this general head, it would present a much larger percentage of the whole number of deaths than any other cause.

Tumors. The number of deaths reported from this cause in 1877, was 4, just half as many as in 1876, and 0.21 per cent. of all from known causes. All the decedents were females; 3 of American, and 1 of foreign parentage. One was an abdominal tumor; one, uterine; and two, ovarian.

Unknown Causes. The number of deaths in 1877 for which no cause was given, was 6, or 0.31 per cent. of all deaths. Never has the number or percentage been so small before. The decedents were 2 males, and 4 females; 4, of American, and 2, of foreign parentage; 4 were children under one year old.

The statistics for the twenty-two years, 1856-1877, are as follows: —

Whole number, 508; annual average, 23.

Males, 265; females, 243; total, 508, or 1.78 per cent. of all.

American parentage, 166; fereign parentage, 342.

Of these 508 decedents, 67.32 per cent. were of foreign parentage.

STILL-BORN.

The following are the usual statistics in regard to this class, for the year 1877: —

Number. There were 145 still-born children in Providence in 1877; 4 less than in 1876. Of the 2,543 children born during the year, the still-born were 5.70 per cent.; or 1 in 17.54.

Sex. Males, 86; females, 59; 6 more males, and 10 less females than in 1876.

Color. White, 134; colored, 11.

Locality. First ward, 18; second, 9; third, 23; fourth, 3; fifth, 12; sixth, 8; seventh, 16; eighth, 19; ninth, 18; tenth, 19.

Season. January, 15; February, 7; March, 8; April, 19; May, 9; June, 6; July, 11; August, 12; September, 12; October, 11; November, 13; December, 22.

The number and percentage in each quarter of the year were as follows: —

Number. Per	cent.	Number.	Per cent.
1st quarter3020	.69 3d	quarter35	24.14
2d quarter3423	3.45 4t	h quarter46	31.72

Parentage. In the following table, the first letter indicates the birthplace of the father; the second, that of the mother.

A. and A6	2 I. and A	A. and S
I. and I4	7 G. and A	I. and S
E. and E	4 B. A. and A	A. and B. A
S. and S	1 A. and I	I. and B. A
F. and F	1 B. A. and I	E. and B. A
B. A. and B. A	3 A. and E	A. and D
	1 I. and E	
P. and P	1 B. A. and E	Total14

A.—American; I.—Irish; E.—English; S.—Scotch; G.—German; B. A.—British American; It.—Italian; F.—French; D.—Dutch; P.—Portuguese.

Of the 2,543 children, born in Providence in 1877, 145, or 5.70 per cent., were still-born. Of these 2,543 children, 1,292, or 50.80 per cent., had American mothers; 714, or 28.08 per cent., had Irish mothers; and 537, or 21.12 per cent., had mothers who were natives of other foreign countries. Of the still-born, the proportions are as follows to the mothers of the different classes: American, 71, or 48.97 per cent.; Irish, 51, or 35.17 per cent.; other foreign, 23, or 15.86 per cent.

From these numbers, it might appear that the largest percentage of the still-born were children of American mothers; but though they are more in number, the proportion of still-born to the whole number of children born to mothers of each class, is much larger among the Irish; as may be seen by making the comparison in the following manner.

Of the 1,292 children of American mothers, 71, or 5.50 per cent., were still-born; of the 714 children of Irish mothers, 51, or 7.14 per cent.; and of the 537 children of other foreign mothers, 23, or 4.28 per cent.

The proportions are: to American mothers, one in 18.20; to Irish mothers, one in 14.00; to other foreign mothers, one in 23.35.

These numbers compare with those of the previous year, as follows: the number and proportion of children born to American mothers were larger in 1877, than in 1876; the number of still-born slightly larger, but the proportion to the whole number less than in the preceding year. Of children of Irish mothers, the number and proportion was less than in 1876; of still-born, the number was less, but the percentage larger. Of children of other foreign mothers, the number and proportion was larger than in 1876; of still-born, the number was a little larger and the percentage a little less than in the preceding year. In the twenty-two years, 1856-1877; there have been born in Providence 41,273 children, of whom 2,464, or one in 16.75, have been stillborn; 5.97 per cent. of the whole number. Of the 2,464 still-born children, 2,311 were white, and 153 were colored; number of males, 1,420, or 57.63 per cent.; females, 1,044, or 42.37 per cent. Reckoning the parentage according to the father's birthplace, 1,019 were American; 1,129, Irish; 309, other foreign; and 7 were of unknown parentage.

GENERAL SUMMARY.

The following recapitulatory table will give the general results of registration in Providence during the 23 years, 1855 to 1877, inclusive:—

TABLE XXI. BIRTHS, MARRIAGES, AND DEATHS.

YEAR.	Population.	Births.	To population, one birth in	Marriages.	Of population, one person married in	Deaths.	Of population, one death in
1855	47,785 48,727 49,457 49,457 50,100 50,666 50,666 52,238 53,024 53,810 54,595 55,709 56,824 64,138	1,600 1,675 1,688 1,724 1,593 1,648 1,725 1,529 1,307 1,344 1,298 1,632 1,632 1,624 1,866	29.86 29.09 29.30 28.68 31.45 30.74 29.37 34.16 40.56 40.04 42.06 34.13 34.99 34.37	636 656 614 560 616 633 547 547 610 739 709 812 855 872	37.56 37.14 40.27 44.15 40.66 40.02 46.31 47.75 43.46 36.40 38.50 34.30 33.23 36.77	991 1,065 925 1,017 899 1,001 1,051 914 1,214 1,281 1,211 1,036 960 1,110	48.22 45.75 53.46 48.63 55.72 50.61 48.21 57.15 43.67 42.01 45.08 53.77 59.19 57.78
1869. 1870. 1871. 1872. 1873. 1874. 1875. 1876. 1877. Average.	66,522 68,904 72,011 75,118 78,225 96,452 100,675 101,500 100,675	1,819 1,889 1,960 2,206 2,128 2,780 2,664 2,631 2,543	36.57 36.48 36.74 34.05 36.76 34.69 37.79 38.58 39.58	930 959 943 1,084 1,150 1,176 1,079 949 953	35.76 35.93 38.13 34.63 34.01 41.01 46.65 53.48 52.82	1,256 1,263 1,254 1,603 1,719 1,987 1,915 1,865 1,938	52.96 54.55 57.42 46.86 45.51 48.54 52.57 54.42 51.95

BIRTHS AND DEATHS BY PARENTAGE.

The following recapitulatory table shows the number of births and deaths, by parentage, in Providence, in each of the twenty-two years from 1856–1877 inclusive. In the three years, 1863, 1864, and 1865, there was an excess of deaths over the births, in the population of American parentage.

TABLE XXII.

YEAR.	AMERI	CAN PARE	NTAGE.	FOREIGN PARENTA			
	Deaths.	Births.	Excess of Births.	Deaths.	Births.	Excess of Births	
1856	541	759	218	524	916	392	
1857	523	701	178	402	987	585	
1858	512	771	259	505	953	448	
1859	479	659	180	420	934	514	
1860	512	731	219	489	917	428	
1861	538	740	202	513	985	472	
1862	460	663	203	454	866	412	
1863	597	528	69	617	779	162	
1864	649	579	—70 l	632	765	133	
1865	631	546	-85	580	752	172	
866	513	726	213	523	906	383	
867	528	725	197	432	899	467	
1868	596	882	286	514	984	470	
1869	627	926	299	629	893	264	
870	642	890	248	621	999	378	
871	643	980	337	611	980	369	
872	832	1,035	203	771	1,171	400	
873	862	1,003	141	857	1,125	268	
874	885	1,217	332	1,102	1,563	461	
875	914	1,180	266	1,001	1,484	483	
876	875	1,169	294	990	1,462	472	
.877	908	1,238	330	1,030	1,305	275	
22 years	14,267	18,648	4,381	14,217	22,625	8,408	

CITY ORDINANCES. - CHAPTER XLVII.

REGISTRATION OF DEATHS.

SECTION 1. There shall be appointed by the City Council a sufficient number of persons to act as undertakers, removable at the pleasure of the City Council.

SECT. 2. Whenever any person shall die in the city, the physician attending in his or her last sickness shall furnish to the undertaker attending the funeral, or to the city registrar, a certificate, giving the name of the person, date of death, and the disease or cause of his or her death.

SECT. 3. No person shall bury, or place in the tomb, or remove from the city, or otherwise dispose of the body of any human being who shall die in the city, without first reporting the death to the city registrar, and obtaining a permit from him.

from him.

Sect. 4. No permit shall be given, as provided in section three, until the city registrar is furnished with the information in relation to the deceased person, required by the laws of the State, for record, so far as the same can be ascertained, together with the physician's certificate of the cause of death, whenever a physician has been in attendance, or a coroner's certificate, whenever a coroner's inquest has been held.

Whenever a permit for burial is applied for, in a case of death without the attendance of a physician, or if it is impossible to obtain the physician's certificate, the city registrar shall investigate the case so far as may be necessary, and when he has obtained satisfactory evidence in relation to the cause and circumstances of the death, he shall sign the certificate and give the required permit. If not satisfied in relation to the cause and circumstances of the death, or if, in his opinion, the public good requires it, he shall report the case to a coroner for investigation. investigation.

SECT. 5. Whenever the body of a human being who has died out of the city shall be brought here for burial, the undertaker, or other person attending the funeral, shall furnish the report required in sections three and four, with the exception of the physician's certificate.

SECT. 6. All funerals shall take place between sunrise and sunset, unless otherwise permitted or directed by the board of health. The top of every coffin deposited in the ground shall be at least three feet below the usual surface

The city registrar shall cause an abstract of the returns of deaths made to him to be published in all the newspapers which contract to do the city printing, monthly, and oftener if required by the board of health. He shall also, on or before the first day of April, annually, prepare and present to the city council a statement of the number of births, marriages and deaths which occurred in the city during the year ending with the thirty-first day of December next preceding, with such other information and suggestions in relation thereto as he may deem useful for the promotion of the public health, and other interests of the city.

SECT. 8. No undertaker or other person shall bury or cause to be buried the body of any deceased person in the city, except in such grounds as are or may be designated as burying-grounds, and authorized to be used as such.

SECT. 9. Every person violating any provision of this chapter shall pay a fine of not less than five and not more than twenty dollars for each offence.

} j i i 1 Vital Statistics, Providence, R. I.

1877.

CITY REGISTRAR'S

TWENTY-THIRD ANNUAL REPORT.

1870.7

CITY DOCUMENT.

[No. 24.

TWENTY-FOURTH ANNUAL REPORT

THOS THE

BIRTHS, MARRIAGES, AND DEATHS

IS IND

CITY OF PROVIDENCE,

TOR THE KANS.

1878,

By EDWIN M. SNOW, M. D.,

SUPPLIES FOR THE ABBADTO AND THEY BRADTELAN.



PROVIDENCE:

PROTERNER PRESS 190, PRINTERS TO THE CITY-1870.



1879.]

CITY DOCUMENT.

[No. 24.

TWENTY-FOURTH ANNUAL REPORT

UPON THE

BIRTHS, MARRIAGES, AND DEATHS

IN THE

CITY OF PROVIDENCE,

FOR THE YEAR

1878,

By EDWIN M. SNOW, M. D.,

SUPERINTENDENT OF HEALTH AND CITY REGISTRAR.



PROVIDENCE:

PROVIDENCE PRESS CO., PRINTERS TO THE CITY. 1879.

THE CITY OF PROVIDENCE.

JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved July 2, 1879.]

RESOLVED, That the twenty-fourth annual report of the City Registrar, for the year ending December 31, 1878, be received, and that six hundred copies thereof, and of the communication accompanying the same, be printed in pamphlet form for the use of the City Council, under direction of the joint committee on printing.

A true copy: witness,

H. V. A. JOSLIN, CITY CLERK.

INTRODUCTION.

CITY REGISTRAR'S OFFICE, PROVIDENCE, June 2, 1879.

To the Honorable the City Council:

Enclosed is my Twenty-Fourth Annual Report upon the Births, Marriages and Deaths, in the City of Providence. The statistics contained in it are for the year ending December 31, 1878. During the year 1878, there were 2,627 children born in Providence; 2,032 persons were married, and 1,989 persons died. As compared with the year 1877, the number of births in 1878 was 84 more; of persons married, 126 more; and of deaths, 51 more.

A low estimate of the population of Providence, in the middle of the year 1878, is 101,000. This gives a ratio of births, marriages and deaths to population as follows:—

One birth in 38.44, or 26 01 in each 1,000.

One person married in 49.70, or 20.12 in each 1,000.

One death in 50.78, or 19.69 in each 1,000.

The average annual rate of mortality in Providence during the 24 years, from 1855 to 1878 inclusive, has been less than 20 in each 1,000 of the population. This rate is very small for a city of the size of Providence, and is based upon full returns of deaths and accurate censuses of population.

Full particulars relating to the causes of death in 1878, together with the usual and additional tables in relation to births, marriages and deaths, are given in the report.

EDWIN M. SNOW, M. D.,
Superintendent of Health and City Registrar.

CONTENTS.

The state of the s	PAGE
BIRTHS	1-1
Wards, months, sex and color, (Table I.). Births in connection with locality, sex and season, (Table II.) Proportion of the sexes born at different seasons.	
Births in connection with locality, sex and season, (Table II.)	1
Proportion of the sexes born at different seasons	-
Number and proportion of the sexes, 25 years. Color:—Births, marriages and deaths in 23 years	
Color :—Births, marriages and deaths in 23 years	4-
Parentage of children born in each ward	4.3
Number and percentage of American and of foreign children, 23 years	6-
Parentage:—Mixture of different nations, (Table IV.). Number of 1st, 2d, etc., children in 1878, and 1855-1877, (Table V.). Average number of children to each mother, 24 years, (Table VI.). Number of the child, with age of the mother, (Table VII.).	3
Number of 1st, 2d, etc., children in 1878, and 1855-1877, (Table V.)	3
Average number of children to each mother, 24 years, (Table VI.)	40 1
Number of the child, with age of the mother, (Table VII.)	10-1
Plurality-births, statistics for 1878	13
Fluranty-birens, 21 years summary	13-16
MARRIAGES	16-23
Marriages at different seasons, 24 years, (Table VIII.)	16-17
Birthplaces of parties married in 1878, (Table IX.)	18
Bride and groom born in same country, etc.	18
Proportions of American, foreign and mixed marriages, 24 years	19
Marriages of colored persons. Ages of persons married in 1878. (Table X)	20
Average age of persons married, 23 years, (Table)	21
Number of times married, (Table XI.)	22
Number of times married, (Table XI.) Denominational. Nativity and residence	22
Marriage and education; signed with a mark	23
Decision	
Deaths to population in 1878; proportion in each class Deaths by sex, condition, color, nativity, parentage, etc., (Table XII.)	24-72
Deaths by sey condition acles proportion in each class	25
Deaths by season; numbers and proportions; recapitulation	26-27
Sex, condition, color, nativity, parentage and locality of decedents, 1878	27-29
A agreement and average are by parentings and investigation vitte	29
Aggregate and average age by parentage, 1878, (Table XIII.)	30-32
Difference between average age of American and foreign decedents, (Table).	31
Percentages at different ages by parentage 1878, Table (XVI)	33
Percentages at different ages by parentage, 1878, Table (XVI.)	34
Percentages at different ages by parentage, 23 years (Table XVII)	35
Deaths by age and parentage in each month and quarter, 1878, (Table XVIII)	36-38
Causes of death, sex, age, parentage and percentages, (Table XIX.)	39-41
Causes of death in connection with season, (Table XX.)	40
Notes on the Causes of Death in 1878	43-70
Cholera Infantum; decedents by sex and parentage, 23 years	40
Consumption; sex and parentage; number of deaths at each age	47-48
Diarrhea and Dysentery; recapitulation for 39 years	49-51
Diarrhoa and Dysentery; recapitulation for 39 years. Diphtheria; statistics for 1878. Diphtheria in Providence, 21 years, (Table). Diphtheria; season and mortality, compared with scarlatina and cronp,	52
Diputheria in Providence, 21 years, (Table)	53
Diputieria; season and mortanty, compared with scariatina and croup,	
(Tables). Diphtheria, epidemic and not epidemic, compared, (Table) Heart, diseases of; decedents by sex and parentage, 23 years, (Table)	54-55
Dipotneria, epidemic and not epidemic, compared, (Table)	56-57 58-59
Heart, diseases of, decedents by sex and parentage, 25 years, (1401e)	59-60
Hooping Cough	62-63
Meningitis, cerebro-spinal; deaths at each age, 15 years	63-64
Scarlatina; statistics for 1878	68
Searlating in Providence 24 years (Table)	67
Small-pox, recapitulatory table, 23 years.	69
Still-born children in 1878	71
Still-born children, 1856–1878	72
Diff-Dota Children and Colored	
GENERAL SUMMARY	73
Births, Marriages and Deaths; numbers and proportions, 24 years, (Table XXI.	73
Births and Deaths by parentage; excess of births, 23 years, (Table XXII.)	74
Come Opposition Del amiso do Project Lanor	95

CITY REGISTRAR'S REPORT.

1878.

BIRTHS.

THE number of children born in the city of Providence during the year 1878, was 2,627.

In Table I. will be found the number born in each ward, in each month and quarter of the year; and the number of each sex, and the number of colored children born in each month and quarter, for the whole city.

TABLE I. BIRTHS, 1878. Wards, Months, Sex, and Color.

PROVIDENCE.					WA	RDS					wı	nole (City.	chil	olore dren ed in cedin	in-
1878.	L	11.	ш,	ıv.	v.	VI.	VII.	viii	ıx.	x.	Males.	Females.	Monthly Totals.	м.	F.	Totals.
January February March	25 25 27	8 4 3	16 22 23	. 13 6 8	14 14 17	8 7 10	24 18 20	29 20 27	26 29 41	50 44 37	128 95 106	88 94 117	216 189 223	4	8 3 4	10 7 6
1st Quarter	87	15	61	27	45	25	62	76	96	134	329	299	628	8	15	23
April	28 20 24	6 1 5	23	11 14 9	11 18 20	8 7 7	19 13 16	30 22 19	38 38 34	46 36 48	122 97 104	100 95 102		8 4 3	4 5 5	12 9 8
2d Quarter	72	12	72	34	49	22	48	71	110	130	323	297	620	15	14	29
July	28 27 34	5 6 4	20 23 21	10 14 10	17 19 26	15 12 7	19 19 17	27 28 17	32 42 45	35 46 62	106 117 133	102 119 110		10	4 4 5	14 8
3d Quarter	89	15	64	34	62	34	55	72	119	143	356	331	687	13	13	26
October November December	28 34 23	8 4 2	24 28 21	8 7 12	21 15 20	12 8 6	21	32 25 33	36 39 40	55 54 37	123 115 112	115 120 107		7	3 3	10
4th Quarter	85	14	73	27	56	26	60	90	115	146	350	342	692	16	10	26
Whole year	333	56	270	122	212	107	225	309	440	553	1,358	1,269	2,627	52	52	104

The number of children born during the year 1878 was 84 more than in the preceding year; of males, 56 more, and of females, 28 more. The first, second, sixth, and ninth wards, collectively, show a loss of 83 in the number of births, and the remaining wards, a gain of 167. In the first quarter of the year there were 56 more births, in the year 1878, than in the corresponding season of 1877; in the second quarter, 54 more; in the third quarter, 12 more; and in the fourth quarter, 38 less. Estimating the population of the city at 101,000 for the middle of the year 1878, it gives a ratio of births to population of one in 38.45; or 26.01 in each 1,000 of the population.

Season. The largest number of births in any one month in 1878, was 243 in September; the smallest, 189 in February. The largest quarterly total was 692 in the fourth quarter; and the smallest, 620 in the second. The number of births in the first six months, was 1,248; in the second six months, 1,379, a difference of 131; in the first half of the year, 110 more than at the same season of the preceding year; and in the second half, 26 less.

The following statement is a comparison of the births in each quarter of the year, in the aggregate for twenty-two years, 1856-1877, inclusive, with the numbers for 1878; also the aggregate for the twenty-three years, 1856-1878:—

	Jan. March.	April. June.	First 6 mos.	July. Sept.	Oct. Dec.	Second 6 mos.	Whole Year.
Twenty-two years	10,094	10,286	20,380	10,191	10,702.	20,893	41,273
1878	628	620.	1,248	687	692.	1,379	2,627
Twenty-three years	10,722	10,906.	21,628 .	10,878	11,394.	22,272	43,900

The following are the monthly totals for the period of twenty-four years, from 1855-1878; the daily average of births in each month is also given, and the daily average for the whole period:—

Births in	January.	February.	March.	April.	May.	June.
24 years, 1855-1878	3,585	8,481	4,013	3,514	3,768	3,987
Daily average	4.82	5.13	5.39	4.88	5.06	5.54
Births in	July.	August.	Sept.	Oct.	Nov.	Dec.
24 years, 1855-1878	3,729	3,822	3,742	3,905	3,879	4,075
Daily average						
Whole number of birth	ıs in 24 years, 40	5,500; daily s	ver age, 5.	19.		

The largest daily average for the whole period is in June; the smallest in March. There were six months in which the daily average was greater than that for the whole period, viz.: March, June, September, October, November, and December.

Sex and Locality. The number of children of each sex, born in each ward of the city, in each quarter of the year 1878, is given in Table II.; also the number of each sex, for the whole year:—

	1st	Qua	rter.	2d	Qua	rter.	3d	Qua	rter.	4th	Que	arter.	Wh	ole Y	ear.
WARDS.	м.	p.	Total.	м.	F.	Total.	M.	F.	Total.	M.	F.	Total.	м.	F.	Total.
ı,	42	45	87	43	29	72	53	36		45	40	85 14	183	150	333
III	22	39	15 61	35	37	12 72	10 27	37	15 64	35	38	73	119	151	270
V	11 20	16 25	27 45	20 25	14 24	49	22 27	12 35	62	15 28	12 28	56	68 100	112	215
VI	12 32	13 30		11 25	23	92 48	20 28	27		28 20 24	36		109	116	
VIII	42 54	34 42		34 62	37 48	71	38 63	34		47 57	43		161 236	148	309
X	85	40		62			68	75		72			287	266	
Whole City	320	290	628	323	207	620	356	331	687	350	342	692	1,358	1,260	2,627

TABLE II. BIRTHS, 1878. Wards, Sex, and Season.

Proportion of the Sexes. Of the 2.627 children, born in Providence, in the year 1878, the number and proportion of each sex were as follows: Males, 1,358; females, 1,269; 107 males to each 100 females; or 51.69 males, and 48.31 females in each 100 children born. This proportion of males is larger than in either of the two years preceding, and greater than the average for 25 years.

The following table shows the number and proportion of each sex in each quarter of the year 1878:—

	Males.	Fema	es.	Males	.]	Fema	les.	Male	s. 1	Females.
JanMarch	329 a	nd 299,	or	110.0	to	100,	or	52.39	to	47.61 in each 100 born
April-June	. 323 a	nd 297,	or	108.7	to	100,	or	52.10	to	47.90 in each 100 born
July-Sept	. 356 a	nd 331,	or	107.6	to	100,	or	51.82	to	48.18 in each 100 born
OctDec,	350 aı	nd 342,	or	102.3	to	100,	or	50.58	to	49.42 in each 100 born
Whole Year	.1,358 a	nd 1,260,	or	107.0	to	100,	or	51.69	to	48.31 in each 100 born

In the aggregate for twenty-three years, 1856-1878 inclusive, the number and proportion of each sex, born in each quarter of the year, were as follows:—

Males	. 1	emales	. 1	Tales.	F	emal	es.	Males	. F	emales.			
JanMarch5,543	and	5,179,	or	107.0	to	100,	or	51.70	to	48.30 in	each	100	born.
April-June5,637	and	5,260,	or	106.9	to	100,	or	51.69	to	48.31 in	each	100	born.
July-Sept5,511	and	5,367,	or	102.7	to	100,	or	50.66	to	49.34 in	each	100	born.
OctDec5,862	and	5,532,	or	106.0	to	100,	or	51.45	to	48.55 in	each	100	born.
Whole Year22,553	and	21,347,	or	105.6	to	100,	or	51.31	to	48.69 in	each	100	born.

In the twenty-five years, 1854-1878 inclusive, there were 46,869 children born in Providence; the number and proportion of each sex in each year, were as follows:—

		_		_	_	_		_			_		_	
	Males		Female	es.	Males.	F	emal	es.	Males	. Fe	emales.			
1854	711 a	nd	658;	or	108.0	to	100;	or	51.91	and	48.09 in	each	100	born.
1855	807 a	nd	793;	or	101.7	to	100;	or	50.41	and	49.56 in	each	100	born.
1856	801 a	nd	784;	or	113.6	to	100;	or	53.19	and	46.81 in	each	100	born.
1857	872 a	nd	816;	or	106.8	to	100;	or	51.66	and	48.34 in	each	100	born.
1858	890 a	nd	834;	or	106.7	to	100;	or	51.62	and	48.38 in	each	100	born.
1859	825 a	nd	768;	or	107.4	to	100;	or	51.79	and	48.21 in	each	100	born.
1860	809 a	nd	839;	or	96.4	to	100;	or	49.09	and	50.91 in	each	100	born.
1861	936 a	nd	789;	or	118.6	to	100;	or	54.26	and	45.74 in	each	100	born.
1862	791 a	nd	738;	or	107.2	to	100;	or	51.73	and	48.27 in	each	100	born.
1863	689 a	nd	618;	or	111.4	to	100;	or	52.71	and	47.29 in	each	100	born.
1864	·663 a	nd	681;	or	97.3	to	100;	or	49.33	and	50.67 in	each	100	born.
1865	691 a	nd	607;	or	113.8	to	100;	or	53.24	and	46.76 in	each	100	born.
1866	849 n	nd	783;	or	108.4	to	100;	or	52.02	and	47.98 in	each	100	born-
1867	830 n	ind	704;	or	104.5	to	100;	or	51.11	and	48.89 in	each	100	born.
1868	944 a	nd	922;	or	102.4	to	100;	or	50.59	and	49.41 in	each	100	born.
1869	945 n	nd	874;	or	108.1	to	100;	or	51.95	and	48.05 in	each	100	born.
1870	967 a	and	922;	or	104.8	to	100;	or	51.19	and	48.81 in	each	100	born.
1871	956 a	ind	1,004;	or	95.6	to	100;	or	48.78	and	51.22 in	each	100	born.
1872	1,079 a	and	1,127;	or	95.7	to	100;	or	48.91	and	51.09 in	each	100	born.
1873	1,109 a	ind	1,019:	or	108.8	to	100;	or	52.11	and	47.89 in	each	100	born.
1874	1,408 п	nd	1,372;	or	102.6	to	100;	or	50.65	and	49.35 in	each	100	born.
1875	1,390 a	ind	1,274;	or	109.1	to	100;	or	52.18	and	47.82 in	each	100	born.
1876	1,359 8	und	1,272;	or	106.8	to	100;	or	51.65	and	48.35 in	each	100	born.
1877	1,302 m	and	1,241;	or	104.9	to	100;	or	51.20	and	48.80 in	each	100	born.
1878	1,358 #	ind	1,269;	or	107.0	to	100;	or	51.09	and	45.31 in	each	100	born.
-	_		-		_		_		_					
25 years2	2,071 8	nd	22,798;	or	105.6	to	100;	or	01.36	and	48.64 11	each	100	born.

Color. There were 104 colored children born in Providence in the year 1878; this was the same number as in the preceding year; but the proportion of the sexes is contrary to the general rule; the number of males and females in 1878 being the same. These births were in the different wards, as follows: First ward, 20; second, 8; third, 8; fourth, 5; fifth, 1; sixth, 11; seventh, 11; eighth, 12; ninth, 28; tenth, none. There were 10 births of colored children in January; 7 in February; 6 in March; 12 in April; 9 in May; 8 in June; 4 in July; 14 in August; 8 in September; 9 in October; 10 in November, and 7 in December.

In 23 years, 1856-1878 inclusive, the number of colored children born in Providence was 1,699; 901 males, and 798 females; a proportion of 112.9 males to each 100 females; or 53.03 males, and 46.97 females in each 100 children born.

The following are the statistics of births, marriages and deaths among the colored population of the city, for the 23 years, 1856-1878:—

Twenty-two years, 1856-1877	95 births774 marriages1,6	81 deaths.
1878	H births 54 marriages 1	06 deaths.
Twenty-three years' total	99 births	787 deaths.

The proportion of births to deaths among this class in 1878, was 100 to 101.9; in the twenty-three years, 1856-1878, it was 100 to 105.2.

PARENTAGE.

The facts on this important subject, given in the next few pages show the relative proportions of the different classes of our population.

Table III. gives the number of children, according to parentage, born in each ward; also the percentage of each nationality for the whole city. When the parents are of different foreign countries the parentage is always reckoned according to the birthplace of the father:—

TABLE III.	Births, 1878.	Wards and	Parentage.

				1	VAI	RDS.					City.	100 ere
PARENTAGE.	1.	11.	m.	ıv.	v.	vı.	vii.	viii	ıx.	x.	Whole C	In each born th were
American	118	40	75	66	62	62	106	164	210	132	1,035	39,40
Irish English, Scotch and Welsh German British American Portuguese Other Foreign	11		110 7 7 8 24 1	7 1 2	75 16 10 9 1	2	63 7 3 6 2 8	64 21 1 9	104 40 13 7 2 10	201 60 21 21 21 2 6	779 177 67 75 31 45	29.65 6.74 2.55 2.86 1.18 1.71
Total Foreign	27		157	34	116	1	89 13	97 25	176 26	311 43	1,174	44.69 6.85
mother	31	2	21	17	23	0	17	23	28	67	238	9.00
Ward totals	333	56	270	122	212	107	225	309	440	553	2,627	100.00

The "other foreign" in this table were as follows: Italian, 13; Swedish, 12; French, 6; Polish, 6; Norwegian, 2; Swiss, 2: and one each of Russian, African, Danish, and Belgian parentage.

The following is a recapitulation, showing the number and percentage of children, born of American, foreign and mixed parentage in each of the twenty-three years, 1856-1878 inclusive; also the same for the entire period:—

Year.		merican. . Per cent.	Forei Number.	gn. Per cent.	Amer. fat For. moth Number. Per	ier.	Amer. m	other.
1856					55			3.52
1857	650.	38.51	937	55.51	51	3.02	50	2.96
1858	712.	41.30	886	51.39	58	3.36	68	3.95
1859	599.	37.60	887	55.69	59	3.70	48	3.01
1860	661.	40.11	857	52.00	69	4.19	61	3.70
1861	654.	37.91	916	53.10	86	4 99	69	4.00
1862	589.	38.52	799	52.26	74	4.84	67	4.38
1863	467.	35.73	711	54.40	62	4.74	67	5.13
1864	520.	38.69	609 .	52.00	60	4.47	65	4.8
1865	476.	36.67.	684	52.70	70	5.39	68	5.24
1866	656.	40.20	821	50.30	70	4.29	85	5.21
1867	645.	39.72	791	48.70	80	4 93	108	6.68
1868	779.	41.75	868	46.51	103	5.52	116	6.25
		45.19			104			
1870	800	42.35	862	45.64	90	4.76	137	7.2
1871					101			
1872					128			
1873					134			
					149			
					181			
					162			
					184			
					180			
23 years' tot	als.17,553	30.98	21,175	48.24	2,310	5.26	2,862	6.5

These percentages vary slightly from year to year; there is usually a much larger percentage of foreign than of American parentage; the years 1869 and 1871 being the only ones in which the American percentage exceeds the foreign; in 1878 the percentage of foreign parentage is considerably larger than in 1877, and all the other classes show a corresponding decrease. The percentage of both classes of mixed parentage, though smaller than in the preceding year, is yet above the average for the whole period.

The following table shows the percentage of children born in Providence, of parents of each nation, for each of the twenty-three years, 1856-1878:—

Per cent. Pe 1856 42.09 2 1857 38.51 2 1858 41.30 2 1859 37.60 5 1860 40.11 5 1861 37.91 2 1862 38.52 5 1863 35.73 2 1864 38.69 6 1865 36.67 4 1866 40.20 3 1867 39.72 1 1868 41.75 1 1869 45.19 3 1870 42.35 3 1871 44.85 5 1872 41.12 3 1873 40.84 36.69	r cent. Per cent 11.79 4.84. 14.32 5.92 12.98 4.82. 5.58 5.34 10.96 6.55 12.61 6.03 11.53 6.02 13.08 6.35	German. t. Per cent. 2.27. 3.73 2.26. 3.20. 2.86 2.49 3.07. 2.98. 2.98.	Per cent.	
1856 42.09 1857 38.51 1858 41.30 1859 37.60 1860 40.11 1861 37.91 1862 38.52 1863 35.73 1864 38.69 1865 36.67 1866 40.20 1868 41.75 1869 45.19 1871 44.85 1872 41.12 1873 40.84	11.79	2.27		
1857 38.51 1858 41.30 1859 37.60 1860 40.11 1861 37.91 1862 38.52 1863 35.73 1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1870 42.35 1871 44.85 1872 41.12 1873 40.84	44.32 5.92 12.98 4.82 5.58 5.34 10.96 6.55 12.61 6.03 11.53 6.02 13.08 6.35	3.732.263.262.862.493.072.982.75		
1858 41.30 1859 37.60 1860 40.11 1861 37.91 1862 38.52 1863 35.73 1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1870 42.35 1871 44.85 1872 41.12 1873 40.84	12.98 4.82 5.58 5.34 10.96 6.55 12.61 6.03 11.53 6.02 13.08 6.35	2.263.262.852.493.072.982.75		
1859 37.60 .4 1860 40.11 .4 1861 37.91 .4 1862 38.52 .4 1863 35.73 .4 1864 38.69 .4 1865 36.67 .4 1866 40.20 .5 1867 39.72 .5 1869 45.19 .5 1870 42.35 .5 1871 44.86 .5 1872 41.12 .5 1873 40.84 .5	.5.585.346.55	3.262.852.493.07	····-	
1860 40.11 41861 37.91 41861 37.91 41862 38.52 41863 36.73 48.69 46.89 4	0.966.55 2.616.03 1.536.02 3.086.35	2.85 2.49 3.07 2.98 2.75	·····-	—1.64 —1.97 —1.04 —1.99
1861 37.91 1862 38.52 1863 35.73 1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1870 42.35 1871 44.85 1872 41.12 1873 40.84	2.616.03 11.536.02 3.086.35	2.49 3.07 2.98 2.75	·······	1.97 1.04 1.99
1861 37.91 1862 38.52 1863 35.73 1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1870 42.35 1871 44.85 1872 41.12 1873 40.84	2.616.03 11.536.02 3.086.35	2.49 3.07 2.98 2.75	·······	1.97 1.04 1.99
1862 38.52 1863 35.73 1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1869 45.19 1871 44.85 1872 41.12 1873 40.84	1.536.02 3.086.35	3.07 2.98 2.75	·····-	1.04
1863 35.73 1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1870 42.35 1871 44.85 1872 41.12 1873 40.84	3.086.35	2.98	·····—···	1.99
1864 38.69 1865 36.67 1866 40.20 1867 39.72 1868 41.75 1869 45.19 1870 42.35 1871 44.85 1872 41.12 1873 40.84		2.75		
1865 36.67 4 1866 40.20 3 1867 39.72 3 1868 41.75 3 1870 42.35 3 1871 44.85 3 1872 41.12 3 1873 40.84 3				
1866. 40.20. 1867. 39.72. 1868. 41.75. 1869. 45.19. 1870. 42.35. 1871. 44.85. 1872. 41.12. 1873. 40.84.				
1867 39.72 1868 41.75 1869 45.19 1870 42.35 1871 44.85 1872 41.12 1873 40.84				
1868 41.75 1869 45.19 1870 42.35 1871 44.85 1872 41.12 1873 40.84 11.2 1873 40.84 11.2 1873 40.84 11.75 1874 11.75 1875 1875 1875 1875 1875 1875 1875 18				
1869 45.19 1870 42.35 1871 44.85 1872 41.12 1873 40.84				
1870 42.35 3 1871 44.85 3 1872 41.12 3 1873 40.84 3	•••••			1.07
1871				
1872 41.12				
1873 40.84				
	2.465.67	2.77	2.31	2.31
	0.507.19	1.88	2.49	2.16
1874 38.42	10 A2 Q 12	2.34	2 . 19	1.221.19
1875	M.UU	2.85	2 . 67	981.43
1876 38.27		2.59	2 . 89	1.221.25
	1.537.69			1.101.30
1878 39.40	31.537.69 31.057.75	1.96		

The British American were included in the "other foreign" until the year 1868; and the Portuguese until 1874.

It will be noticed that the proportion of children born of American parents has changed but slightly in the whole period of twenty-three years. It was 42.09 per cent. in 1856, and 39.40 in 1878. In 1869, 1870, and 1871 the percentage was greater than in 1856. The extremes of the percentage of American parentage, in the 23 years, were 35.73 in 1863, and 45.19 in 1869. The percentage of Irish children kept pretty uniform, with a slight increase, from 1856 to 1865. Since the last date it has rapidly decreased, the percentage 28.08, in 1877, being the smallest in the twenty-three years.

Table IV. shows more definitely the parentage of the children born in 1878, as it gives the birthplace of both parents:—

TABLE IV. BIRTHS, 1878. Parentage.

	BIRTHPLACES OF MOTHERS.									
BIRTHPLACES OF FATHERS.	United States.	Ireland.	England, Scotland, and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number according to the father's birthoades		
United States. Ireland. England, Scotland and Wales Germany. British America. Portugal and Western Islands Other Countries.	28	95 732 32 6 13	51 32 131 4 4	3 1 48	29 13 11 2 58	30	2 1 3 7 1 35	1,215 890 237 82 103 35 56		
Whole number according to the mother's birthplace		882	222	53	118	30	49	2,627		

From this table it will be seen that of the 2,627 children born in Providence, in 1878, 1,035, or 39.40 per cent. of all were of purely American parentage; in addition to these there were 180 who had American fathers and foreign mothers; and 238 with foreign fathers and American mothers, making a total of 1,215, with American fathers, and 1,273 with American mothers; in both classes a smaller number than in the preceding year. There were 1,034 children, or 39.36 per cent. of all, whose parents were natives of the same foreign country; 140 children, or 5.33 per cent. whose parents were natives of different foreign countries; and 418, or 15.91 per cent. whose parents were one American and one foreign. In regard to these numbers and percentages it may be said, that of the 35 children whose parents were of "other countries" than those named in the table, the parents were not in every instance of the same foreign country; but as in many cases they were, and as it would be difficult to distinguish them, the countries not being specified in the table, they have always been included in that class, in giving these percentages.

NUMBER OF THE CHILD.

Table V. shows how many of the children born in 1878 were the first, second, third, etc., children of their mothers; also the same statistics for the 42,873 children born in the twenty-three years preceding, 1855-1877 inclusive:—

TABLE V. BIRTHS, 1878. Number of the Child.

NUME	E	R ()F (CHILI	о.	1878.	23 years, 1855-1877.	NUMBER	OF	CE	IILD.	1878.	23 years, 1855-1877.
First ch	ld	of	the	mothe	er	529	10,734	Twelfth chile	of	the	mother	16	167
Second	**	**	15	16		520	8,899	Thirteenth "	**	**	- 44	5	64
Third		**	44	- 64		456		Fourteenth "	**		**	2	28
Fourth		**	14	**		357	5,079	Fifteenth "	- 64	**	**		14
Fifth	11.		48	44		229	3,716	Sixteenth "	**	11	**		3
Sixth	**	**	**	44		198	2,677	Seventeenth"		11.	**	1	6
Seventh	44	**	**	**		122	1,894	Eighteenth "		**	**	1	
Eighth	**		46	**		91	1.185	Nineteenth "	48	44	44		4
Ninth	16	44	**	**		49	791					2000	
Tenth	**		**	**		38	517	Whole numb	er e	of cl	hildren.	2,627	42,873
Eleventh			**	**		13	275						

There were 529 children in 1878 who were the first children their mothers had borne; this was 20.14 per cent. of the whole number of children born during the year. Of the 42,873 children born in the 23 years. 1855-1877 inclusive, 10,734 or 25.04 per cent. were the first children of their mothers.

The next table shows the average number of children to each mother in each of 24 years, 1855-1878:—

Table VI. Births, 1855 to 1878.

Average number of children to each mother.

Years.	No. of Mothers.	No. of A	Aggregate of children. Average.
1855		1,600	• • • • • • • • • • • • • • • • • • • •
	•	•	5,562 3.35 children to each mother.
		1,688	
1858	1,706	1.724	5.6853.33 children to each mother
1859	1,570	1,593	5,6183.58 children to each mother.
	-	1.648	-
1861	1.694	1,725	6,0173.55 children to each mother.
1862	1,512	1,529	5.3223.52 children to each mother.
	•	1,307	•
	-	1.344	•
1865	1,279	1,208	4,625 3.60 children to each mother
866	1,614	1,632	5,7443.56 children to each mother
	•	1,624	· ·
		1,866	
1869	1,797	1,819	5,954 3.30 children to each mother.
1870	1,859	1,889	6,1453.31 children to each mother
1871	1,937	1,960	6,4203.31 children to each mother
1872	2,174	2,206	7,3463.38 children to each mother.
		2,128	
1874	2,738	2,780	8,9603.27 children to each mother.
1875	2,634	2,664	8,9773.41 children to each mother.
1876	2,604	2,631	8,9143.42 children to each mother
1877	2,516	2,543	8,7053.46 children to each mother.
1878	2,597	2,627	9,4173.63 children to each mother.
24 years	44,931	45,500	154,372 3.44 children to each mother.

The number of twin births in Providence, in 1878 was 30, which makes the number of mothers 2,597. These 2,597 mothers had borne, in all, 9,417 children, an average of 3.63 children to each mother.

This table shows how many children these mothers had already borne, and the average number for each; but it does not show the average number of children the same mothers may have in a life-time.

Table VII. shows the age of the mother in connection with the number of the child; also how many first, second, third, etc., children were born in 1878 to mothers at the respective ages given; and the number of mothers at each age, and the aggregate and average number of children they had borne.

Table VII. Births, 1878. Number of child, with age of mother.

				N	UMI	BER	OF	CI	IIL	D.					others a	bore in	they	number of a to mothers
AGE OF THE MOTHERS IN YEARS.		2d.	3d.	4th.	5th.	6th.	7th.	8th.	Mrn.	11th.	12th.	13th.	14th.	17th.	Number of mothers a each age.	Children they 1878.	Total children they had borne.	Average number of children to mothers at each age.
14. 16. 16. 17. 18. 18. 19. 19. 19. 19. 19. 19. 19. 19. 19. 19	11 8 20 30 44 44 45 *57 *43 30 24 19 11 11 11 11 11 11 11 11 11 11 11 11	8 14 15 52 38 50 53 50 44 *40 23 42 8 14 15 6 *13 9 2 2 3 3 3 3 3 2 2 2 2 3 8 1 1 1 1 1 1 1 2 2 3 3 3 3 3 3 3 3 3 3 3	20 24 34 47 *48 56 *27 43 *27 18 23 14 16 *10 6 5 4 5	9 10 22 *23 36 26 41 19 39 *21 11 26 10 *15 6 7	7 10 18 13 *36 9 24 *20 20 19 8 10 10 3 *8	3 6 9 6 19 7 *22 *21 18 21 *14 5 15 14	1 3 6 14 5 14 11 11 *12 6 7 3 8	3 *4 1 8 12 10 *9 6 8 6 5 10 	42445232.	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	200				128 70	16 17 10	207 238 293 378 422 450 616 414 824 350 677 6547 6547 408 308 423 307 409 1124	1.26 1.26 1.24 1.60 2.20 2.28 2.69 3.39 4.70 5.12 5.12 5.12 6.00 7.75 6.40 6.13 6.20 6.20 6.20 6.20 6.20 6.20 6.20 6.20

^{*}One of these mothers bore twins. † Two of these mothers bore twins.

The reference marks in this table indicate the age of the mother and the number of the child in cases of plurality birth; thus at the age of 22 years two mothers bore twins, and in both cases they were the second and third children.

The oldest woman bearing a child in Providence in 1878, was 50 years of age and had borne five children before. The number of women forty years old and over, who bore children in 1878, was 119; they had borne 842 children, or an average of 7.08 children to each mother. From July 1, 1863, to December 31, 1878, a period of fifteen years and six months, during which the age of the mothers has been recorded, there have been 1.554 women of forty years and over, who bore children; the number of children they had borne was 11,881, or an average of 7.64 children to each mother.

During this period of fifteen years and six months, there have been 31,576 children borne by mothers whose age was given. The number of children borne by mothers of each age during this time is as follows:—

Age of Mother. Children.	Age of Mother. Chile	iren. Age of Mother. Children
14 8	271,85	2 40 736
15 18	282,21	0 41 207
16 47	291,58	0 42 242
17 138	302,54	6 43 138
18 352	311,07	1 44 90
19 568	321,47	1 45 94
20 915	331,24	2 46 25
211,057	341,03	4 47 24
221,545	35 1,58	5 48 9
231,705	36 93	7 49 3
241,881	37 77	4 50 5
252,133	38 84	6
261,959	39 53	4 All known ages 31,576

The mothers of these children were divided as to age as follows:—

Under twenty years
Twenty, and under twenty-five
Twenty-five, and under thirty
Thirty, and under thirty-five
Thirty-five, and under forty
Forty, and under forty-five
Forty-five and over
Total

Of the 31,576 children born during this period of years, 17,963 or 56.89 per cent. of all were borne by mothers under thirty years of age; 12,040, or 38.13 per cent. by mothers between thirty and forty; and 1,573 or 4.98 per cent., by mothers of forty years and over.

PLURALITY BIRTHS.

There were 30 cases of plurality births in Providence, in 1878; this was three more than in 1876 or 1877; all were twin births.

The following statistics are recorded in regard to them: — Cases. Whole number 30; children born, 60.

Locality. First ward, 2; second, none; third, 5; fourth, 2; fifth, 2; sixth, 1; seventh, 3; eighth, 3; ninth, 5; tenth, 7.

Season. January, 2; February, 2; March, 3; April, 2; May, 1; June, 2; July, 4; August, 3; September, 3; October, 2; November, 3; December, 3.

Sex. In thirteen cases both children were males; in ten cases both were females; and in seven there was one child of each sex, making 33 male and 27 female children.

Color. In every case the twins were of white parents.

Number of the Child and Age of the Mother. In three cases they were the first and second children of mothers respectively, 23, 24 and 25 years old; in four cases they were the second and third children of mothers, two of whom were 22 years old; one 28 years old, and one 35; in four cases the third and fourth children of mothers 27, 29, 31 and 36 years old; in three cases the fourth and fifth children of mothers 26, 33 and 37 years old; in three cases the fifth and sixth children, of mothers 30, 33 and 40 years old; in five cases the sixth and seventh children of mothers of 32, 33, 36, 40 and 44 years; in two cases, the seventh and eighth children of mothers of 26 and 36 years; in three cases, the eighth and ninth children of mothers of 30, 35 and 43 years; in two cases, the ninth and tenth children of mothers of 39 and 42 years; and in one case the eleventh and twelfth children of a mother 39 years old. In twenty cases, or just two-thirds of the whole number, the mothers were thirty years old or over.

Parentage. In only six cases were both of the parents of American birth; in fourteen cases, (nearly half,) the parents were both

Irish; in two cases both parents were English, and in two, both German; in one, both were British American; in three cases, where the father was American, the mothers were respectively Scotch, Irish and British American; in two cases where the fathers were Irish, the mothers were respectively English and British American. This gives six American mothers; fifteen Irish; three English; two German; three British American, and one Scotch.

TWENTY-FOUR YEARS' SUMMARY OF PLURALITY BIRTHS.

During twenty-four years, 1855-1878 inclusive, the cases of plurality births occurring in Providence have been 560; of which 551 were of twins, and nine of triplets; whole number of children, 1,129. There have been no triplets born in Providence since 1873, when there were two cases.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children, and one female; making 9 male and 18 female children in the nine cases.

Of the 551 cases of twins, there were 177 in which both children were males; in 189, both were females; and in 185, there was one child of each sex; total, 539 male, and 563 female children. Of the 1,129 children in all the cases of plurality births 548 were males, and 581 females.

Color. In all the cases of triplets, the parents were white. Of the 551 cases of twins, in 533 the parents were white, and in 18 they were colored.

Parentage. Of the triplets, in one case, the parents were both American; in five cases, both were Irish; in one case, both German; in one, the father was English and the mother Irish; and in one, the father Scotch and the mother Irish. There was but one American mother in the nine cases.

The following statistics of the 521 cases of twin births, give the nativity of both parents, the first letter representing the father's birthplace and the second the mother's.

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British American; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish.

Parentage.	No.	Parentage	No.	Parentage.	No.	Parentage.	No.
A. and A I. and I E. and E S. and S G. and G B. A. and B. A. D. and D. P. and P. It. and It	218 25 4 15 10	E. and A S. and A B. A. and A G. and A Sw. and A A. and I E. and I	12 6 1 12	B. A. and I Sw. and I It. and I P. and I A. and E I. and E S. and E	1 1 1 3 6	I. and S E. and S I. and W	3 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

This table shows that of the 551 cases of twin-births, in 215 cases, or 39.02 per cent. of all, the mothers were born in the United States; in 249 cases, or 45.19 per cent., the mothers were born in Ireland, and in 87 cases, or 15.79 per cent., the mothers were natives of other foreign countries. Of the 560 cases of plurality births in the twenty-four years, the American mothers numbered 216, and the foreign mothers, 344.

Proportion of plurality births to whole number. The following statement is a comparison of the year 1878, with the twenty-three years previous, as to the whole number of births, and the proportion of plurality births:—

90 mann 1955 1	Single births.		-	Whole No. Births.	Whole No. Children.
23 years, 1855-1 1878					· <u>-</u>
24 years' total	44.371	551	<u> </u>	44.931	45,500

Proportion of cases of plurality births: -

	Twins.	Triplets.
1855-1877	1 in 81.3	1 in 4,703.7
1878	1 in 86.6	
1855–1878	1 in 81.5	1 in 4,992 8

Proportion of children of plurality births:-

1855–1877	Twins.	Triplets.
	1 in 43.7	•
1855-1878	<u>1 in 41.3</u>	1 in 1,685.2

Proportion of plurality births by parentage. The whole number of births in the twenty-four years, 1855–1878 inclusive, was 44,931. The following statement gives the number of births of American, foreign and mixed parentage; also the number and proportion of plurality births in each class:—

	Whole No. births.	Plurality cases. 1 in Per cent.
American parents		18199.50, or 1.00
Foreign parents	21,698	32367.18, or 1.49
American and Foreign	5,223	5693.27, or 1 07
_		
Total	. 44 ,931 . .	56080.23, or 1.25

Of the 560 cases of plurality births in this period of twenty-four years, the 181 cases of American parentage were 32.32 per cent.; the 323 cases of foreign parentage were, 57.68 per cent.; and the 56 cases of mixed parentage were 10.00 per cent.

MARRIAGES.

The number of marriages in Providence during the year 1878, was 1,016.

Table VIII. gives the number for each month and quarter of the year 1878, and for each of the six years, 1872-1877 inclusive; also the aggregate and average number for each month and quarter for the period of twenty-four years, 1855 to 1878 inclusive:—

TABLE VIII. MARRIAGES. Season.

MONTHS.	1878.	1677.	1876.	1875.	1874.	1873.	1872.	24 years. 1855–1878.	24 years' average.
January February March	73	68 65 38	79 93 44	75 79 71	93 87 56	105 111 49	97 83 53	1,699 1,526 1,010	71 63 42
First quarter	222	171	216	225	236	265	233	4,235	176
April May June		83 85 83	70 80 94	84 82 100	95 113 101	87 91 107	93 85 94	1,650 1,686 1,762	65 70 73
Second quarter	249	251	244	266	309	285	272	4,998	208
July	94 62 93	86 76 87	56 62 93	101 73 111	88 480 129	105 63 108	91 71 83	1,519 1,281 1,803	63 54 75
Third quarter	249	248	211	285	297	276	245	4,603	192
October November December	99 115 82	99 102 82	95 109 74	107 114 82	127 114 93	118 125 81	124 121 89	2,018 2,162 1,600	84 90 67
Fourth quarter	296	283	303	334	334	324	334	5,780	241
Yearly totals	1,016	953	949	1,079	1,176	1,150	1,084	19,616	817

The number of marriages in 1878 was 63 more than in 1877, and 67 more than in 1876. Estimating the population in the middle of the year 1878 at 101,000, the ratio of persons married to the population was one in 49.70 or 20.12 persons in each thousand of the population. There were 77 married to each one hundred children born; and 102 persons married to each 100 deaths.

Season. As is generally the case, the larger number of these marriages occurred in the latter half of the year; the largest quarterly total being 296 in the fourth quarter; the smallest quarterly total was 222, in the first quarter. The largest number in any one month was 115 in November; the smallest, 62 in August. The smallest monthly total is generally found in March; in the 24 years, 1855-1878, in which these reports have been published, there have been only four exceptions to this fact. In 1859, August had the same number as March, and July a smaller number; in 1862, July had the same number as March, and August a smaller number; in 1870, August had the smallest monthly total; while in 1878, both August and May have a smaller total than March, and February has the same number. According to the aggregate number of marriages for 24 years, as given in Table VIII., of each 100 marriages in Providence, about 25 occur in the winter months, December, January and February; 22 in the spring; 23 in the summer, and 30 in the autumn.

The following statement gives the number and percentage of marriages in each quarter of the year 1878, and in the aggregate for the corresponding seasons for the 23 years previous, 1855–1877 inclusive:—

	16	378.	1855 -	— 1877.
	Marriages.	Per cent.	Marriages.	Per cent.
January-March	. 222	21.85	4,013	21.58
April-June	. 249	24.51	4,749	25.53
July-September				
October-December	. 296	29.13	5,484	29.48
Total	.1,016	100.00	18,600	100.00

In table IX. are given the birthplaces of the brides and bridegrooms in each of the 1,016 marriages recorded in Providence in the year 1878:—

TABLE IX. MARRIAGES, 1878. Nativity of the Parties.

	BIRTHPLACES OF BRIDES.							
BIRTHPLACES OF GROOMS.	United States.	Ireland.	England, Scot- land and Wales.	Germany.	British America.	Portugal and Western Islands.	Other countries.	Whole number of C
United States. Ireland. England, Scotland and Wales. Germany British America. Portugal and Western Islands. Other Countries.	601 33 28 4 12 2 6	35 140 12 2 2 1 4	32 11 29 1 1 1	12 12 1	15 6 1 8	2	3 7	689 189 75 22 22 11
Whole number of brides	686	196	75	15	30	2	12	1,01

Bride and Groom born in the same Country. The number of marriages in which both parties were born in the same country was 799, or 78.64 per cent. of all; this is a little smaller proportion of this class than in 1877. In 601 marriages, or 59.15 per cent. of all, both bride and groom were born in the United States, this is 1.96 per cent. more than in 1877; in 140, both were born in Ireland; this is 13.78 per cent. of all, and 1.85 per cent. less than the same class in the preceding year; in 29 marriages, both parties were born in England, Scotland, or Wales; in 12, both were born in Germany; in 8, both in British America; in 2, both in Portugal or the Western Islands, and in 7, both were born in other countries not specified in the table.

Bride and Groom born in different Countries. There were 217 marriages or 21.36 per cent. of the whole number, in which the bride and groom were natives of different countries; thus 35 men born in the United States married women who were natives of Ireland; 32 men married women who were natives of England, Scotland or Wales; of the 686 grooms born in the United States, only 85 or 12.39 per cent. married natives of foreign countries; while of 184 who were born in Ireland, 44, or 23.91 per cent. married natives of other countries; of the 75 born in England, Scotland, or Wales, 46 or 61.33 per cent. married natives of other countries. In regard to the brides who were born in the United States, the numbers are the same as in the case of the grooms, the whole number being 686, and the number who married natives of other

countries, 85, or 12.39 per cent. Of the 196 brides born in Ireland 56, or 28.57 per cent. married men of other countries. The number of brides born in England, Scotland or Wales was also the same as the number of grooms, 75, and the proportion marrying natives of other countries, 61.33 per cent. From these figures it seems that the natives of the United States are more apt to marry their own country people than the natives of other countries are.

The 19 grooms, natives of "other countries," were as follows: 5 born in Sweden; 3 in St. Helena; 2 each in France and Italy; and 1 each in Denmark, Spain, Norway, Greece, Switzerland, Bermuda, and the West Indies. The 12 brides born in "other countries" were 6 in Sweden; 2 in France; and 1 each in Norway, Switzerland, St. Helena, and the Sandwich Islands.

The number and percentage of American, foreign, and mixed marriages in Providence, are shown in the following table for each of the 24 years, 1855-1878 inclusive. By mixed marriages are understood those where one of the parties was of American, and one of foreign birth:—

		rican.	Fore	ign.	Mixed.
Year.	Number.	Per cent.	Number.	Per cent.	Number. Percent.
1856	. 308	48.73	246	41.93	59 9.34
1856	. 352	54.74	212	32.97	7912.29
1857	384	62.54	175	28.50	55 8.96
1868	283	50.54	214	38.21	6311.25
1859	333	54.50	224	36.66	54 8.84
1560	336	54.19	225	36.29	59 9.52
1961	304	55.58	182	33.27	6111.15
1862	288	52.65	179	32.72	8014.63
1863	342	56.07	192	31.47	7612.46
1864	427	57.78	211	28.55	10113.66
1865	417	58.82	202	28.49	9012.69
1886	471	58.00	244	30.05	9711.95
1867	485	56.73	269	31.46	10111.81
1868	530	60.78	215	24.66	12714.56
1869	530	56.99	286	30.75	11412.26
1970	552	57.56	262	27.32	145
l871	523	55.46	278	29.48	14215.06
1672	591	54.52	305	28.14	188 17.34
1873	652	56.69	317	27.57	181 15.74
BC4	648	66.10	346	29.42	18215.48
1875	586	54.31	320	29.66	17316.03
1876	547	57.64	241	25.39	16116.97
877	545	57.19	233	24.45	17518.36
1878	661	59.15	245	24.12	17016.73
A veeze	11.035	56.27	5,842	97.79	2.733

The proportion of American marriages has steadily increased for some years, and that of foreign marriages proportionately decreased; the proportion of mixed marriages in 1878 was less than in the two years preceding, but still considerably above that of the entire period. The apparent discrepancy between the whole number of marriages as given here and in Table VIII., for the same period is explained in the report for 1875.

Marriages of colored persons. There were 54 marriages of colored persons in Providence, in 1878; this is more than in any year since 1874. The aggregate and average age of the parties in these marriages were as follows:—

Address of the Control of the Contro	
El malas A	1 500 mans 4 manus and 97 20
or mates	1,506 years Average age, 27.89.
to demote the second	1 010
of lemales	1,342 years

The whole number of marriages of colored persons in the twenty-three years, 1856-1878, was 828; the aggregate and average age, as follows:—

828 males Aggregate age, 24,225 years	Average age, 29.26.
828 females Aggregate age, 21,712 years	

Age of persons married. The next table gives the number and percentage of brides and grooms in each division of ages: —

TABLE X. MARRIAGES, 1878. Age of the Parties.

	AGES OF THE BRIDES.							number soms.	Jo :	
AGES OF THE GROOMS.	Under 20 years.	20 to 25.	25 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.	Whole num of Grooms	Percentuge Grooms.
Under 20 years	15 86 44 16	5 198 147 48 10 2	1 41 112 85 14 1	1 4 21 75 27 5 1	1 1 8 15 14 5	2 4 1 1	1 1 2	1	22 330 325 232 60 27 9 2	2.17 32.48 31.98 22.86 6.75 2.66 0.88 0.20
Whole number of Brides	161	410	254	134	44	8	4	1	1,016	
Percentage of Brides	15.85	40.35	25.00	13.19	4.33	0.79	0.39	0.10		100.00

Of the 1,016 grooms, 352, or 34.65 per cent., were less than 25 years old; and of the brides, 571, or 56.20 per cent.

The aggregate and average age of all the persons married in Providence in 1878 were as follows:—

1 Ald males	1		1
1,VD (1986)	ARRITEMIC MET,	2,500 1623	Average age, 25.39
1.016 females	Approprie non.	25.786 Tears	Average age, 25.40
.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			to the state of th

The following table shows the aggregate and average age of the persons married in Providence, in each of the twenty-three years from 1856 to 1878 inclusive:—

	Males.	Females.
Years.	No. Aggregate age. Average age.	No. Aggregate age. Average age.
866	639 18,104 years28.33 years.	
857	607 16,753 years27.59 years.	
856	500 15,850 years28.29 years.	. 560 13,702 years24.47 years,
869	611 16,940 years 27.81 years.	. 611 14,616 years 23.92 years.
860	620 17,125 years27.60 years.	. 620 14,667 years23.66 years.
261	547 15,448 years 28.24 years.	. 547 13,316 years 22.33 years.
962	547 15,773 years28.83 years.	. 547 13,565 years21.80 years.
	610 17,208 years 28.21 years.	. 610 14,792 years
1864	739 21,361 years28.90 years.	
1865	709 20,539 years28.96 years.	. 709 17,605 years24.83 years.
1866	812 22,801 years28.06 years	
1867	855 21,443 years28.56 years.	· · · · · · · · · · · · · · · · · · ·
1808	· •	
	930 26,549 years28.55 years	
	950 27,343 years28.51 years	
	943 26,743 years28-37 years	
		. 1,084 26,875 years21.76 years.
		. 1,150 28,701 years24.96 years.
		. 1,176 29,510 years25.09 years.
		. 1,079 27,000 years25.11 years.
		. 949 23,640 years24.91 years.
		. 953 24,002 years25.28 years.
1678	1,016 29,460 years28.99 years	1,016 25,796 years25.40 years.
	8,967541,208 years28.53 years	

In 1856, the ages of 17 males and 19 females, and in 1857, the ages of 7 males and 8 females, were not given. This will account for the apparent discrepancy in the first lines of this table.

Number of the marriage. Table XI. shows the number of the marriage of each party, in each of the 1,016 marriages in 1878:—

Table XI. Marriages, 1878. Number of times married.

GROOMS. Number of the Marriage.	BRIDES.	Whole		
	First.	Second.	Third.	Grooms.
First	734 125 2	67 67 9	6 2 2	807 194 13
Fifth	1	1		1
Whole number of Brides	862	144	10	1,016

Of these 1,016 marriages, there were 734 cases, or 72.24 per cent., in which it was the first marriage of both parties; 67, or 6.59 per cent., in which it was the second marriage of both. In both classes the percentage is smaller than in 1877; but of marriages, where it was the second of the groom, and the first of the bride, the number and proportion in 1878 were much larger than in the preceding year; in 1877 they were 88, or 9.23 per cent. of all; in 1878, 125, or 12.30 per cent. There was one instance in 1878 where it was the fifth marriage of the groom and the first of the bride; and one was the sixth of the groom and second of the bride.

Denominational. The number of marriages solemnized in 1878 by clergymen of the various religious denominations, or by civil authority, was as follows:

Roman Catholie	Lutheran 12
Calvinistic Baptist	Advent Christian 2
Free Will Baptist 41	Evangelical Advent 10
Methodist154	New Jerusalem 8
Episcopalian129	Latter Day Saints 1
Congregational 92	Jewish 1
Unitarian 34	Friends' Marriage 1
Universalist	Justice of Supreme Court 1
Presbyterian 9	Denomination not specified 5
Christian 16	_
Bethel Pastor 11	Total

Nativity and Residence. Of the 1,016 marriages in Providence, in 1878, there were 75, or 7.38 per cent in which both bride and groom were natives of the city. In addition to these, there were 157 marriages in which the bride only was a native of Providence; and 121 in which the groom only was born here. This gives a total of 428 persons married who were born in Providence; this is 21.06 per cent. of the whole number of persons married. There

were 100 marriages in which both parties were non-resident; this is 9.84 per cent of all.

Marriage and Education. The number of persons married in Providence in 1878 was 2,032; they were divided as to sex and nativity as follows:—

Born in United States	Males, 686	.Females, 686	Total, 1,372
Born in foreign countries	Males, 330	.Females, 330	.Total, 660
ŭ		·	
Total		1.016	2.032
		,	

The number and proportion of each class who signed the marriage certificate with a mark were as follows: —

1878. Males, born in United States		Signed with a mark.	100.
Females, born in United States			
Total, born in United States		55	4 01
Males, born in foreign countries	330	59	17.88
Females, born in foreign countries	330	107	32.42
Total, born in foreign countries	660	166	25.15

Of the 55 born in the United States, who signed with a mark, 14, (8 males and 6 females), were colored persons; and 32 others, (8 males and 24 females), were of foreign parentage; this leaves 9 persons, (6 males and 3 females), of white American parentage, who signed the marriage certificate with a mark. This is 5 more of this class than in 1877.

These statistics for the nineteen years, 1859-1877 inclusive, are as follows:—

1859-1877. Males, born in United States	Whole number married.	Signed with a mark.	In each 100.
Females, born in United States	•		
Total, born in United States	20,521	1,220	5.94
Males, born in foreign countries Females, born in foreign countries			
Total, born in foreign countries			

The number and proportion of those who signed with a mark in 1878, were less in all classes than in 1877; of those born in the United States, 9 less, 5 males and 4 females; of those born in foreign countries, 23 less; males, 25 less, and females, 2 more. In 1877, the whole number of persons married was 1,906; of those who signed with a mark, 253, or 13.27 in each hundred; in 1878, the number of persons married was 2,032; of those who signed with a mark, 221, or 10.88 in each hundred.

DEATHS.

The number of deaths in Providence in 1878 was 1,989; this was 51 more than in 1877, and is a larger number than has ever been recorded in these reports, though the number in 1874 was nearly as large, 1,987. The number of males in 1878 was 47 more than in 1877; the number of females 4 more; of whites, 50 more; of colored, 1 more; of natives of the United States, 54 more; of natives of foreign countries, 3 less; of American parentage, 19 less; of foreign parentage, 70 more.

The average death-rate in Providence for the twenty-four years, 1855-1878 inclusive, was one in 50.80 of the population. In 1878 it was as follows in the different classes of the population:—

	Population, 1875.	Deaths, 1878.	Deaths to To population, each 1,000 one death in population.
Whole number	100,675	1,989	50.6219.76
Males	48,701	969	50.2619.90
Females	51,974	1,020	50.9519.63
Whites	97,188	1,883	51.6119.37
Colored	3,487	106	32.9030.40
Born in the United States	71,438	1,565	21.91
Born in Ireland	18,458	313	58.9716.96
Born in England, Scotland and Wa	ales 6,110	63	96.9810.31
Born in Germany	1,246	14	89.00
Born in other foreign countries	3,423	34	100.68 9.93
Total of foreign born	29,237	424	68.9614.46
American parentage	47,316	889	53.2218.79
Irish parentage	37,440	795	47.0921.23
English, Scotch and Welsh parents	nge 9,636	172	
German parentage	2,516	40	62.9015.90
Other foreign parentage	3,767	y3	40.5124.69
Total of foreign parentage	53,359	1,100	48.5820.62

Table XII., on the next page, gives some general statistics in regard to the deaths in 1878:—

TABLE XII. DEATHS, 1878.

Sex, condition, color, nativity, parentage, locality, season.

		_		_						_			
1878.	January.	February.	March	April.	May.	June.	July.	August.	September.	October.	November.	December.	Whole year.
Whole number of Deaths	208	175	158	164	131	128	203	175	142	160	165	180	1,989
Sez.— Males Females	98 110	84 91	74 84	88 76	66 66	52 76	111 92	83 92	70 72	78 82	84 81	82 98	969 1,020
Condition.— Married	48 136 18 6	37 117 13 6	44 98 11 4	32 100 19 12 1	39 71 14 7	37 69 14 8	9	35 118 16 6	49 77 11 4	36 107 14 3	43 107 11 4	47 103 19 11	489 1,250 169 74 7
Color.— Whites Colored	197 11	168 7	151 7	149 15	125 6	121 7	189 14	170 5	136 6	150 10	160 5	167 13	1,883 106
Nativity.— United States	171 27 6 1 2	133 29 9 2 2	131 22 2	131 26 4	93 32 3	95- 24- 4- 1- 2-	24 5 1 3	145 20 6 1 2	112 21 5 1 3	127 22 5 5	120 36 6 1 1	138 30 8 1 3	1,565 313 63 14 28 2
Parentage.— American Irish English and Scotoh German British American Portuguese Other foreign	101 71 21 5 5	61 86 16 3 5 4	77 58 13 5 4	80 62 12 3 4 2	58 58 8	51 58 11 2 3	4 3 4	70 74 17 4 4 2	75 46 10 2 5	60 73 16 4 8 1	65 76 15 3 4	99 56 14 5 4	889 795 172 40 49 21 23
Locality, Wards	31 4 17 14 10 9 21 19 34 41 8	29 5 23 9 15 12 14 9 22 31 6	16 3 22 11 21 7 11 12 20 29 6	26 9 16 6 14 10 9 13 17 33	17 2 9 10 15 7 9 23 15 20 4	10 7	25 11 23 11 27	25 3 17 11 16 7 14 19 30 24	21 15 4 15 22 27 22 27 24 4	10 8 15 10 13 14 22 16 23 25	26 5 11 9 17 7 15	28 4 5 11 13 10 18 22 36 28 5	264 62 182 109 189 108
Totals	208	175	158	164	131	128	203	175	142	160	165	180	1,989

Season. Contrary to the usual rule, the largest number of deaths was in the first quarter of the year instead of the third; the smallest number was in the second quarter. The following statement gives the number of deaths in each quarter of each of the last six years; also the whole number and the average for each quarter for the period of twenty-three years, 1855–1877 inclusive:—

	1878.	1877.	1876.	1875.	1874.	23 years 1855-1877. 1873. Aggregate. Av'ge.
January-March	541	411	383	488	444	427 6,978303
April-June	423	440	419	401	501	407 6,486
July-September	520	557	573	531	556	504 8,833384
October-December	505	530	490	495	486	381 7,178312
Whole year	1,989	1,938	1,865	1,915	1,987	1,71929,4751,281

The number and percentage for each quarter of the year 1878 are given below; also the same statistics for the period of twenty-three years, 1855-1877 inclusive:—

	1878		23 years,	1855-1877	
	Deaths.	Per cent.	Deaths.	Per cent.	
January-March	. 541	27.20	6,978	23.67	
April-June	423	21.27	6,486	22.01	
July-September	520	26.14	8,833	29.97	
October-December	. 505	25.39	7,178	24.38	
Totals	1.000	100.00	20.475	100.00	

The number and percentage of deaths in each quarter of each year during the period of twenty-four years, 1855-1878, were as follows:

	January-M	farch.	April-J	ine.	July-8		Oct.	
	Deaths. Pe	r cent.	Deaths.	Per cent.	Deaths.	Per cent.	Deaths.	Per ct
1855	213	21.49	208	.20.99	339	.34.21	231	23.31
1856	308	28.92	244	.22.91	296	.27.80	217	20.37
1857	214	23.14	173	.18.70	286	.30.92	252	27.24
1858	279	7.43	278	.27.34	257	.25.27	203	19.90
1859	213	23.69	205	.22.80	270	.30.04	211	23.47
1860	243	24.27	209	.20.88	297	.29.67	.,252	25.18
1861	240	22.84	236	.22.45	336	.31.97	239	22.74
1862	197	21.55	192	.21.01	271	.29.65	254	27.79
1863	287	23.64	248	.20.43	370	.31.14	301	24.7
1864	302	23.58	273	.21.31	403	.31.46	303	23.6
1865		24.69		.19.82				
1866							COMPAND THE	23.3
1867						.20.79	214	
1868			244					
1869				.19.90		.28.91		
1870						*****		
1871				.20.98				
				.22.71				
				.23.68				
				.25.21				
1875				.20.94			495	of the second
			419					
1877		21.21				.28.74		27.34
1878	5415	7.20	423	.21.27	520	.26.14	505	25.31

It is an established rule, with very few exceptions, that the largest number and percentage of deaths occur in the third quarter

of the year; the few exceptions have been when epidemic diseases have prevailed extensively at other seasons; as in 1856, when scarlatina, croup, and pneumonia prevailed in the first quarter; in 1858, the first and second quarters show a larger percentage than the third, owing to the prevalence of scarlatina and measles. The year 1878 presents another marked exception to this rule, the number and percentage of the first quarter being the largest of the year; this is owing to the fact that a large proportion of the deaths from measles, bronchitis, pneumonia and congestion of the lungs occurred at that season. The number is smaller in the third quarter than it has been since 1873, and the percentage less than in any year since 1869. One reason for this is that the number of deaths from cholera infantum and diarrhea, which usually prevail largely in the third quarter, was less in 1878 than in any year since 1871. The smallest percentage of the year is generally found in the second quarter, though there are many more exceptions to this than to the rule just spoken of. The prevalence of some epidemic has frequently raised the percentage of this quarter above that of the first or fourth, but it has never been made the largest in the year. The years 1856 and 1858 have already been mentioned and in 1874 also, there was a large percentage in the second quarter owing to many deaths from scarlatina, consumption, pneumonia and congestion of the lungs. In 1878, the percentage for this quarter was the smallest of the year.

Sex and Parentage. The decedents in 1878 were divided as to sex and parentage as follows:—

			•
Acteign benencede			
Total	Q40 meles	020 females	Total 1.080

The number of females was 51 more than of males; of American parentage, 7 more; of foreign parentage, 44 more. The number of foreign parentage is 211 more than of American parentage; of males, 87 more; of females, 124 more, the excess of females among this class in 1878 being much greater than among the decedents of American parentage.

The decedents of both classes during the twenty-four years, 1855-1878 inclusive, were as follows, in regard to sex:—

Condition. Comparing the decedents in the different classes of civil condition, with those of 1877, we note the following differences:

Married, 76 less in 1878 than in 1877; single, 115 more; widows, 5 more; widowers, 6 more; divorced, 1 more.

Color. The number of deaths among the whites in 1878 was 50 more than in 1877; among the colored, 1 more. The number of births among the colored people of Providence in 1878 was 104; the proportion of births to deaths, 100 to 101.9.

In the twenty-four years, 1855-1878 inclusive, there have been among our colored population 1,743 births, and 1,841 deaths; a proportion of 100 to 105.6.

Nativity and Parentage. Table XII. shows how great is the difference between the number of decedents registered according to nativity and those registered according to parentage. It is an important distinction, as many who are born in this country, are yet, as children of foreign parents, under the same social and hygienic conditions and influences as the class more strictly called foreign because born in foreign countries. The number of decedents of American birth in 1878, was 1,565, or 78.68 per cent. of all; the number of American parentage, 889, or 44.70 per cent. The number of American birth was 54 more than in 1877; the number of American parentage, 19 less.

The number of decedents of foreign birth in 1878, was 424, or 21.32 per cent. of all; of foreign parentage, 1,100, or 55.30 per cent.; compared with the numbers for 1877, the number of decedents of foreign birth in 1878, was 3 less; of foreign parentage, 70 more.

Of the 4 born in "other countries" than those named in the table, one was born in Italy; one in Turkey, and two in St. Helena.

Of the 23 of "other foreign" parentage, 5 were of Italian parentage; 4, of St. Helena; 3, of Polish; 2 each, of Swedish, Dutch and French; and one each of Danish, Turkish, Swiss, West Indian, and South American parentage.

Locality. Table XII. gives the number of deaths in each ward of the city and in Public Institutions, in each month and for the whole year. In the first ward the number was 23 less than in 1877; in the second, 5 less; in the third, 13 less; in the fourth, 3 less; in the fifth, 20 more; in the sixth, 13 less; in the seventh, 8 more; in the eighth, 15 more; in the ninth, 56 more; in the tenth, 2 more; in Public Institutions, 7 more.

The number of decedents in each ward, according to parentage, was as follows:—

Wards.	American.	Foreign.	Wards.	American.	Foreign
I	106	158	VII	75	9:
II	47	15	V111	121	6
ш	76	105	IX	146	15
IV	63	16	x	76	25
v	64	125	Public Inst	itutions, 44	4
VI	71	37		<u> </u>	
Who	le city				1.10

Average Age. Table XIII. shows the number of deaths, and the aggregate and average age according to parentage, in each month of the year 1878:—

TABLE XIII. DEATHS, 1878.

Aggregate and Average Age, and Parentage.

	American Parentage.			Fore	ign Pare	ntage.	Total American and Foreign.		
1878.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.
January February March March April. May June July August September October November December	101 61 77 80 58 51 92 70 75 60 65	3,183 1,483 2,250 2,580 1,843 2,061 1,967 2,183 2,274 1,703 2,170 3,027	31.51 24.31 29.22 32.25 31.77 40.41 21.38 31.18 30.32 26.38 33.38 30.58	107 114 81 84 73 77 111 105 67 100 100 81	2,364 2,489 1,604 2,219 2,123 1,942 1,879 1,818 1,805 2,060 2,398 2,477	22.09 21.83 19.80 26.42 29.06 26.22 16.93 17.31 26.94 29.60 23.98 30.58	208 175 158 164 131 128 203 175 142 160 165 180	5,547 3,972 3,854 4,799 3,966 4,003 3,846 4,001 4,079 3,763 4,568 5,504	26.67 - 22.70 24.39 29.26 30.27 31.27 18.95 22.86 22.52 27.68 30.58
Whole year	889	26,724	30.06	1,100	25,178	22.89	1,989	51,902	26.09

The average age of all decedents in 1878, was 26.09 years, which was 1.65 years less than in 1877; the average age of American decedents in 1878, 30.06 years, or 1.16 years less than in 1877; of decedents of foreign parentage, 22.89 years, or 1.78 years less than in 1877.

By comparing the different months in the two years, we find many variations. Among the decedents of American parentage, the average age in each of the first five months of 1878 is much less than in the corresponding months of 1877; in June the average age is much higher in 1878; in July it is slightly less, and in

all the remaining months of the year, more than in the previous year. Among the decedents of foreign parentage the average age was less in 1878 than in 1877, in the months of January, February, March, June, July, August, October and November; and greater in the months of April, May, September, and December.

Table XIV. is a recapitulation, and gives the same statistics for the period of twenty-three years, 1856-1878:—

TABLE XIV. DEATHS, TWENTY-THREE YEARS, 1856-1878.

Recapitulation. Aggregate and Average Age by Parentage.

37	7			1	S. I mic	ntage.	Total American and Foreign.		
YEARS,	Number of	Aggregate	Average	Number of	Aggregate	Average	Number of	Aggregate	Average
1856-1878.	Deaths.	Age.	Age.	Deaths.	Age.	Age.	Deaths.	Age.	Age.
January February March April May June June July August September Occember	1,340	45,836	34.20	1,166	28,984	24.86	2,506	74,820	29.86
	1,225	41,459	33.84	1,137	25,103	22.08	2,362	66,562	22.18
	1,276	44,457	34.84	1,162	28,739	24.73	2,438	73,196	30.02
	1,254	42,497	33.88	1,186	30,469	25.69	2,440	72,966	29.91
	1,165	42,287	36.30	1,081	27,871	25.78	2,246	70,158	31.23
	985	34,049	34.57	1,030	24,673	23.96	2,015	58,722	29.14
	1,357	34,706	25.58	1,624	29,602	18.23	2,981	64,308	21.57
	1,499	41,015	27.36	1,733	31,265	18.04	3,232	72,280	22.36
	1,341	39,327	29.33	1,460	28,023	19.19	2,801	67,350	24.04
	1,185	39,008	32.92	1,303	29,734	22.81	2,488	68,742	27.63
	1,215	40,591	33.41	1,218	30,752	25.25	2,433	71,343	29.32
	1,314	47,465	36.12	1,217	31,127	25.58	2,531	78,592	31.05

A careful study of these two tables will show marked differences between the average age of the two classes of decedents in the different months. In the year 1878, (Table XIII.,) the highest average age of decedents of American parentage was 40.41 years, in June; the lowest, 21.38 years, in July. Of the decedents of foreign parentage, the highest average age was 30.58 years, in December; the lowest, 16.93 years, in July. It is an almost universal rule in Providence that the average age of the decedents of foreign parentage is less than that of American decedents; the month of December in 1878 is one of the rare exceptions; the average age of both classes in that month being 30.58 years; the other months of the year follow the general rule; the greatest difference being 15.19 years, in June. The difference between them in the whole year is 7.17 years.

Table XIV. shows differences of the same nature in the period of twenty-three years, 1856-1878 inclusive. The average age of the American decedents exceeds in every month, and in the whole year, that of the decedents of foreign parentage. In this table we find the highest average age of American decedents to be 36.30 years, in May; the lowest, 25.58 years, in July. Of foreign decedents, the highest average age is 25.78 years, in May; and the lowest, 18.04 years, in August. The highest average age of foreign decedents is but little above the lowest average among the Americans. The greatest difference between the two classes in any one month, for the long period of years, was 11.76 years, in February; the least difference, 7.35 years, in July. The difference between the two classes for the entire year, in the long period was 9.88 years.

These differences may be more plainly seen by putting the figures in the form of a table; it will be seen that in December, 1878, the average age was the same for both classes; in all other cases the average age of the decedents of foreign parentage, was less than that of the American decedents:—

Difference in years, between the average age of American and foreign decedents.

		1878.		23 years, 1856–1878.				
Months.	Averag	e Age.	1	Averag	e Age.			
	American.	Foreign.	Difference.	American.	Foreign.	Difference		
January	31.51	22.69	9.42	34.20	24.86	9.34		
February	24.31	21.83	2.48	33.84	22.08	11.76		
March	29.22	19.80	9.42	34.84	24.73	10.11		
April	32.25	26.42	5.83	33.88	25.69	8.19		
May	31.77	29.08	2.69	36.30	25.78	10.52		
June	40.41	25.22	15.19	34.57	23.95	10.62		
Jaly	21.38	16.93	4.45	25.58	18.23	7.35		
Atrust	31.18	17.31	13.87	27.36	18.04	9.32		
September	30.32	26.94	3.38	29.33	19.19	10.14		
October	28.38	20.60	7.78	32.92	22.81	10.11		
Sovember	33.38	23.98	9.40	33.41	25.25	8.16		
December	30.58	30.58	0.00	36.12	25.58	10.54		
Whole year	30.06	22.89	7.17	32.50	22.62	9.88		

Generally the difference is greater in the long period of time than in the single year; but in January, June, August and November, it is the reverse. The greatest contrast is in December, where for the year 1878 there is no difference; while in the long period of years the difference between the average age of the two classes is 10.54 years.

The largest monthly aggregate of deaths, (Table XIV.), was as follows: American parentage, 1,499, in August; foreign parentage, 1,733, also in August; the smallest monthly aggregate was in June, for both classes, American, 985; foreign, 1,030.

The percentages of the total deaths in the different quarters, for the period of twenty-three years, 1856-1878 inclusive, were as follows:—

1st quarter.	2d quarter.	3d quarter.	4th quarter
American parentage25.31	22.46	27.69	24.51
Foreign parentage22.62	21.53	31.45	24.40

Table XV. gives the number of decedents, and the aggregate and average age, according to parentage, in each of the twenty-three years, 1856-1878 inclusive, and for the entire period:—

Table XV. Deaths, 1856-1878.

Recapitulation. Aggregate and average age by parentage.

	Americ	an Pare	ntage	Foreig	n Paren	tage.	Total	America Foreign.	n and
YEARS.	Deaths.	Aggregate Years.	Average Years.	Deaths,	Aggregate Years.	Average Years.	Deaths.	Aggregate Years.	Average Years.
856	541	14.554	26.90	524	8,258	15.75	1,065	22,819	21.43
857	523	16,314	31.19	402	8,007	19.91	925	24,321	26.29
858		15,910		505	8,478	16.78	1,017	24,388	23.98
859	479	14,993	31.30	420	8,640	20.57	899	23,633	26.22
860		16,660		489	9,751	19.94	1,001	26,411	26.3
861		16,327	30.34	513	9,605	18.72	1,051	25,932	24.67
862		14,589	31.71	454	10,904	24.21	914	25,583	28.0
863		19,398	32.49	617	13,757	22.29	1,214	33,155	27.3
905		21,732 19,628	33.48	632 580	11,365 11,991	17.98 20.67	1,281	33,097 31,619	26.1
866		18,906	36.85	523	12,762	24.40	1,036	31,668	30.5
867		19,258	36.47	432	12,140	28.12	960	31,407	32.7
868	596	21,779		514	12,539	24.30	1,110	34,318	30.9
869		20,490		629	14,776	23.49	1,256	35,266	28.0
370		21,078	32.83	621	14,393	23.18	1,263	35,471	28.0
371	643	22,725		611	17,005	27.83	1,254	39,730	31.6
372		27,572	33.14	771	19,160	24.88	1,603	46,732	29.1
373		28,675		857	10,061	22.24	1,719	47,736	27.7
874		27,588	31.17	1,102	24,802	22.51	1,987	52,390	26.3
875		30,419		1,001	24,302	24.28	1,915	54,721	28.5
876		29,025		990	23,959	24.20	1,865	52,984	28.4
877		28,353 26,724	31.22	1,030	25,410 25,178	24.67 22.89	1,938	53,763 51,902	27.7
36	009	40,643	30,00	1,100	20,170	24.00	1,000	01,002	2010
3 years' total	15,156	492,697	39.50	15,317	346,342	90 60	30,473	830,039	27.5

The difference between the two classes ranges from 6.55 years in 1877, to 15.50 years in 1864. The difference in the entire period of time is 9.88 years.

Number and percentage of decedents at different ages. Table XVI. gives the number and percentage, according to parentage, of decedents at different periods of life, in 1878, and in the twenty-three years, 1856-1878 inclusive:—

Table XVI. Deaths, 1878.

Percentage at different ages, according to parentage.

		rican ntage. 78.		eign ntage. 78.	Total can a eign.	Ameri- nd For- 1878.	23 1	ears, 0 1878.
AGKS.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Under 1 year		18.90 7.98 9.90	205 103 161		373 174 249	18.76 8.74 12.52	2,745	9.01
Total under 5	327	36.78	409	42.64	796	40.02	11,562	37.94
5 to 10	58 27	6.52 3.04	25	2.27	138 52	6.94 2.62	1,501 629	4.93 2.06
20 to 30	56 49	2.92 7. 11 5.51	37 103 113	10.27	162	3.16 8.49 8.14	1,039 3,079 2,786	3.41 10.11 9.14
40 to 50		6.75 8.10 8.44	69 73 51	6.27 6.64 4.64	129 145 126	6.49 7.29 6.34	2,347 2,181 2,118	7.70 7.16 6.95
70 to 80	79 40	8.88 4.50	62 13	5.64	141 53 15	7.09 2.67	1,994 1,022	6.5 <u>4</u> 3.35
Total		100.00		100.00			30,473	. ——

By Table XIII. we saw that the average age of decedents of both classes was less in 1878 than in 1877; it corresponds with that fact to find by Table XVI. that the percentage of decedents under five years of age was larger in both classes in 1878, than in the preceding year. Of American decedents, the percentage of this class, in 1878, was 36.78, or 3.85 per cent. more than in 1877; of foreign decedents, the percentage was 42.64, or 1.38 per cent. more than in 1877; of the whole number of decedents, 40.02, or 266 per cent. more than in 1877.

The following recapitulation gives the percentage, according to parentage, of the decedents under five years old, in each of the twenty-three years, 1856-1878 inclusive:—

	American. Under five years.	Foreign. Under five ye a rs.	Total American and Foreign. Under five years.
856	37.15 per cent	54.95 per cent	45.91 per cent
857	32.31 per cent		t46.75 per cent.
858	35.35 per cent		
859	28.60 per cent		37.82 per cent
1860	32.81 per cent	48.87 per cen	t40.66 per cent
1861	35.87 per cent	52.63 per cen	t41.05 per cent
1862	31.08 per cent	42.07 per cent	t36.54 per cent
1863	28.47 per cent	46.84 per cen	t37.81 per cent
1864	28.81 per cent	51.11 per cent	39.82 per cent
1865			
			t
1867		36.35 per cen	t29.59 per cent
	•	•	t34.69 per cent
			t36.72 per cent
			32.22 per cent
			t35.27 per cent
			36.71 per cent
		•	t39.31 per cent
	•	-	t36.66 per cent
			t38.96 per cent
			t
			t40.02 per cent

During the thirty-nine years, 1840-1878 inclusive, the number of decedents in Providence was 43,618, of whom 17,297 were less than five years old; the number and percentage at each age under five years were as follows:—

Whole number of decedents	43,618
Under one year	8,517, or 19.53 per cent. of all.
One, and under two years	4,528, or 10.38 per cent. of all.
Two, and under five years	4,252, or 9.75 per cent. of all.
Total, under five years	17,297, or 39.66 per cent. of all.

Table XVII. shows the number and percentage according to parentage, of decedents at different periods of life, in the twenty-three years, 1856-1878 inclusive:—

Table XVII. Deaths, 1856-1878.

Number and Percentage at different Ages according to Parentage.

	America age, 23 year 183	78.	Foreign I 23 years,	1990-1979		
AGE.	Number of Deaths.	Percentage in each divi- sion of age.	Number of Deaths.	Percentage in each divi- sion of age.	Number of Deaths.	Percentage in each divi- sion of age.
Under one year	2,638 992 1,135	17.41 6.54 7.49	3,374 1,753 1,670	22.03 11.45 10.90	6,012 2,745 2,805	19.73 9.01 9.20
Total under 5	4,765	31.44	6,797	44.38	11,562	37.94
5 and under 10 10 and under 15 15 and under 20	753 341 560	4.96 2.24 3.69	748 288 479	4.88 1.88 3.13	1,501 629 1,039	4.93 2.06 3.41
20 and under 30	1,421 1,318 1,182	9.37 8.69 7.79	1,658 1,468 1,165	10.83 9.58 7.61	3,079 2,786 2,347	10.11 9.14 7.70
50 and under 60 60 and under 70 70 and under 80 80 and under 90	1,251 1,289 1,358 774	8.25 8.50 8.96 5.16	930 829 636 248	6.07 5.41 4.15 1.62	2,181 2,118 1,994 1,022	7.16 6.95 6.54 3.35
90 and over	144	0.95	71	0.46	215	0.71
Totals	15,156	100.00	15,817	100.00	30,473	100.00

By Tables XIII. and XIV. it was shown that the average age of decedents of foreign parentage is almost always considerably less than that of American decedents. By Tables XVI. and XVII. we see how much larger is the percentage of deaths of young children in the former class than in the latter, both in the single year, and in the long series of years. The decedents of American parentage under five years of age, in the twenty-three years, 1856–1878, were 31.44 per cent. of the whole number; of foreign parentage, the decedents of this class were 44.38 per cent. of all. From 20 to 30 years of age, and from 30 to 40, the percentage of decedents was larger among the foreign; but beyond that age, the larger percentage is always of American parentage, 39.61 per cent. of the decedents living to the age of 40 years or more, while of the foreign decedents only 25.32 per cent. lived beyond that age.

By comparing Tables XVI. and XVII., it will be found that the percentage of decedents less than five years old, in 1878, was 2.08 per cent. above the average for the whole period of twenty-three years; the decedents of American parentage, 5.34 per cent. above the average; and the decedents of foreign parentage, 1.74 per

cent. below the average. Of those who lived to the age of fifty years or more, the percentage of American parentage in 1878 was 31.04 years, or 0.78 per cent. below the average of that class, as found in Table XVII.; the percentage of foreign parentage was 18.55, or 0.84 per cent above the average. This corresponds with the facts of Tables XIII. and XIV., that the average age of all the decedents of American parentage in 1878 was 2.44 per cent. below the average for the period of twenty-three years, 1856–1878; and the average age of all the foreign decedents in 1878 was 0.27 per cent. above the average.

Deaths by age and parentage in each month and quarter of the year. Table XVIII. gives the number of deaths at each age and of all ages, according to parentage, in each month and quarter of the year 1878, and shows the comparative mortality at different seasons of the year in the two classes of the population. The whole number of deaths during the year was 1,989; of American parentage 889; of foreign parentage, 1,100; divided as follows in the different quarters of the year:—

	1st quarter.	2d quarter.	3d quarter.	4th quarter.
American parentage	26.88	21.26	26.66	25.20
Foreign parentage	27.45	21.27	25.73	25.55

As has already been stated in the remarks on Table XII., page 25, the year 1878 is an exception to the general rule, in that the largest percentage of mortality was in the first quarter of the year, among both classes of decedents. It is also contrary to the usual order to find a larger percentage in the third quarter, among the decedents of American parentage than among the foreign. By the percentages given on page 32, it will be seen that in a long period of time it is the reverse of this. Table XVIII. shows also the mortality at different ages. Of decedents of American parentage, there were 237 in the third quarter; of this number, 70, or 29.53 per cent., were under one year old; this percentage is a little less than in 1877; the number of decedents of American parentage, under one year old, in the whole year 1878, was 168; of this number the 70 who died in the third quarter were 41.66 per cent., or 4.02 per cent. less than in 1877. The number of decedents of American parentage in the first quarter of the year was 239; of this number, 43, or 17.99 per cent., were less than one year old; this is 3.78 per cent. more than those of the same class in the first quarter of 1877; of the 168 children under one year old, who died in the course of the year, the 48 who died in the first quarter were 25.60 per cent., or 8.31 per cent. more than in 1877.

Among the decedents of foreign parentage, these statistics are as follows: number of decedents in the third quarter, 283; number under one year old, 86, or 30.38 per cent., 0.11 per cent. less than in 1877; number of decedents under one year old, in the whole year, 205; number in third quarter, 86, or 41.95 per cent., 3.86 per cent. less than in 1877. Number of decedents in the first quarter, 302; number under one year old, 49, or 16.22 per cent., 2.47 per cent. less than in 1877. Number of decedents under one year old in the year 1878, 205; number in the first quarter, 49, or 23.90 per cent.; 4.20 per cent. more than in 1877.

Causes of death; sex, age, parentage, and percentage from each cause. Table XIX. gives the number of deaths in 1878 from each cause, according to sex and parentage, and in each division of age; also the percentage from each cause, of all from known causes.

Season and disease. Table XX. also gives the number of deaths from each cause, but is different in detail, from Table XIX. It gives the number from each cause in each month and quarter, and in the whole year, and shows the comparative mortality from different diseases at different seasons.

Table XVIII. Deaths by age and parentage in each month, 1878.

					Di	VISI	ONA	L PI	ERIO	DS O	F LI	FE.					
1	YEAR.	Under 1.	1 to 2.	2 to 5.	5 to 10.	10 to 15.	15 to 20.	20 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.	80 to 90.	90 to 100.	100 & over	Ave Acres
-	January February March	18 15 10	12 6 7	8 10 8	8 2 11	2 1 1	2	3 3 5	5 4 5	7 5 3	9 6 8	11 4 9	8 2 7	6 3 1			
ı	1st quarter	43	25	26	21	4	4	11	14	15	23	24	17	10	2		
	April	13 9 5	7 2 1	10 7 4	3 3	2 4 2	1 3 4	9 4 3	4 1	4 4 6	3 4 5	6 7 4	9 1 10	7 4 2	2		1
	2d quarter	27	10	21	10	8	8	16	9	14	12	17	20	13	4		ı
The same of the sa	July	39 19 12	4 7 7	5 2 5	2 2 4	2	2 3 4	9 6 7	5 1 7	6 6 4	5 4 9	5 6	9 5	3 3 2		···	
	3d quarter	70	18	12	9	5	9	22	13	16	18	17	18	8	1	1	000
	October November December	9 7 12	7 4 7	5 9 15	8 4 6	2 2 6	1 1 3	5 6 6	2 5 6	4 5 6	5 5 9	3 6 8	6 8 10	2 2 5	1		
	4th quarter	28	18	29	18	10	5	17	13	15	19	17	24	9	2		1
	Whole year	168	71	88	58	27	26	66	49	60	72	75	79	40	9	1	-
	January February March	12 24 13	14 17 14	24 17 14	9 7 4	2 2 1	6 4 2	6 6 9	10 9 5		11 8	8 5 4	9 3				
	1st quarter	49	45	55	20	5	12	21	24	9	23	17	18	3	1		,
	April	20 9 12	5 3 5	12 6 11	6 3 5	1 4 2	1 3 4	6 10 7	7 15 9	5 5 7	4 7 5	4 2 4	11 4 6	2	1		
	2d quarter	41	13	29	14	7	8	23	31	17	16	10	21	3	1		1
	July August September	43 35 8	8 13 8	12 11 5	7 4 3	4	3 2 3	1 13 6	13 8 11	12 6 5	5 7 6	3 2 2	3 4	1 2			-
	3d quarter	86	29	28	14	- 5	8	20	32	23	18	7	10	3			
	October November December	12 11 6	9 4 3	20 20 9	10 12 10	5 1 2	522	11 13 15	11 11 4	3 7 10	3 8 5	5 5 7	4 5	1 1 2	1	1	-
	4th quarter	29	16	49	32	8	9	39	26	20	16	17	13	4	2	1	
	Whole year	205	103	161	80	25	37	103	113	69	73	51	62	13	4	1	1,
	otal Amer. and For.	373	174	249	138	52	63	169	162	129	145	126	141	53	13	2	1,

from each cause to whole mamber from known Ratio of deaths .latoT PARENTAGE Discuse, sex, age, parentage, and percentage from each cause. Foreign. American. • pun 06 So and nader 90. under 80. pun 02 under 70. DIVISIONAL PERIODS OF LIFE. ban 00 50 and ander 60. 40 and or 50, under 50. under 40, bun og **** **** **** pan oz 15 and under 20. 10 and ander 15. under 10, pus g under 5, pur z 1 and under 2. Year. Under 1 JefoT. B-400 Childbirth
Puerperal Convulsions
Cholera Infantum.
Cholera Infantum. Males. Apoplexy
Addren
Addren
Balder
Brain, Congestion of
Brain, Diseases of
Inflammation of
Bronchilis
Cancer, various
of Stomach Falls Poleoning Rallroads.... (various)
Asphyxia
Burns and Scalds Aneurlsm..... CAUSES OF DEATH. TABLE XIX. Abscesses... Childbirth

DEATHS, 1878.

Table XIX.—Continued. Deaths, 1878. Disease, sex, age, parentage and percentage from each cause.

		SKX.			-	Q	IV18	LONA	DIVISIONAL PERIODS OF LIFE	SR10	DS C	J. J.	IPE.	-		1	PAI	PARENTAGE	GE.	mber mye
CAUSES OF DEATH.	Males.	Femules.	Total	Luder 1	I and pinder 2, ban 2	.ë rabnu bna ë	for and for and	ban di	20 and 20.	30 and	40 and 40, 40 and under 50,	50 and under 60.	60 and 40 and 60 and	70 and under 80.	80 and under 90.	90 and	American.	Foreign.	Total.	
Marrhera, Chronic	1368	+011 8		i= :	:05	1 :000	1.01	1:4:	. DE 1				:		::	Lili	105	141	24	51
Smphysema. Sateritis Caltensy	-1-4	-10	n g c				:-	: :-	* PF :	- 01		- 04 04	!!		: :	Ш	10010	101	1120	16.
Erysipelas Exposure and Neglect	(00 OI)	140)	P-41	01.01					11								200 H	4400		
	38.	2 a 2 .	2010	iii	iii	* : :	• ! !	N :00	2 01	: -	:				:		4 . 2	5 24.80	2000	3.96
Heat. Hemorrhage	+									:	::			11						
30	25 64 5			П		Ħ			1				:	11	Ш		- 00			
doping Cough.	17.00	24.24	328	14	100	100	10100		Щ	ш	Ш	:	111	Ш	Ш	Ш	ដន	929		2.18
lydrophobia. nsanity. ntemperance	H 50 4	. # °	121	Ħ			: :	4		: 01	. 02 0		100				F	-66	-4.	
Delirium Tremens	(B) FF	100	09 01		İī	100		11	11	44	ii,		:	11	H		DE-10	-	01017	
Bright's Disease of	200	13	1 33 -	1	i	1	11	::	-	4 00	2	-1-	010	110	2.00		17	16		1.67
Aver Diseases of Inflammation of	100	. 20	1200		ń					_	. 04			:	· ·		9-		108	
daloga, Diseases of faiformations	4.58		- N	108	-	-		#	11		11		11		11			2001	-	

TABLE XIX.—CONTINUED.	DEATHS, 1878.	18, 1	878.	7	Jise	486,	se	, a	ا ۾	nar.	en te	10e	as	ž.	rce	2	ge J	Disease, sex, age, parentage and percentage from each cause	each	Car		1
		BKX.				1	1710	NOI	DIVIAIONAL PRRIODS OF LIPE	KK	800	0	LIFE					PARENTAGE.	TAG		icathi ores to mber	u.
CAUSES OF DEATH.	Males.	Females.	Total.	Under 1 Year.	bas t S 19bau bas 2	.č rəbau bas č	.01 19ban San 01	.dl 19bar bas dl	.08 rebau bas 08	.06 raban 30 and	.04 19bag bas 04	.05 19bag bas 05	.08 reban bas 08	.07 19bau bas 07	.08 rebau bas 08	.09 19bau bas 08	American.		Foreign.	Total.	Trom each ca	trom kno
Menlettis, Cerebro Spinal Murder and Homidde	84-8	a≈ :⊈:	85-2	7	2:::	9 : : :	60 N	-6		<u> : ::</u> 	<u> </u>		<u> </u>	: : : : : : : : : : : : : : : : : : :	: : : : : : : : : : : : : : : : : : :	: : : : : : : : : : : : : : : : : :	 	<u>:</u> ഇന-ജ	8 · 8	82-2		2888
Paralysis Peritonitis Franciusy Preumonia	29 : 3	¥ <u>r-8</u>	12-12		<u> </u>	 -			:- :-		- i A : F	ଷର ଅ	 	<u>~:</u>	<u> </u>	: : :	::::	22: 22	er-2	¥ 2- 2		1258
Congestion of L	α : σ×		3-25	= : :	- 	≅ ::8		•	:::-	::: -	:::	::-		_ 	*		+	<u> </u>	<u>a-n</u>	2-05		3,8,8,8
Serofula Septicemia Skin, Diseases of	3887		F 01 80	N	- : :	[] 	1 : : :	म	· N : :	: : : :					-		::::	• • • • •		- www		855
Spine, someon,	- 81 - 4	. N M C	<u> </u>				N -			;;;; ;		::- :	::	: : 	<u> </u>		<u>:</u> ::::	N	- 000	-445		ន់ន់ន់ដ
	4	88	K8-41							e : : :	A :	- ev	- : : :					8777 7	هم . ۵۰	V 3141		ន់ដន់ន់ន
Tetains and Trismas Naccentium Throat, Diseases of Thrush Thrush Tuberculosis	N- 0-	2	-8Fe	N HAH				81		· · · · · ·		্ মূল		N						<u>850</u>		ទទន់ខ្លួន
Uterus, Diseases of		10 00 :	-50	:==		8			:::	- ; ;	- : :	N : :	- 	: : : : : : : : : : : : : : : : : : :	<u> </u>		:::	∞∞ -	۳ ۾ :	99-		8
Totals	8	1,020	1,980	2	2	2	8	8	8	8	162 129	2	155	126 141		13	192	880 1,	1,100	1,989	2	100.00

a

867-1951428-25185-9782885-282858-2928-9858

CAUSES OF DEATH.	January.	February.	March.	First Quarter.	April.	May.	Jane	Second Quarter.	.ylnt.	August.	September.	Third Quarter.	October.	Хочетьет.	December.	Fourth Quarter,
Accidents, (all kinds). Apoplexy. Brain, Diseases of Bronchtits. Cancer, (all kinds). Child-burth and Puerperal Diseases. Cholese Infuntum	00041-10	10000000		2428440	0.47-0 :01	800000	0004004	122822004	Second			1 2 2 4 · V 2 5		व्यक्तक का क		0210244
Cholera Morbus. Consumption. Convulsions.	72	21	24	72	20	. 25.01	25.		-20		65			1		
Croup Diarrhera, Acute and Chronic. Disturberia	40,4	19: 0	2.0	20.20	9-18-		100	02.45			11"			# N N N		
Enterities Erystpitas Ferer, Tynhoid	OF OF		8	1000			iii [*]	2								
Heart, Diseases of Hooping Cough Hydrocephalus	1.00	644	400	723			2000	an I		018		200			500	
Insunity Interperance and Delirium Tremens Kidneys, Diseases of. Liver, Diseases of.	. 0.00	- 1	01.10											:01 50 01	0 400	
Marasmus Measles Menlightis, Cerebro-Spinal. Old Age and Senile Guarene	"F "	10 St - 4	10 M 101 10						-11		Ш	_ ! !		- 11	11	1
ion of	400	77	ra co m	5000		04-100	0170	4408	:		90			i E		3200
Suidde Teething Unknown. Other others a given	1 - 25	Hene			1 7 7 8		1111	8	11	1		gm++g				
	900	126	AGN	641	164	1	326	1	808	1	140	1°	1	1-	1	19

NOTES ON THE CAUSES OF DEATH,

DURING THE YEAR 1878.

These notes explain more particularly some of the facts that are given in Tables XIX. and XX., and also give some additional statistics; and compare these with the statistics for the twenty-three years, 1856–1878 inclusive.

The percentage of deaths from the different causes is always reckoned upon the number from known causes.

The whole number of deaths in Providence, in 1878, was 1,989, of which 16, or 0.80 per cent. were from unknown causes; this is 0.59 per cent. more than in 1877. The number of which the cause was stated, and on which all the percentages are reckoned, was therefore 1,973. During the twenty-three years, 1856–1878, the number of deaths was 30,473, of which 524, or 1.72 per cent. were from unknown causes, and 29,949 were from known causes. During the thirty-nine years, 1840–1878, the number of deaths was 43,618, of which 1,526, or 3.50 per cent. were from unknown causes, and 42,092 from known causes.

Table XX. gives the number of deaths from different causes in each month and quarter of the year, and shows what diseases prevail in the different seasons. A large proportion of the deaths from bronchitis occurred in the first quarter of the year; also of the deaths from measles, pneumonia and congestion of the lungs; about two-thirds of the deaths from diphtheria were in the first and last quarters of the year; the deaths from consumption were very evenly distributed through the year; the deaths from all diarrhocal diseases, were chiefly in the third quarter; as usual, and the deaths from typhoid fever, in the third and fourth quarters of the year.

Abscesses. The number of deaths from abscesses in 1878, was 5, or 0.25 per cent. of all from known causes; this was one more and 0.04 per cent. more than in 1877. They were specified as

follows: one psoas abscess, one abdominal, one retro-pharyngeal, and one each of the neck and the bladder; besides these there were others which were reckoned under the head of disease of the organ affected, as "disease of the brain, lungs or liver." The 5 decedents were 3 males and 2 females; all of foreign parentage. The statistics for the 23 years, 1856-1878, are as follows:—

Number of deaths from all known causes, 29,949.

From abscesses: males, 47; females, 39; total. 86, or 0.29 per cent. of all.

American parentage, 36; foreign parentage, 50.

Accidents. The number of deaths resulting from accidents in 1878, was 50, or 2.53 per cent. of all from known causes. They are recorded in the table under the following heads: asphyxia, 2; burns and scalds, 7; drowning, 16; falls, 7; poisoning, 3; railroad, 3. Besides these, 12 were from various causes not specified in the table; one was killed by the fall of a bank of earth; one by the fall of a building; one by blasting; one was run over by a dray; one caught on a shaft; three by injury to the brain, in what way received was not specified; one by injury to the spine. Of those by poisoning, two were by opium and one from an overdose of laudanum.

The statistics as to sex and parentage were as follows:-

Males, 38; females, 12; total, 50, or 2.53 per cent. of all. American parentage, 20; foreign parentage, 30.

For the twenty-three years, 1856-1878:-

Males, 655; females, 210; total, 865; or 2.89 per cent. of all.

American parentage, 344; foreign parentage, 521.

Aneurism. There were 4 deaths from aneurism in 1878, or 0.20 per cent. of all from known causes; in two cases it was aneurism of the aorta; in one, popliteal, and in one it was not specified; the decedents were all males of foreign parentage. The statistics for the twenty-three years, 1856-1878, are as follows:—

Males, 20; females, 3; total, 23, or 0.08 per cent. of all. American parentage, 8; foreign parentage, 15.

Apoplexy. This was reported as the cause of 42 deaths in 1878, or 2.13 per cent. of all from known causes. Males, 25; females, 17; American parentage, 30; foreign, 12. (Vide "Paralysis.")

The following are the statistics for 23 years, 1856-1878:—

Males, 301; females, 319; total, 620, or 2.07 per cent. of all. American parentage, 446; foreign parentage, 174.

Brain, Diseases of. The number of deaths reported from diseases of the brain in 1878, was 67, or 3.40 per cent. of all from known causes; this was 15 less, and 0.85 per cent. less than in 1877; 7 were from "congestion"; 42, from "inflammation." The number of each sex and parentage, as follows:—

Males, 37; females, 30; total, 67, or 3.40 per cent. of all. American parentage, 27; foreign parentage, 40.

For the twenty-three years, 1856-1878:-

Males, 638; females, 526; total, 1,164, or 3.89 per cent. of all.

American parentage, 613; foreign parentage, 551.

Bronchitis. There were 47 deaths from bronchitis in 1878, or 2.38 per cent. of all from known causes. Of these decedents, 20, or 42.55 per cent., were less than one year old.

Males, 18; females, 29; total, 47, or 2.38 per cent. of all. American parentage, 19; foreign parentage, 28.

For the twenty-three years, 1856-1878:-

Males, 160; females, 209; total, 369, or 1.23 per cent. of all.

American parentage, 164; foreign parentage, 205.

Cancers. The number of deaths from cancers in 1878 was 48, or 18 less than in 1877, and 2.44 per cent. of all. Of these decedents, 9 had cancer of the breast; 8, cancer of the stomach, and 10, cancer of the uterus. The 21 included in the table under the head of "various," were as follows: rectum, 3; liver, 2, and one each of the throat, eye, tongue, kidney, thigh, pylorus, bowels, pancreas, sub-maxillary glands, testicle, left femur, and one ovarian; four were not specified. The decedents were divided as to sex and parentage as follows: males, 16; females, 32; American parentage, 26; foreign 22.

The statistics for the 23 years, 1856-1878, are:

Males, 179; females, 516; total, 695, or 2.32 per cent of all.

American parentage, 461; foreign parentage, 234.

The deaths in each of the last nine years have been:-

	Number.	Per cent
1870		19.06
1871		15.66
1872		
1873		13.57
1874		13.76
1875		15.79
1876	284	15.34
1877		15.22
1878		15.46
		- -

The number and percentage of the decedents from consumption in each division of age, were as follows for the year 1878, and for the period of 23 years, 1856 to 1878 inclusive:—

Periods.	Under 1 year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	70 and ander 80.	80 and under 90.	TOTAL
1878. Number	.33	1.31	.33	1.31	2.95	31 10.16	97 31.80	70 22.95			100	12 3.93	1000	305 100.00
1856–1878. Number	63		54	53	99	100	1,556 30.66		1.0	-		127	11	5,075

The largest percentages are between the ages of 20 and 50 years; 66.23 per cent. of the decedents in 1878 being within those limits; 51.81 per cent. lived beyond the age of 30. The average age of decedents from all causes in 1878 was 26.09 years. Of the decedents from consumption in the twenty-three years, 1856–1878 inclusive, 53.65 per cent. lived beyond the age of 30 years; the average age of all decedents during that period was 27.53 years.

Convulsions. There were 56 deaths from convulsions in 1878, or 2.84 per cent. of all from known causes. This was 18 more deaths and 0.87 per cent. more than in 1877. As usual, most of these deaths were of very young children, 54 of the decedents being under 5 years old; one was between 5 and 10 years; and one between 15 and 20.

Males. 33; females, 23; total, 56, or 2.84 per cent. of all. American parentage, 20; foreign parentage, 36.

The statistics for twenty-three years, 1856-1878, are as follows:

Males, 438; females, 351; total, 789, or 2.63 per cent. of all.

American parentage, 326; foreign parentage, 468.

Croup. There were 38 deaths from this cause in 1878; 2 less than in 1877, and 1.93 per cent. of all from known causes. The decedents were as follows in regard to age, sex and parentage: under 1 year, 4; between 1 and 2 years, 11; between 2 and 5 years, 18; between 5 and 10 years, 5. Males, 18; females, 20; American parentage, 18; foreign parentage, 20.

For the twenty-three years, 1856-1878, the statistics are:-

Males, 350; females, 346; total, 696, or 2.32 per cent. of all. American parentage, 285; foreign parentage, 411.

Other statistics in relation to croup are given hereafter, in comparison with those of diphtheria.

Debility. This was reported as the cause of 37 deaths in 1878, including 12 from premature birth; the decedents were 33 under one year old; one between one and two years, and the other three were 50 years or over. The statistics are:—

Males, 20; females, 17; total, 37, or 1.88 per cent. of all. American parentage, 23; foreign parentage, 14.

For the twenty-three years, 1856-1878:-

Males, 287; females, 253; total, 540, or 1.80 per cent. of all. American parentage, 224; foreign parentage, 316.

Diarrhoea and Dysentery. The number of deaths from these diseases in 1878 was as follows:—

	Male.	Female.	American.	Foreign.	Total. Percentage.
Acute Diarrhœa	12	9	12	9	211.06
Chronic Diarrhea	3	4	5	2	70.36
Dysentery	3	8	6	5	110.56
•	_	_	_	_	
Total	18	21	23	16	391.98

The number of deaths was 16 less than in 1877; the percentage, 0.87 less. It is very remarkable to find the number of decedents of American parentage out-numbering those of foreign parentage,

as the reverse of this has generally been true. In this connection may be given the statistics concerning the deaths from the following diseases of a like nature, in 1878:—

Cholera Infantum			American.	•		_
Cholera Morbus	2	1		3	3	0.15
Total	_		_		_	

Combined with the figures already given on page 49, it makes the total of deaths from these five diarrheal diseases in 1878, as follows:

Males, 62; females, 68; total, 130, or 6.59 per cent. of all. American parentage, 57; foreign parentage, 73.

The number of deaths is 74 less than in 1877; the number in that year was 204, or 10.56 per cent. of all. In 1878 there were 44 less males; 30 less females; of American parentage, 23 less; of foreign parentage, 51 less.

The statistics of these five diseases for the twenty-three years, 1856-1878, are as follows:—

Number of deaths, 3,495; annual average, 152.

Males, 1,750; females, 1,745; total, 3,495, or 11.67 per cent. of all.

American parentage, 1,416; foreign parentage, 2,079.

The proportion of decedents of foreign to those of American parentage, in the twenty-three years, was 147 to 100, or 59.48 of foreign, and 40.52 of American parentage in each 100. One or two reasons for this may be given. The most of the decedents from these causes are generally young children; thus in 1878, 93 of the decedents, or 71.54 per cent., were less than two years old; and the proportion between the two classes, 128 of foreign to 100 of American parentage, or 56.15 of foreign, and 43.85 of American parentage in each 100. But the number of children born of foreign parentage is also much larger than the number of American parentage. In 1878 the number of children born was 2,627; of American parentage, 1,215; of foreign parentage, 1,412, or 116 of foreign to 100 of American parentage; or 53.75 of foreign and 46.25 of American parentage in each 100.

In the twenty-three years, 1856-1878, the number of children born was 43,900; of American parentage, 19,863; of foreign

parentage, 24,037; or 121 of foreign to 100 of American parentage, or 54.75 of foreign, and 45.25 of American parentage in each 100. The proportions of the two classes of decedents during this period have already been given, and it is easy to see why the foreign should so largely exceed the American, since there are so many more children living of the former than of the latter class.

The following table shows the number of deaths from each of these causes, with the number from the whole class of causes, and their percentage of all from known causes, reported in Providence during the period from 1840 to 1854 inclusive; during each year, from 1855 to 1878 inclusive, and in the aggregate during the whole period of 39 years, 1840 to 1878 inclusive:—

5	•	entery. Enteritis. Com 796	
4 5 2 1 5 1 5	37	38 35	140
5 2 2 1 2 5 5	29	35	13211.1 12413.4 989.6 798.7 11811.7 12611.9
2 2 1 2 5 5	38 27 28 21 22 31 35	29 5 12 8 13 4 18 9 32 6 22 9	124
2 1 2 5 5	27 28 21 22 31	12 8 13 4 18 9 32 6 22 9	98
1 2 5 1	28 21 22 31 35	13 4 18 9 32 6 22 9 77 9	798.7 11811.7 12611.9 10711.7
2 5 1 5	21 22 31 35	18 9 32 6 22 9 77 9	11811.7 126 11.9 10711.7
5 1 5	22 31 35	32 6 22 9 77 9	126 11.9 10711.7
1 5	31 35	22 9 77 9	10711.7
5	35	77 9	
			19215.8
2	20		100 101
_	~ 4	40 13	16012.4
7	34	62 4	15712.9
17	28	44 9	14513.9
2	15	. 43 10	
3	24	15 8	12010.8
6	24	23 6	124 9.8
3	22	16 15	14911.9
6	24	16 18	
10	52	35 14	
			165 9.6
6			
4			20611.1
6	34	21 21	20410.5
	28	11 18	130 6.5
	6 3	3 50 6 36 4 40 5 34 3 28	3 50 17 25 6 36 14 31 40 22 18 6 34 21 21

It will be noticed that the number of deaths from each of the five causes was less in 1878 than in the previous year; and the percentage from the five causes combined was less than in any other year given in the table.

The number of deaths from known causes in the thirty-nine years, 1840-1878, was 42,092.

Deaths from Asiatic cholera are not included in these statistics. They have been as follows: in 1849, 158; in 1854, 159; in 1866. 14; total, 331.

Diphtheria. This disease, though not causing quite so many deaths as in 1877, still prevailed in 1878 to so great an extent as to cause serious anxiety, and much inquiry as to the cause, and the means, if possible, of prevention. The number of decedents from it was 246, which was 49 less than in 1877; the particulars were as follows:—

Males, 136; females, 110; total, 246, or 12.47 per cent. of all.

American parentage, 105; foreign parentage, 141.

These deaths were in the different seasons of the year, as follows: first quarter, 81; second, 46; third, 32; fourth, 87. The mortality continues to be mostly among children, 226 of the decedents or 91.87 per cent., being less than ten years old. The first cases of diphtheria reported in Providence were in 1858; for the twenty-one years, 1858–1878 inclusive, the statistics regarding it, are as follows:—

Deaths from all known causes in 21 years, 28,021.

From diphtheria, 986; annual average, 47.

Males, 497; females, 489; total, 986, or 3.52 per cent. of all.

American parentage, 473; foreign parentage, 513.

The 246 deaths from diphtheria in 1878 are more than five times the annual average, and form 24.95 per cent. of the whole number from this disease in the twenty-one years.

In the following table are given the statistics as to sex, parentage, season and locality, for each of the twenty-one years, 1858-1878, and for the whole period:—

DEATHS FROM DIPHTHERIA IN PROVIDENCE, TWENTY-ONE YEARS.

БіритневіА.	1858.	1859.	1860.	1801.	1862.	1863.	1864.	1805.	1866.	1867, 1868.		1860, 1870.		1871.	1872	1873.	1874.	1875.	1876, 1877.		1878.	Total.
Whole number	00.00	1.96	8.6	272	1.82	3.54	=17	1.35	=8	0.03	0.46	28	0.80	198	221	88	88.	14 0	111	2816	246	3,52
MalesFemales		000	21	212	43	875	85	51.4	100	62.00	10	00.00	100	210	120	20	99	600	23	25	136	407
PARENTAGE: AmericanForeign	क्रम	==	0.9	510	11.0	80	28	99	610		10	101%	40	42	28	13.8	91	# 9	28	129	192	473 513
	П	O1 HH 410 22	01-H 010-000	स्त्र स्त्र स्त्राध्यम्	∞ H H H − ∞ ∞	10 PRESENTED 11 4100	01010000000000000000000000000000000000	- 4	H HM H 100M	a- - -	00 14 1044		H HHMM 10		∞404- 2 104 4		0101H 010H H001H	w	ZSgonmonor; a	おおおおおいったおおおお	生まなお言る。自己的問題	12588389898981
Docality: Ward II. III. IV. VII. VIII.	-E- 000	121 00 2000 100 4	Q+0 00+	ннюноюн	# 20 + 0 m	residence		- 01010001-	H	, ox 60	юнннии 4	60	# : #HHHH00	0-01-00000	0-3c253	H NAH 120		901240010F30	8222222288	8422532258	56888886116	

The subject of the connection of season with the mortality from diphtheria is exceedingly interesting and important of itself, and still more so when the mortality from this disease is compared with that of scarlatina and croup. The following table shows the number and the percentage of deaths in Providence, from diphtheria, scarlatina, and croup, in each month and quarter of the year. The figures for scarlatina and croup include the aggregate mortality from these diseases during the twenty-four years, from 1855 to 1878 inclusive. The figures for diphtheria are for twenty-one years, from 1858 to 1878 inclusive.

		HERIA. -1878.		ATINA. 1878.	CROUP. 1855-1878.		
Монтня.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	
January	91	9.23	194	12.52	92	12.71	
February	64	6.49	172	11.10	84	11.60	
March	77	7.81	150	9.68	78	10.78	
1st quarter	232	23.53	516	33.30	254	36.09	
AprilMay	67	6.79	142	9.17	60	8.29	
May	62	6.29	147	9.49	40	5.52	
June	57	5.78	141	9.10	25	3.45	
2d quarter	186	18.86	430	27.76	125	17.26	
July	36	3.65	91	5.88	27	3.73	
August	37	3.75	65	4.20	11	1.52	
September	85	8.62	69	4.46	40	5.52	
3d quarter	158	16.02	225	14.54	78	10.77	
October	152	15.42	92	5.94	69	9.53	
November	145	14.71	119	7.08	92	12.71	
December	113	11.46	167	10.78	106	14.64	
4th quarter	410	41.59	378	24.40	267	36.88	
Totals	986	100.00	1,549	100.00	724	100.00	

Dividing the year into periods of four months, the numbers and proportions are as follows:—

Seasons.	DIPHT	HERIA.	SCARL	ATINA.	. CROUP.		
SEASUNS.	Number.	Per cent.	Number. Per cent.		Number.	Per cent.	
1st 4 months	192	30.32 19.47 50.21	658 411 417	42.47 28.67 28.86	314 103 307	43.38 14.22 42.40	

Dividing the year into two periods, the numbers and proportions are:—

SEABONS.	DIPHT	HERIA.	SCARL	ATINA.	· CROUP.		
DEADURG.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	
1st 6 months	418 568	42.39 57.61	946 603	61.06 38.94	379 345	52.35 47.65	

The general results of these figures are the same as given in previous reports, showing marked and characteristic differences in the three diseases, in respect to the season of their prevalence and mortality.

Diphtheria shows a large excess of mortality in the fall of the year, more than half of all the deaths from this disease occurring in the last four months of the year; and 57.61 per cent. of all, in the last six months.

Scarlatina shows a large excess of mortality in the first part of the year, 42.47 per cent. of all the deaths occurring in the first four months of the year, and 61.06 per cent. of all, in the first six months.

Croup shows but a slight difference between the first quarter and the last quarter of the year, the mortality evidently depending more upon severity of cold than upon other causes.

In the aggregate for the whole period included in the tables, the smallest monthly mortality from scarlatina and croup, was in August; from diphtheria the mortality was slightly less in July than in August. The greatest monthly mortality was: from diphtheria, 15.42 per cent., in October; from scarlatina, 12.52 per cent. in January; from croup, 14.64 per cent. in December.

If we notice the preceding table, "Deaths from Diphtheria in twenty-one years," we shall see that though there were some deaths every year from the beginning of the disease in 1858 to the year 1875, the number each year was small.

There were only five deaths from the disease in each of three years, during the period, and the greatest number in any year was forty-two. The whole number of deaths from diphtheria in the 18 years from 1858 to 1875 inclusive, was only 334, or an average of 18.5 annually. But in 1876 a severe epidemic of the

disease began in Providence, and in the three years from 1876 to 1878 inclusive, there were 652 deaths from diphtheria, or nearly twice as many in three years as in the 18 years previous.

It naturally became an interesting question to know the difference in the statistics of the disease when epidemic and when not epidemic. I have, therefore, had the following table prepared which shows, side by side, certain statistics of diphtheria in Providence, first, for 18 years, from 1858 to 1875 inclusive, when the disease was not at any time epidemic; second, for the three years, from 1876 to 1878 inclusive, when the disease prevailed as a very severe epidemic, and third, for the two periods combined:—

		HERIA. -1875.		HKRIA. -1878.	DIPHTHERIA. 1858–1878.		
Months.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	
January	31 32	9.28 9.58	60	9.20 4.91	91 64	9.23 6.49	
March	32	9.58	45	6.90	77	7.81	
1st quarter	95	28.44	137	21.01	232	23.53	
April	19	5.69	48	7.36	67	6.79	
May	15	4.49	47	7.21	622	6.29	
June	23	6.89	34	5.21	57	5.78	
2d quarter	57	17.07	129	19.78	186	18.86	
July	19	5.09	17	2.61	36	3.65	
August	10	2.00	27	4.14	37	3.75	
September	25	7.49	60	9.20	85	8.62	
3d quarter	54	16.17	104	15.95	158	16.02	
October	40	11.98	112	17.18	152	15.42	
November	45	13.47	100	15.34	145	14.71	
December	43	12.87	70	10.74	113	11.46	
4th quarter	128	38.32	282	43.26	410	41.50	
Totals	334	100.00	662	100.00	986	100.00	
Males	162	48.50	335	51.38	497	50.71	
Females	172	51.50	317	48.62	480	49.29	
American	196	58.68	277	42.48	473	47.97	
Foreign	138	41.32	375	57.52	513	52.03	

It will be seen that the percentages of deaths in the different months vary greatly in the first two columns; January is nearly the same in both columns; February and March show a greatly reduced mortality, as compared with other months, when the disease is

30.32

19.47

50.21

epidemic; April and May show an increased mortality, and June and July a largely decreased mortality when the disease is epidemic. After July there is an increased proportional mortality, when the disease is epidemic, in each month until December. By quarters of the year there is a decreased comparative mortality when the disease is epidemic, in the first and third quarters, and an increased mortality in the second and fourth quarters.

By periods of four months the percentages of mortality were as follows:—

1858	1875.	1876-	1878.	1858–1878.		
Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	

DIPHTHERIA.

By periods of six months, the percentages of mortality were as follows:—

34.13 20.06 45.81

114

67 153

DIPHTHERIA.

Seasons.	1858	1875.	1876-	-1878.	1858–1878.		
SEASONS.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	
1st 6 months 2d 6 months		45.51 54.49	266 386	40.79 59.21	418 568	42.39 57.61	

These figures show a marked difference in the season of mortality from diphtheria when the disease is epidemic, and when not epidemic. They also confirm and make more marked the general law in regard to the season of diphtheria as stated on page 55 of this report, and in previous reports.

If we refer again to the table, page 56, we find, in regard to sex and parentage very great differences in the mortality of diphtheria, when epidemic and when not epidemic, the proportions being almost exactly reversed. Thus in the first period, when diphtheria was not epidemic, the decedents were 48.50 males, and 51.50 females in each 100; but in the second period when the disease

SEASONS.

1st 4 months.....

3d 4 months.....

months.....

was severely epidemic, the decedents were 51.38 males, and 48.62 females in each 100. So also in respect to parentage, when the disease was not epidemic, there were 58.68 decedents of American, and 41.32 of foreign parentage in each 100; but when the disease was epidemic, the figures were almost reversed, and there were 42.48 decedents of American, and 57.52 of foreign parentage in each 100.

The subject of the causes of diphtheria is one of the most important that can occupy the attention of those interested in the public health, at this time. Any statistics that may possibly throw light upon the subject are therefore important.

Exposure and Neglect. Four deaths in 1878 were ascribed to exposure or neglect; two of the decedents were children under one year old; one was between 50 and 60 years, and one between 70 and 80. Two were males, and two females; 1 of American, and 3 of foreign parentage; total 4, or 0.20 per cent. of all from known causes.

Fever, Typhoid. There were 47 deaths from this cause in 1878; 4 more than in 1877, and 14 more than in 1876.

Males 19; females, 28; total, 47, or 2.38 per cent of all. American parentage, 27; foreign parentage, 20.

In the following statistics for the twenty-three years, 1856-1878, besides the deaths from typhoid, are included a few that have been reported in some years from "fever," "bilious fever," and about 30 from "typhus fever":—

Whole number, 982; annual average, 43.

Males, 508; females, 474; total, 982, or 3.28 per cent. of all.

American parentage, 544; foreign parentage, 438.

Heart, Diseases of. The number of deaths from diseases of the heart in 1878 was 83; five of which were specified in the table as "hypertrophy"; the whole number was 10 less than the number in 1877.

Males, 39; females, 44; total, 83, or 4.21 per cent. of all. American parentage, 45; foreign parentage, 38.

During the twenty-three years, 1856-1878, the number of deaths from diseases of the heart, was 1,367, or 4.56 per cent. of all

from known causes; an annual average of 59. The following are the particulars for each year:—

	81	x.	PAR	ENTAGE.	Whole		
	Males.	Females.	American.	Foreign.	Number.	Percentage	
856	8	13	16	5	21	2.03	
857	13	18	22		31	3.45	
1858	13	23	28	8	36	3.61	
1859	13	11	17	7	24	2.76	
1860	15	16	21	10	31	3.23	
1861	23	18	29	12	4l	4.05	
1862	26	23		16	49	5.54	
1863	23	19	30	12	42	3.54	
1864	26	82	44	14	58	4.63	
1865	18	29	33	14.	47	3.95	
1866	19	21	27	13	40	3 . 95	
1867	26	23	35	14	49	5 . 19	
1868	25	27	34	18	52	4.79	
1869				27			
1870	28	31	33	26	59	4.72	
				31			
1872				29			
1873	46	54	59	41	100	5.88	
				42			
				41			
				37			
				40			
				38			
	_			-			
23 years' to	tal.654	713	863		1,367	4.56	

Hemorrhage. The number of deaths reported from this cause in 1878, was 14; 5 more than in 1877, and 0.70 per cent. of all from known causes; males, 5; females, 9; American parentage, 5; foreign, 9. So far as specified, 6 were from hemorrhage from the lungs; 2, hemorrhage from the stomach; 2, umbilical; 1, uterine; 1 from the bowels, and 1 from hemorrhoids.

Hernia. This was stated as the cause of 5 deaths, in 1878; 2 males, and 3 females; 3 of American, and 2 of foreign parentage; total, 5 or 0.25 per cent. of all from known causes.

Hooping Cough. The number of deaths from this cause in 1878, was 43; this is a larger number than has ever before been recorded in these reports. The decedents were as follows in regard to age: 17 under one year; 15, between 1 and 2 years; 9 between 2 and 5 years; 2, between 5 and 10 years. There were deaths from this cause in each month of the year, except May, but the largest number occurred in the third quarter, viz.: 28, or 53.49 per cent. of all; the deaths in each month were as follows:—

January 2	August11
February 4	September 6
March 3	October 4
April 2	November 1
May	December 3
June 1	The second secon
July 6	Total43

Males, 20; females, 23; total, 43, or 2.18 per cent. of all.
American parentage, 24; foreign parentage, 19.

In more than three-fourths of the cases, the hooping cough was accompanied or followed by some other disease, which, as being only induced by the cough, or fatal because of weakness consequent upon the cough, was not accounted as the primary or real cause of death.

Thus in 14 cases it was accompanied by convulsions; in 12, by pneumonia; in 2, by congestion of the lungs, and in one case each, by atelectasis, bronchitis, pleurisy, diarrhœa and enteritis; in 10 cases hooping cough alone was returned as the cause of death.

In one case it was accompanied by measles, and the death assigned to that cause; and in one case by marasmus, which was accounted the cause of death, because it is a lingering disease by which a child might gradually waste away, after the cough, as the primary cause, was entirely removed.

The statistics for the twenty-three years, 1856-1878, are as follows:-

Whole number, 371; annual average, 16.

Males, 159; females, 212; total, 371, or 1.24 per cent. of all.

American parentage, 153; foreign parentage, 218.

Hydrocephalus. There were 39 deaths from hydrocephalus in 1878; this was 18 more than in the preceding year.

Males, 17; females, 22; total, 39, or 1.98 per cent. of all. American parentage, 20; foreign parentage, 19.

For the twenty-three years, 1856-1878, the statistics are :-

Whole number, 698; annual average, 30.

Males, 379; females, 319; total, 698, or 2.33 per cent. of all.

American parentage, 353; foreign parentage, 345.

Insanity. The number of deaths reported from this cause in 1878, was 17; the largest number that has ever been recorded. Males, 3; females, 14; American parentage, 11; foreign parentage, 6; total, 17, or 0.86 per cent. of all from known causes.

The statistics for the twenty-three years, 1856-1878, are:-

Whole number, 236; annual average, 10.

Males, 112; females, 124; total, 236, or 0.79 per cent. of all.

American parentage, 170; foreign parentage, 66.

As has been frequently stated in these reports, so large a number of deaths from insanity is owing to the fact that Butler Hospital for the insane is within the limits of the city; and a large majority of the deaths occur there, many of the decedents not being residents of the city. It is owing to this also, that so large a proportion of the decedents are of American parentage; that class being more largely represented there than the foreign.

Intemperance. There were 9 deaths from this cause in 1878; 2 of which were from delirium tremens. Number of males, 6; females, 3; American parentage, 3; foreign, 6; total, 9, or 0.46 per cent. of all from known causes.

The following statistics are for the twenty-two years, 1857–1878:

Whole number from known causes, 28,921.

From intemperance, 222; annual average, 10.

Males, 161; females, 61; total, 222, or 0.77 per cent. of all.

American parentage, 64; foreign parentage, 158.

Of the 222 decedents, 72.52 per cent. are males; and 71.17 per cent. are of foreign parentage.

Kidneys, Diseases of. The number of deaths in 1878 from diseases of the kidneys, was 47; of which 33 were from Bright's disease.

Males, 30; females, 17; total, 47, or 2.38 per cent. of all. American parentage, 23; foreign parentage, 24.

The statistics for twenty-three years, 1856-1878:—

Whole number, 487; annual average, 19.

Males, 260; females, 177; total, 437, or 1.46 per cent. of all.

Liver, Diseases of. The deaths from diseases of the liver numbered 18, or 0.91 per cent. of all from known causes; males, 9; females, 9; American parentage, 10; foreign, 8.

For the twenty-three years, 1856-1878, the numbers are as follows:—

Whole number, 337; annual average, 15.

Males, 168; females, 169; total, 337, or 1.13 per cent. of all.

American parentage, 166; foreign parentage, 171.

Malformations. Twenty-two deaths were reported from this cause in 1878; 6 more than in 1877, and 1.12 per cent. of all from known causes; number of males, 15; of females, 7; of American parentage, 10; of foreign, 12.

They were specified as follows: in ten cases, malformation of the heart; in 6, spina bifida; in 3, of the anus; and one each of the palate, lungs, and rectum.

The following are the statistics for the twenty-three years, 1856-1878:—

Males, 110; females, 64; total, 174, or 0.58 per cent. of all.
American parentage, 114; foreign parentage, 60.

Marasmus. This was returned as the cause of 41 deaths in 1878; 14 more than in 1877, and 2.08 per cent. of all from known causes; males, 25; females, 16; American parentage, 14; foreign, 27.

The statistics are as follows for the twenty-three years, 1856-1878:—

Whole number, 590: annual average, 26.

Males, 816; females, 274; total, 590. or 1.97 per cent of all.

American parentage, 296; foreign parentage, 294.

Measles. This disease has appeared in Providence as an epidemic at very irregular intervals, and in 1878 was the cause of 50 deaths, a larger number than ever before recorded. The following table shows the number in each year since 1856:—

1856 1	1862 1	1868 4	1874 4
1867 2	186316	1869 6	1875 0
185840	1864 7	187016	1876 0
1859 0	1865 7	1871 1	1877 2
18600	1866 9	1872 7	1878 50
1861 6	1867 0	187328	
			Total, 23 years207

The decedents in 1878 were divided as to age, as follows: 11 under 1 year; 14, between 1 and 2 years; 15, between 2 and 5 years; 9 between 5 and 10; and 1 between 10 and 15.

Forty-five of the deaths, or 90 per cent. of all, were in the first quarter of the year; 11, in January; 22, in February; and 12 in March; there were 4 deaths in April, and 1 in June; none in the last six months of the year.

It may be said of this as it has already been said of hooping-cough, that in a large majority of cases it is attended or followed by some other disease, which yet could not be considered as really the cause of death. In only 8 cases was measles alone returned as the cause; in 15 cases, it was accompanied by pneumonia; in 8, by croup; in 7, by bronchitis, (which, in 2 instances was specified as "capillary"); in 7, by convulsions; in 3, by meningitis; and in 1 each, by hooping-cough, and uremia; there was one case in which it was followed by marasmus, and the death was assigned to that cause.

The 50 deaths were 2.53 per cent. of all from known causes. Males, 25; females, 25; American parentage, 16; foreign, 34.

The decedents were 68 per cent. of foreign parentage, and 32, of American; this has generally been the rule, as may be seen from the following statistics for the twenty-three years, 1856-1878:

Males, 100; females, 107; total, 207, or 0.69 per cent. of all.

American parentage, 60; foreign parentage, 147.

In the long period of time the decedents are 71.01 per cent. of foreign, and 28.99 of American parentage.

Meningitis, Cerebrò-Spinal. The number of deaths from this cause in 1878 was 7, the same as in 1877.

Males, 4; females, 3; total, 7, or 0.36 per cent. of all. American parentage, 3; foreign parentage, 4.

Since 1864, when the first deaths were reported in Providence, from this cause, the number in each year has been as follows:—

1864 2	1870 0	1876 4
1865 2	187112	1877 7
1866 1	187210	1878 7
1867 0	187839	
1868 1	187410	Total, 15 years101
1869 0	1875 6	

Whole number from known causes in 15 years, 22,112; from cerebrospinal meningitis, 101.

Males, 55; females, 46; total, 101, or 0.46 per cent. of all.

American parentage, 48; foreign parentage, 53.

The number and percentage of deaths in each division of age, and according to sex and parentage, were as follows, for the 15 years:—

AGES.	Under 1 Year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	Males.	Females.	American.	Foreign.	Total.
No.	10	9	10	22	12	13	13	6	2	1	1	99	46	48	63	101
PER CENT.	4.05	4.95	00.0	24.76	11.88	12.87	12.87	8.91	6.93	0.09	66.0	54.46	45.54	47.52	62.48	100.00

Of these 101 decedents, 57, or 56.44 per cent. were less than fifteen years old.

Murder and Homicide. Only one death was returned under this head in 1878, one American male, under one year old, was killed by arsenic; 0.05 per cent. of all from known causes.

Old Age. This was reported as the cause of 61 deaths in 1878. Of these decedents, 14 were over 90 years of age; 6 more than in 1877; of these 14, 5 were males and 9 females, and two of the latter were reported as 105 and 107 years old respectively. Their names were Henrietta Brown, colored, born in Maryland, died September 22, 1878, aged 107 years, 5 months, 7 days; and Mary Dunn, white, born in Ireland, died December 4, 1878, aged 105 years.

Males, 18; females, 43; total, 61, or 3.09 per cent. of all. American parentage, 38; foreign parentage, 23,

The following are the statistics for the twenty-three years, 1856-1878:—

Whole number, 1,190; annual average, 52.

Males, 389; females, 801; total, 1,190, or 3.97 per cent. of all.

American parentage, 735; foreign parentage, 455.

Of the 1,190 decedents, 67.31 per cent. are females; and 61.76 per cent. are of American parentage.

Only a small proportion of those who live to an advanced age are reported under this head, as may be seen from the following numbers:—

During the year 1878, there were 209 decedents of the age of 70 years or over; 129 of American, and 80 of foreign parentage; this was 10.59 per cent. of all from known causes.

In the twenty-three years, 1856-1878, the number of decedents of 70 years old and over, was 3,231, or 10.79 per cent. of all from known causes; 2,276 of American, and 955 of foreign parentage.

Paralysis. Though not in reality a distinct disease, but a symptom, of which the primary cause should be specified more frequently than it is, yet paralysis alone is often reported as the cause of death. It was reported in 1878 as the cause of 24 deaths, the same number as in each of the two years preceding.

Males, 10; females, 14; total, 24, or 1.22 per cent. of all.

American parentage, 18; foreign parentage, 6.

The statistics for the twenty-three years, 1856-1878, are as follows:

Whole number, 878; annual average, 16.

Males, 193; females, 185; total, 878; or 12.62 per cent. of all.

American parentage, 286; foreign parentage, 92.

These deaths being of a similar nature with those from apoplexy, the statistics for the two causes are here given for each of the twenty-three years, 1856-1878:—

Years.	Apoplexy.	Paralysis.	Total.	Years.	Apoplexy.	Paralysis.	Total
1856		2	23	1868	22	24	46
1857		9	27	1869	32	16	48
1858	15	3	18	1870	28	24	52
1850	16	6	22	1871	33	28	61
1860	17	11	28	1872	31	20	51
1861	14	11	25	1873	36	21	57
1862	19		24	1874	36	23	59
1863		8	28	1875	41	35	76
1864	13	16	29	1876	44	24	68
1865	20	18	38	1877	50	24	74
1866	21	8	29	1878	42	24	66
1867	31	18	49				
			T	otals, 23 y	'rs620	378	998

The statistics of both causes, as to sex and parentage, for the twenty-three years, are:—

Whole number, 998; annual average, 43.

Males, 494; females, 504; total, 998, or 3.33 per cent. of all.

American parentage, 732; foreign parentage, 266.

The decedents are very equally divided as to sex. but very unequally as to parentage; 73.35 per cent. being of American, and only 26.65 per cent. of foreign parentage.

Peritonitis. The deaths from this cause in 1878, numbered 17; 7 more than in 1877, and 0.86 per cent. of all from known causes; males, 10; females, 7; American parentage, 10; foreign, 7.

The statistics for the twenty-three years, 1856-1878, are :-

Whole number, 201; annual average, 9.

Males, 90; females, 111; total, 201, or 0.67 per cent. of all.

American parentage, 114; foreign parentage, 87.

Pneumonia. The number of deaths from pneumonia, in 1878, was 118; from congestion of the lungs, 22; total, 140. This is a great increase over the previous year, when the number was only 98; but it is less than in 1876 or 1875. The deaths in 1878 were distributed through the year as follows:—

1st quarter	3d quarter17
2d quarter 30	4th quarter 37

The decedents are mostly very young, or quite advanced in years, and very few in middle life; thus 62, or 44.28 per cent. were under 10 years of age; and 50, or 35.71 per cent. were over 50 years.

Males, 63; females, 77; total, 140, or 7.10 per cent of all. American parentage, 70; foreign parentage, 70.

The following are the statistics for the twenty-three years, 1856-1878, inclusive:—

Whole number, 2,157; annual average, 94.

Males, 1,116; females, 1,041, total, 2,157, or 7.20 per cent. of all.

American parentage, 1,094; foreign parentage, 1,063.

Scarlatina. The number of deaths from this cause in 1878, was 57; more than in any year since 1875; the number in 1877 was only 12. The following table gives the details as to sex, parentage, season and locality for each of the 24 years, 1855-1878, and for the entire period:—

SCARLATINA IN PROVIDENCE, TWENTY-FOUR YEARS.

Total.	1,649	743	652 897	2552544 1458 1558 1558 1558 1558 1558 1558 1	243 243 243 263 203 203 199 146 146
1878.	67 2.89	ងន	412	9939-99-2009 9939-99-2009	#+# Murula
1877.	110	***	60.00	B 844 +04	00 H000
1876.	1.87	22	#2	+66 40 -17	4
1875.	4.91	28	23	235e5eu4uuva	2001-824592
1874.	268	55	123	822222222	2285402828
1873.	132	38	28	************	802080128
1872.	1.20	*1	81		4 H410
1871.	1.69	23.80	93	**************************************	
1870.	35	28	28	Brond wa a	H 410 V O V
1869.	138	35	22 88	8751×4+1574×	***************************************
1808.	6.60	នគ	22		₩ - a-a+a+a
1867.	1.00	101	1-00		***
1806.	0.29	01	21-1	- - -	- 6
1865.	108	88	66	お記記をおり4 311	825283
1864.	141	38	96	140048808188	#8884E
1863.	2.78	28	12		∞e∞≈≈≈
1862.	1.68	900	. I		
1801.	2.76	22	22	+++000	2000-000-
1800.	172	101	119	s - es	434 900
1820.	46	20	12.41	@@####### @	@-5w95w
1858.	52.23	28	88	**********	5257455 515227455
1867.	3.56	22	18	Ø-40404-01	B201-40-
1856.	144	28	22	記記器数字記字のおよりの	8782848
1855.	4.37	88	22		8-22523
SCARLATINA.	Whole number	Males	PARENTAGE: American Foreign	January January February Rebruary March April. June July August September October December	LOCALITY: Ward III IV VI VIII

The decedents in 1878, were divided as to age, as follows: under 1 year, 5; between 1 and 2 years, 7; between 2 and 5 years, 23; between 5 and 10 years, 18; between 10 and 15 years, 2; between 15 and 20 years, 1; and between 20 and 30 years, 1. There were 6 deaths in the first quarter of the year; 8, in the second; 18, in the third; and 25, in the fourth.

Males, 25; females, 32; total, 57, or 2.89 per cent. of all. American parentage, 17; foreign parentage, 40.

The statistics for the twenty-three years, 1856-1878, are:

Whole number, 1,507; annual average, 66.

Males, 721; females, 786; total, 1,507, or 5.03 per cent. of all. American parentage, 633; foreign parentage, 874.

Scrofula. There were 7 deaths from scrofula, in 1878; males, 3; females, 4; American parentage, 6; foreign, 1; total, 7, or 0.86 per cent. of all from known causes.

Small-Pox. There have been no deaths from small-pox in Providence, since 1875. For the twenty-three years, 1856-1878, the statistics are as follows:—

Males, 58; females, 21; total, 79, or 0.26 per cent. of all. American parentage, 48; foreign parentage, 81.

The particulars for each year are as follows:-

	Sex.		PARKN		Whole	_
	Males.	Females.	American.			Percentage
			5			
			• • • • • • • • • • • • • • • • • • • •			
			•••••			
			5			
			3			
			• • • • • • • • • • • • • • • • • • • •			
i.862	2	····· 2·····	1	3	4	0.45
			8. <i>.</i>			
l866	9	2	8. <i></i>	3	11	0.98
			• • • • • • • • • • • • • • • • • • • •			
1867	1	• • • • • • • • • • • • • • • • • • • •	1		1	0.11
1866	1	••••••		• • • • • • • • • • • • • • • • • • • •	1	0.09
1869	3	• • • • • • • • • • • • • • • • • • • •	. 3		3	0 24
1870		2		2	2	0.16
1871		8		2	2	0.24
1872	9	4	9	4	18	0.82
1873	12	2	8	6	14	0.82
1874						
1875	1	••••••		1	1	0.05
1876	•••••	• • • • • • • • • • • • • • • • • • • •		•••••	• • • • • • • • • • • • •	
1877			• • • • • • • • • • • • • • • • • • • •			
1878		• • • • • • • • • • • • • • • • • • • •				
	-	-	-	=	=	-
is years' total.	95		48	81	79	0.25

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Spine, Disease of. The number of deaths from this cause in 1878 was 4, or 0.20 per cent. of all from known causes; 2 males, and 2 females; 1 of American, and 3 of foreign parentage.

Spleen, Disease of. There was one death from this cause in 1878; a male, of foreign parentage.

Stomach, Diseases of. The number of deaths from diseases of the stomach in 1878, was 14; 6 less than in 1877, and 0.71 per cent. of all from known causes; males, 5; females, 9; American parentage, 3; foreign parentage, 11.

Suicide. There were 7 cases of suicide in 1878; 5 less than in 1877, and 0.36 per cent. of all from known causes. Males, 4; females, 3; American parentage, 2; foreign, 5. In three cases it was by laudanum; in two by pistol shot; in one by drowning; and in one by cutting the throat.

The statistics for the twenty-three years, 1856-1878, are:-

Males, 98; females, 41; total, 139, or 0.46 per cent. of all. American parentage, 79; foreign parentage, 60.

Teething. The number of deaths reported from this cause in 1878, was 7; the same number as in each of the two years preceding, and 0.36 per cent. of all from known causes. Six of the decedents were males, and 1 female; 4 of American, and 3 of foreign parentage.

Tuberculosis. The number of deaths ascribed to this cause in 1878, was 17; three less than in 1877, and 0.86 per cent. of all from known causes; males, 9; females, 8; of American parentage, 9; of foreign, 8.

There are other tubercular diseases, such as consumption, hydrocephalus, and others, which if united under one general head, would show a much larger percentage of the whole number of deaths, than any other cause.

Tumors. The number of deaths from tumors in 1878, was 6; 2 more than in 1877, and 0.30 per cent. of all from known causes.

Males, 1; females, 5; American parentage, 1; foreign, 5. Three were ovarian; one, abdominal; and one of the neck. There were also others that were classed under the head of disease of the organ affected.

Uterus, Disease of, was the cause of 5 deaths in 1878; 0.25 per cent. of all from known causes; all the decedents were of foreign parentage.

Unknown causes. The number of deaths in 1878, of which the cause was not stated, was 16, or 0.80 per cent. of all deaths. This was 10 more than in 1877; the decedents were 8 males, and 8 females; 4 of American, and 12 of foreign parentage. Twelve were children under one year old.

The statistics for the twenty-three years, 1856-1878, are:-

Whole number, 524; annual average, 23.

Males, 273; females, 251; total, 524, or 1.72 per cent. of all.

American parentage, 170; foreign parentage, 354.

Of the 524 decedents, 67.56 per cent. were of foreign parentage.

STILL-BORN.

The statistics in regard to this class for 1878, are as follows:-

Number. The number of still-born children in Providence, in 1878, was 139; 6 less than in 1877, and 5.29 per cent. of the 2,627 children born during the year, or one in 18.90.

Sex. The number of males, was 81; of females, 58; 5 less males, and 1 less female than in 1877.

Color. White, 132; colored, 7.

Locality. First ward, 22; second, 6; third, 14; fourth, 10; fifth, 10; sixth, 9; seventh, 14; eighth, 18; ninth, 17; tenth, 19.

Season. January, 12; February, 13; March, 19; April, 15; May, 6; June, 6; July, 10; August, 17; September, 12; October, 8; November, 9; December, 12.

The number and percentage in each quarter of the year were as follows:—

	Number.	Per cent.	Numbe	r. Per cent.
lst quarter	44	81.66	8d quarter89.	28.06
2d quarter	27	19.42	4th quarter29.	20.86

Parentage. In the following table, the first letter indicates the birth-place of the father; the second, that of the mother:—

A. and A	is It. and It	UL and E
I. and I	0 E. and A 5	S. and E 1
		A. and B. A 3
		F. and B. A 1
		1 A. and D 1
		2 F. and I 1
I. and A	1 A. and E	1 —
	1	Total139

A.—American; I.—Irish; E.—English; S.—Scotch; G.—German; Sw.—Swedish; It.—Italian; B. A.—British American; F. French; D.—Dutch.

There were 2,627 children born in Providence in 1878, of whom 139, or 5.29 per cent. were still-born. Of these 2,627 children, 1,273, or 48.46 per cent. had American mothers; 882, or 33.57 per cent. had Irish mothers; and 472, or 17.97 per cent. had mothers who were natives of other foreign countries. Of the 139 still-born children, 61, or 43.88 per cent. had American mothers; 64, or 46.04 per cent. had Irish mothers; and 14, or 10.07 per cent. had mothers who were natives of other foreign countries.

Of the 1,273 children, with American mothers, 61, or 4.79 per cent. were still-born; of the 882 with Irish mothers, 64, or 7.26 per cent. were still-born; of the 472 with mothers of other nationalities, 14, or 2.97 per cent. The proportions are as follows: to American mothers, one still-born in 20.87; to Irish mothers, one in 13.78; to other foreign mothers, one in 33.71.

These numbers may be compared with those of 1877 as follows: the number and proportion of children born to American mothers were less than in the previous year; the number of still-born, and the proportion of these to the whole number, were also less.

Of children of Irish mothers, the number and proportion of all children born were larger in 1878 than in 1877; also the number and proportion of still-born children. Of the children of other foreign mothers, the number and percentage were larger in 1878 than in the preceding year; of still-born the number and percentage very much less. These comparisons are made with the figures for 1877, as corrected, there having been a slight error in the report for that year.

For the twenty-three years, 1856-1878, the statistics are as follows:-

Number of children born, 43,900.

Number of still-born, 2,603, or 5.93 per cent. of all, or one in 16.87. Of these 2,603 still-born children, 2,443 were white, and 160 were colored; males, 1,501, or 57.66 per cent.; females, 1,102, or 42.34 per cent. Reckoning the parentage according to the fathers' birth-place, 1,090 were American; 1,181, Irish; and 325 other foreign; 7 were of unknown parentage.

GENERAL SUMMARY.

The following recapitulatory table will give the general results of registration in Providence during the 24 years, 1855 to 1878, inclusive:—

TABLE XXI. BIRTHS, MARRIAGES, AND DEATHS

YEAR.	Population.	Births.	To population, one birth in	Marriages.	Of population, one person married in	Deaths.	Of population, one death in
	45.505				07.50	991	48.22
1×65	47,785	1,600	29,86	636 656	37.56 37.14	1,065	45.75
1857	48,727	1,675 1,688	29.09 29.30	614	40.27	925	53.46
1858	49,457	1,724	28.68	560	44.15	1,017	48.63
1559	50,100	1,593	31.45	616	40.66	899	55.72
1860	50,666	1,648	30.74	633	40.02	1,001	50.61
1961	50,666	1,725	29.37	547	40.31	1,051	48.21
1962	52,238	1.529	34.16	547	47.75	914	57.15
1463	53,024	1,307	40.56	610	43.46	1,214	43.67
1964	53,810	1,344	40.04	739	36.40	1,281	42.01
1866	54,595	1,298	42.06	709	38.50	1,211	45.08
1:66	55,709	1,632	34.13	812	34.30	1,036	53.77
1867	56,824	1,624	34.90	855	33.23	960	59.19
Þ86	64,138	1,866	34.37	872	36.77	1,110	57.78
1:69	66,522	1,819	36.57	930	35.76	1,256	52.96
1879	68,904	1,889	36.48	959	35.93	1,263	54.55
1871	72,011	1,960	36.74	943	38.13	1,254	57.42
1872	75,118	2,206	34.05	1,084	34.63	1,603	46.86
1873	78,225	2,128	36.76	1,150	34.01	1,719	45.51
1874	96,452	2,780	34.69	1,176	41.01	1,987	48.54
1×75	100,675	2,064	37.79	1,079	46.65	1,915	52.57
1×76	101,500	2,631	38.58	949	53.48	1,865	54.42
1577	100,675	2,543	39.58	953	52.82	1,938	51.95 50.78
1678	101,000	2,627	38.45	1,016	49.70	1,989	50.78
Average	66,595	1,896	35.12	819	40.66	1,311	50.80

BIRTHS AND DEATHS BY PARENTAGE.

The following recapitulatory table shows the number of births and deaths, by parentage, in Providence, in each of the twenty-three years, from 1856-1878 inclusive. In the three years, 1863, 1864, and 1865, there was an excess of deaths over the births, in the population of American parentage:—

TABLE XXII.

	AMERIC	AN PARI	INTAGE.	FOREIGN PARENTAGE.				
YEAR.	Deaths.	Births.	Excess of Births.	Deaths.	Births.	Excess of Births		
1866	541	759	218	524	916	392		
1867	523	701	178	402	987	585		
1858	512	771	259	505	953	448		
1869	479	659	180	420	934	514		
1860	- 512	731	219	489	917	428		
1861	538	740	202	513	985	472		
1862	460	663	203	454	866	412		
1863	597	528	-69	617	779	162		
864	649	579	-70	632	765	133		
865	631	546	-85	580	752	172		
866	513	726	213	523	906	383		
867	528	725	197	432	899	467		
868	596	882	286	514	984	470		
869	627	926	200	629	893	264		
1870	642	890	248	621	999	378		
871	643	980	337	611	980	360		
872	832	1,035	203	771	1,171	400		
1873	862	1,003	141	857	1,125	268		
1874	885	1,217	332	1,102	1,563	461		
1875	914	1,180	266	1,001	1,484	483		
876	875	1,169	294	990	1,462	472		
1877	908	1,238	330	1,030	1,305	275		
1878	889	1,215	326	1,100	1,412	312		
23 years	15,156	19,863	4,707	15,317	24,037	8,720		

CITY ORDINANCES.—CHAPTER XLVII.

REGISTRATION OF DEATHS.

- SECTION 1. There shall be appointed by the City Council a sufficient number of persons to act as undertakers, removable at the pleasure of the City Council.
- SEC. 2. Whenever any person shall die in the city, the physician attending in his or her last sickness shall furnish to the undertaker attending the funeral, or to the city registrar, a certificate, giving the name of the person, date of death, and the disease or cause of his or her death.
- SEC. 3. No person shall bury, or place in the tomb. or remove from the city, or otherwise dispose of the body of any human being who shall die in the city, without first reporting the death to the city registrar, and obtaining a permit from him.
- Sec. 4. No permit shall be given as provided in section three, until the city registrar is furnished with the information in relation to the deceased person, required by the laws of the State, for record, so far as the same can be ascertained, together with the physician's certificate of the cause of death, whenever a physician has been in attendance, or a coroner's certificate, whenever a coroner's inquest has been held.

Whenever a permit for burial is applied for, in a case of death without the attendance of a physician, or if it is impossible to obtain the physician's certificate, the city registrar shall investigate the case so far as may be necessary, and when he has obtained satisfactory evidence in relation to the cause and circumstances of the death, he shall sign the certificate and give the required permit. If not satisfied in relation to the cause and circumstances of the death, or if, in his opinion, the public good requires it, he shall report the case to a coroner for investigation.

- Src. 5. Whenever the body of a human being who has died out of the city shall be brought here for burial, the undertaker, or other person attending the funeral, shall furnish the report required in sections three and four, with the exception of the physician's certificate.
- SEC. 6. All funerals shall take place between sunrise and sunset, unless otherwise permitted or directed by the board of health. The top of every coffin deposited in the ground shall be at least three feet below the usual surface thereof.
- SEC. 7. The city registrar shall cause an abstract of the returns of deaths made to him to be published in all the newspapers which contract to do the city printing, monthly, and oftener if required by the board of health. He shall also, on or before the first day of April, annually, prepare and present to the city council a statement of the number of births, marriages and deaths which occurred in the city during the year ending with the thirty first day of December next preceding, with such other information and suggestions in relation thereto as he may deem useful for the promotion of the public health, and other interests of the city.
- SEC. 8. No undertaker or other person shall bury or cause to be buried the body of any deceased person in the city, except in such grounds as are or may be designated as burying-grounds, and authorized to be used as such.
- SEC. 9. Every person violating any provision of this chapter shall pay a fine of not less than five and not more than twenty dollars for each offence.



	•	
		•
•		
•		

Vital Statistics, Providence, R. I.

1878.

CITY REGISTRAR'S

TWENTY-FOURTH ANNUAL REPORT.

CITY DOCUMENT. [No. 18,

TWENTY-FIFTH ANNUAL REPORT

BIRTHS, MARRIAGES, AND DEATHS

CITY OF PROVIDENCE:

BY EDWIN M. SNOW, M. D.,

CONTRACTOR OF BUILDING STORY OF BUILDING





1880.]

CITY DOCUMENT.

[No. 18.

TWENTY-FIFTH ANNUAL REPORT

UPON THE

BIRTHS, MARRIAGES, AND DEATHS

IN THE

CITY OF PROVIDENCE:

FOR THE YEAR

1879,

By EDWIN M. SNOW, M. D..

SUPERINTENDENT OF HEALTH AND CITY REGISTRAR.



PROVIDENCE:

PROVIDENCE PRESS CO., PRINTERS TO THE CITY.

1880.

THE CITY OF PROVIDENCE.

JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved June 2, 1880.]

RESOLVED, That the twenty-fifth annual report of the City Registrar, for the year ending December 31, 1879, be received, and that six hundred copies thereof, and of the communication accompanying the same, be printed in pamphlet form for the use of the City Council, under direction of the joint committee on printing.

A true copy: witness,

HENRY V. A. JOSLIN, CITY CLERK.

INTRODUCTION.

CITY REGISTRAR'S OFFICE, PROVIDENCE, May 20, 1880.

To the Honorable the City Council:

Herewith I send my Twenty-Fifth Annual Report upon the Births, Marriages, and Deaths in the city of Providence. The statistics are for the year ending December 81, 1879. During that year 2,522 children were born in Providence, 2,142 persons were married, and 2,026 persons died in the city. Compared with the year 1878, the number of births was 105 less; of persons married, 110 more; and of deaths, 37 more.

Estimating the population of Providence, in the middle of the year 1879, at 103,000, we have the ratio of births, marriages, and deaths for the year, to population, as follows:—

One birth in 40 84, or 24.49 in each 1,000. One person married in 48.08, or 20.80 in each 1,000. One person died in 50.84, or 19.67 in each 1,000.

The twenty-five annual reports that I have presented as City Registrar, give the statistics of 48,022 children born, of 41,432 persons married, and of 33,490 persons who have died in the city of Providence. This is a total of 122,944 names that are contained in the records, with very full and minute particulars relating to each.

EDWIN M. SNOW, M. D.,

Superintendent of Health and City Registrar.

CONTENTS.

Departs	PAGE.
Wards, months, sex and color, (Table I.).	1-17
Births in connection with season, sex and locality, (Table II.),	2-3
Births in connection with season, sex and locality, (Table II.) Proportion of the sexes born at different seasons	3
Number and proportion of the sexes, 26 years	4
Color:—Births, marriages and deaths in 24 years	5
Parentage of children born in each ward, (Table III.)	6.
Parentage - Mixture of different nations (Table IV)	7-8
Parentage:—Mixture of different nations, (Table IV.) Number of 1st, 2d, etc., children in 1879, and 1855-1878, (Table V.)	10
Average number of children to each mother, 25 years, (Table VI.)	11
Number of the child, with age of the mother, (Table VII.)	12-14
Average number of children to each mother, 25 years, (Table VI.) Number of the child, with age of the mother, (Table VII.). Plurality-births, statistics for 1879. Plurality-births, 25 years' summary.	14-15
Plurality-births, 25 years' summary	15-17
MARRIAGES	18-27
Marriages at different seasons, 25 years, (Table VIII.)	18-19
Marriages at different seasons, 25 years, (Table VIII.)	20
Bride and groom born in same country, &c Proportions of American, foreign and mixed marriages, 25 years	20-21
Proportions of American, foreign and mixed marriages, 25 years	22
Marriages of colored persons. Age of persons married in 1879, (Table X.)	100
Average age of persons married, 24 years, (Table)	23
Average age of persons married, 24 years. (Table)	25
Nativity and residence	26
Marriage and education; certificate signed with a mark	26-27
DEATHS	28-85
Deaths to population in 1879; proportion in each class	28
Deaths by sex, condition, color, nativity, parentage and locality,	
(Table XII.)	29
Deaths by season; numbers and proportions; recapitulation	30-31
· Sex, condition, color, nativity, parentage and locality of decedents.	no no
Aggregate and average age by parentage, 1879, (Table XIII.)	32-33
Aggregate and average age by parentage, 24 years, (Tables XIV.	13.8
Aggregate and average age by parentage, 24 years, (Tables XIV. and XV.)	35-38
Difference between average age of American and foreign decedents.	
(Table)	37
Percentages at different ages by parentage, 1879, (Table XVI.) Percentages under five years of age by parentage, 24 years, (Table)	39
Percentages at different ages by parentage, 24 years, (Table)	40
Deaths by age and parentage in each month and quarter, 1879, (Table	71
XVIII)	42-44
Causes of death, sex, age. parentage and percentages, (Table XIX.)	45-50
Causes of death in connection with season, (Table XX.)	51-52
NOTES ON THE CAUSES OF DEATH IN 1879	53-82
Cholera Infantum; decedents by sex and parentage. 24 years (Table)	56-57
Consumption; sex and parentage; number of deaths at each age	57-59
Diarrhœa and dysentery; recapitulation for 40 years	60-63
Diphtheria; statistics for 1879	63
Diphtheria in Providence, 22 years, (Table)	64
cronp (Tables)	65-66
Diphtheria, epidemic and not epidemic compared, (Table)	67-69
Heart, diseases of, decedents by sex and parentage, 24 years (Table)	70-71
Meningitis, cerebro-spinal; deaths at each age, 16 years Scarlatina; statistics for 1879. Scarlatina in Providence, 25 years. (Table).	75
Scarlatina; Statistics for 1878.	78
Small-pox; recapitulatory table, 24 years	78 70 80
Unknown causes	82
Still-born children in 1879	83
Still-born children, 1856-1879	84-85
GENERAL SUMMARY	
Births, Marriages and Deaths; numbers and proportions, 25 years,	
(Table XXL)	80
Births and Deaths by parentage; excess of births, 24 years, (Table	-
XXII.)	87
Conv. Oppression. Per opped no Protognation	200

CITY REGISTRAR'S REPORT.

1879.

BIRTHS.

There were 2,522 children born in the city of Providence during the year 1879.

Table I. gives the number of children born in each ward of the city, in each month and quarter of the year; also the number of each sex, and the number of colored children born in the whole city, in each month and quarter:—

TABLE I. BIRTHS, 1879. Wards, Months, Sex, and Color.

PROVIDENCE. 1879.					WA	RDS					W	iole C	ity.	chil	olore dren ed in cedi	in-
	1.	11.	111.	ıv.	v.	VI.	vII.	VIII	ıx.	x.	Males.	Females.	Total.	м.	r.	Total.
January February March	33 25 18	8	32	10 7 9	21 18 12	3 8 10	15 17 16	24 21 22	37 28 29	36 38 47	117 95 97	80 107 91	197 202 188	6 1 5	6 3	7
1st Quarter	76	20	63	26	51	21	48	67	94	121	309	278	587	12	15	27
April May June	31 33 30		12	11 8 14	16 14 23	7 10 11	20 22 10	18 23 31	33 28 37	39 49 52	111 93 127	93 112 112	205		5 4 7	13 6 16
2d Quarter	94	23	. 55	33	53	28	52	72	98	140	331	317	648	19	16	35
July August September	35 28 20	5 9 4	15 23 21	10 3 13	13 28 20	11 7 8	14 20 11	24 18 35	38 41 41	47 38 47	108 100 109	115			1 1	934
3d Quarter	83	18	59	26	61	26	45	77	123	132	317	333	650	10	-6	16
October November December	27 17 21	7 9 4	20 11 22	8 9	30 15 23	13 5 10	24 16 16	29 16 36	35 24 36	54 52 46	124 101 115	66	167	5 2 10	3	9 4 13
4th Quarter	65	13	53	26	68	28	56	81	95	152	340	297	637	17	9	26
Whole Year	318	74	230	111	233	103	201	297	410	545	1.297	1,225	2,522	58	46	104

The number of births in 1879 was the smallest since 1873; it was 105 less than in the year 1878; the number of males, 61 less; of females, 44 less. In the second quarter of the year the number of births was 28 more than in the corresponding season of the year previous; in all the other quarters the number was much less. Only the second and fifth wards show any increase over 1878; the second ward having 18, and the fifth ward 21 more births in 1879; the other wards, collectively, had 144 less births than in the preceding year. The population of the city in the middle of the year 1879 is estimated at 103,000; the 2,522 births therefore give a ratio of one in 40.84 of the population, or 24.49 in each thousand.

Season. The largest monthly total of births in 1879 was 247 in October; the smallest, 167 in November. The largest quarterly total was 650 in the third quarter; the smallest, 587, in the first quarter.

The difference between the number of births in the first six months and the last six months of 1879 was 52, which was less than in several years previous; in 1878 the difference between the two parts of the year was 131; in 1877 it was 267. The number of births in the first six months of 1879 was 1,235, or 13 less than in the corresponding season of 1878; in the last six months of the year, the number was 1,287, or 92 less than in 1878.

The following figures give the numbers of births in each quarter of the year 1879, compared with the number for the corresponding seasons in the aggregate of twenty-three years, 1856-1878; also the numbers for the twenty-four years, 1856-1879, inclusive:—

	Jan. Iarch.		First. 6 mos.	July. Sept.	Oct. Dec.	Second 6 mos.	Whole Year.
Twenty-three years1	0,722	.19,906	21,628	.10,878	11,394	22,272	43,900
1879	587	. 648	1,235	. 650	637	1,287	2,522
Twenty-four years1	1,309	11,554	22,863	11,528	12,031	23,559	46,422

For the period of twenty-five years, 1855-1879, inclusive, the monthly totals of births in Providence were as follows: the daily average is also given for each month, and for the whole period:—

Births in	January.	February.	March.	April.	May.	June.
25 years, 1855-1879		8,683	4,201	3,718	. 8,978	4,226
Daily average	4.88	5.22	5.42	4.96	5.1 3	5.63
Births in	July.	. August.	Sept.	Oct.	Nov.	Dec.
25 years, 1855-1879	8,941	4,087	8,965	4,152	4,046	4,298
Daily average	5.09	5.91	5.29	5.36	5.89	5.55

The highest daily average for the whole period was in June; the lowest in January.

Sex and Locality. Table II. shows the number of children of each sex, born in each ward of the city, in each quarter of the year, and in the whole year 1879:—

TABLE II. BIRTHS, 1879. Wards, Sex, and Season.

	1st Quarter.			2d Quarter.			3d Quarter.			4th Quarter.			Whole Year.		
WARDS.	M.	F.	Total.	M.	F.	Total.	M.	F.	Total.	M.	F.	Total.	м.	F.	Total.
	36	40	76 20	49	45 12	94 23	40	43	83 18	30	35	65 13	155	163 38	318
III	33 14	30	63 26	11 27 15	28	55	33	26 13	59 26	25 13	28 13	53	118		230
v	29	22	51	24	18 29 13	53 28	18 29	32 13		43 14	25 14	68	125	108	233
VIII	24	24	48	15 26 34 49	26 38	52 72	13 22 40	23 37		30 47	26	56 81	102	99 136	201
ix x	56 57	38 64	94	49 81	49 59	98 140	57 59	66 73	123	52 81	43 71	95 152	214 278	196 267	410
Whole City.	309	278	587	331	317	648	317	333	650	340	297	637	1.297	1,225	2,525

Proportion of the Sexes. The whole number of children born in Providence in 1879 was 2,522, of whom 1,227 were males, and 1,225 females; this gives a proportion of 105.9 boys, to 100 girls; or 51.43 boys, and 48.57 girls in each 100 children born.

The number and proportion of each sex in the different quarters of the year 1879 were as follows:—

	Males.	Females.	Males	. Femal	es. Males.	Females.	
JanMarch	809 and	278, 01	111.1	to 100,	or 52.64 to	47.86 in eac	h 100 born.
April-June	331 and	817, or	104.4	to 100,	or 51.08 to	48.92 in eac	h 100 born.
July-Sept	817 and	833, or	95.2	to 100,	or 48.77 to	51.23 in eac	h 100 bo rn .
OctDec	840 and	297, or	114.5	to 100,	or 53.3 8 to	46 62 in eac	h 100 born.
_	—						
Whole Year	1,297 and	1,225, or	105.9	to 100, o	or 51. 48 to	48.57 in eac	h 100 born.

The same statistics for the period of twenty-four years, from 1856-1879 inclusive, are as follows: The whole number of children born during this period was 46,422; males, 23,850; females, 22,572.

```
Males. Females. Males. Females. Males. Females.

Jan -March...... 5,852 and 5,457, or 107.2 to 100, or 51.75 to 48.25 in each 100 born.

April-June...... 5,968 and 5,586, or 106.8 to 100, or 55 12 to 44.88 in each 100 born.

July-Sept...... 5,828 and 5,700, or 102.2 to 100, or 50.56 to 49.44 in each 100 born.

Oct.-Dec...... 6,202 and 5,829, or 106.4 to 100, or 51.55 to 48.45 in each 100 born.
```

Whole Period...23,850 and 22,572, or 105.2 to 100, or 51.38 to 48.62 in each 100 born.

In the twenty-six years, 1854 to 1879 inclusive, the number of children born in Providence was 49,391; males, 25,368; females, 24,023. The number and proportion of each sex in each year and in the whole period were as follows:

```
Males.
               Females. Males. Females. Males. Females.
1854..... 711 and
                658; or 108.0 to 100; or 51.91 and 48.09 in each 100 born.
1855 .... 807 and
               793; or 101.7 to 100; or 50.41 and 49.56 in each 100 born.
1857...... 872 and 816; or 106.8 to 100; or 51.66 and 48.34 in each 100 born.
1858..... 890 and
                834; or 106.7 to 100; or 51.62 and 48.38 in each 100 born.
1859 ..... 825 and
                768; or 107.4 to 100; or 51.79 and 48 21 in each 100 born.
1860...... 809 and 839; or 96.4 to 100; or 49.09 and 50.91 in each 100 born.
1862 ...... 791 and 738; or 107.2 to 100; or 51.73 and 48.27 in each 100 born.
1863...... 689 and 618; or 111.4 to 100; or 52.71 and 47.29 in each 100 boon.
1864..... 663 and
                681; or 97.3 to 100; or 49.33 and 50.67 in each 100 born.
                607; or 113.8 to 100; or 53.24 and 46.76 in each 100 born.
1865 ..... 691 and
1866...... 849 and 783; or 108.4 to 100; or 52.02 and 47.98 in each 100 born.
1867...... 830 and 794; or 104.5 to 100; or 51.11 and 48 89 in each 100 born.
1869...... 945 and 874; or 108.1 to 100; or 51.95 and 48.05 in each 100 born.
1870...... 967 and 922; or 104.8 to 100; or 51.19 and 48.81 in each 100 born.
1871...... 956 and 1,004; or 95.6 to 100; or 48.78 and 51.22 in each 100 born.
1872...... 1,079 and 1,127; or 95.7 to 100; or 48.91 and 51.09 in each 100 born.
26 years ...... 25,368 and 24,023; or 105.6 to 100; or 51.36 and 48.64 in each 100 born.
```

Color. The number of colored children born in Providence in 1879 was 104; the number having been the same now in three successive years; 58 were males, and 46 females; a proportion of 126.1 male to each 100 female children; or 55.77 males and 44.23 females in each 100 born. The births in the different wards were as follows: First ward, 13; second, 10; third, 8; fourth, 2; fifth, 1; sixth, 18; seventh, 10; eighth, 14; ninth, 27; tenth, 1. The following is the number in each month: January, 12; February, 7; March, 8; April, 13; May, 6; June, 16; July, 9; August, 3; September, 4; October, 9; November, 4; December, 13.

In the twenty-four years, 1856-1879 inclusive, there were 1,808 colored children born in Providence; 959 males, and 844 females; a proportion of 113.6 males to each 100 females; or 53.19 males, and 46.81 females in each 100 children born.

The statistics of births, marriages and deaths among the colored people of the city, in the 24 years, 1856–1879 inclusive, are as follows:

Twenty-three years, 1856-1878	1,699 births828 m	arriages1,787 deaths.
1879	104 births 43 m	arriages 112 deaths.
Twenty-four years, 1856-1879	1,808 births871 m	arriages1,899 deaths.

In 1879 there were 107.7 deaths to each 100 births among the colored people; in the twenty-four years, 1856-1879, 105.3 deaths to each 100 births.

PARENTAGE.

The statistics on the subject of parentage given in the following pages, show the variety of nations represented in our population, and the relative proportions of the different classes.

Table III. shows the number of children, according to parentage, born in each ward; also the number and proportion of each nationality in the whole city. In cases where the parents were of different foreign nationalities, the parentage is always given according to the birth-place of the father:—

TABLE III. BIRTHS, 1879. Wards and Parentage.

				-	WA	RDS						- 2
PARENTAGE.	1.	11.	ш.	ıv.	v.	VI.	VII.	viii	ıx.	x.	Whole City	In each 100 born there
American	114	55	69	61	58	74	87	153	230	128	1,029	40.80
lrish	96	6	82	24	76	13	60	53	82	190	682	27.0-
English, Scotch and Welsh	17	2	4	2	18	2	6	23	22	76	172	6.8
German	5	2	5		11	1	1		12	13	50	1.9
British American	13		3	4	13	1		12	5	13	64	2.5
Portuguese			18				4		2		24	.00
Other Foreign	14		3	5	7		3	2	6	3	43	1.7
Total Foreigu	145	10	115	35	125	17	74	90	129	295	1,035	41.0
American father and Foreign mother	30	3	14	5	25	1	13	23	24	61	199	7.5
Foreign father and American mother	29	6	32	10	25	11	27	31	27	61	259	10.2
Ward totals	318	74	230	111	923	103	901	997	410	545	2,522	100:00

The "other foreign" in this table were as follows: Italian, 12; Polish and Swedish, 8 each; Russian, 4; Austrian and French, 3 each; Danish, 2; St. Helena and Norwegian, 1 each; and one which was not given any more definitely than "European."

The proportion of American and foreign children born, differs very greatly in the different wards of the city. Thus in the Third, Fifth, and Tenth wards there were more children born of Irish than of American parents. The aggregate of foreign children born was greater than of American children in the First, Third, Fifth, and Tenth wards. The aggregate of children of American fathers in the whole city, amounts to 48.69 per cent., a little less than one-half the whole number.

The following table gives the number and percentage of children of American, foreign, and mixed parentage in each of the

Color. The number of colored children born in Providence in 1879 was 104; the number having been the same now in three successive years; 58 were males, and 46 females; a proportion of 126.1 male to each 100 female children; or 55.77 males and 44.23 females in each 100 born. The births in the different wards were as follows: First ward, 13; second, 10; third, 8; fourth, 2; fifth, 1; sixth, 18; seventh, 10; eighth, 14; ninth, 27; tenth, 1. The following is the number in each month: January, 12; February, 7; March, 8; April, 13; May, 6; June, 16; July, 9; August, 3; September, 4; October, 9; November, 4; December, 13.

In the twenty-four years, 1856-1879 inclusive, there were 1,808 colored children born in Providence; 959 males, and 844 females; a proportion of 11,8.6 males to each 100 females; or 53.19 males, and 46.81 females in each 100 children born.

The statistics of births, marriages and deaths among the colored people of the city, in the 24 years, 1856–1879 inclusive, are as follows:

In 1879 there were 107.7 deaths to each 100 births among the colored people; in the twenty-four years, 1856-1879, 105.8 deaths to each 100 births.

PARENTAGE.

The statistics on the subject of parentage given in the following pages, show the variety of nations represented in our population, and the relative proportions of the different classes.

Table III. shows the number of children, according to parentage, born in each ward; also the number and proportion of each nationality in the whole city. In cases where the parents were of different foreign nationalities, the parentage is always given according to the birth-place of the father:—

	American. Per cent.			German.		Other Portuguese. for'n. Per cent. Per ct.
1856	42.09	41.79	4.84	2.27		2.27
1857	38.51	44.32	5.92	3.73		1.54
1858	41.30	42.98	4.82	2.26		1.33
1859	37.60	45.58	5.34	3.26		1.51
1860	40.11	40.96	6.55	2.85		1.64
1861	37.91	42 61	6.03	2.49		1.97
1862	38.52	41.53	6:02	3.07		1.54
1863	35.73	43.08	6 35	2.98		1.99
1864	38.69	41.36	6.55	2.75		1.34
1865	36.67	42.99	5.93	1.85		1.93
1866	40.20	39.52	6.25	2 21		2.32
1867	39.72	37 81	5.66	3.02		2.91
1868	41.75	36.01	6 48	1.88	1.07	1.07
1869	45.19	34.14	4.01	1.81	1.32	1.38
1870	42 35	34.68,	6.19	1.96	1.91	
1871	44.85	32.25	4.54	1.63	2.09	1.73
1872	41.12	32.46	5.67	2.77	2.31	2.31
1873	40.84	30.50	7.19	1.88	2.49	2.16
1874	38.42	.32.63	8.13	2.34	2.19	1.221.19
1875	37.50	.31.53	7.69	2.85	2.67	981.43
1876	38.27	.31.05	7.75	2.59	2.89	1.221.25
1877	41.45	.28.08	6.96	1.96	2.56	1.101 30
1878	39.40	.29.65	6.74	2.55	2.86	1.181.71
1879	40.80	.27.04	6.82	1.98	2.54	961.70

The British American were included in the "other foreign" until the year 1868; and the Portuguese until 1874.

The lowest percentage of American parentage in any year was 35.73 in 1863; the highest, 45.19 in 1869. The lowest percentage of Irish parentage was 27.04 in 1879; the highest, 45.58 in 1859. The other classes vary but little from year to year.

The parentage of the children born in 1879 is shown more exactly in Table 1V., which gives the birth-places of both parents.

TABLE IV. BIRTHS, 1879. Parentage.

	вп	RTH-	PLAC	ES OF	мо	THER	s.	aecording birth-
BIRTH-PLACES OF FATHERS.	United States.	Ireland.	England, Scot- land and Wales.	Germany.	British America.	Portugal and Western Islands.		Whole number accorto to the father's birth place.
United States	1,029	107	55	8	32		2	1,228
Ireland	126	635	31		16			808
England, Scotland and Wales	69	25	135	1	11			241
Germany	24	2	4	39	2		3	74
British America	30	8	5		51			94
Portugal and Western Islands						24		24
Other Countries	10	1	3	2	2		35	53
Whole number according to the mother's birth-place	1,288	778	238	45	114	24	40	2,522

The number of children of purely American parentage in 1879 was 1,029, or 40.80 per cent. of all; in addition to these there were 199 with American fathers and foreign mothers; and 259 with American mothers and foreign fathers; a total of 1,228 with American fathers; and 1,288 with American mothers; in both classes a larger number than in 1878. The number of children whose parents were natives of the same foreign country was 919, or 36.44 per cent. of all; of children whose parents were natives of different foreign countries, 116, or 4.60 per cent.; of children whose parents were, one American and one foreign, 458, or 18.16 per cent. of the whole number.

NUMBER OF THE CHILD.

Table V. shows how many of the 2,522 children born in Providence in 1879 were the first, second, third, &c., children of their mothers; also the same facts for the 45,500 children born in the twenty-four years preceding, 1855-1878 inclusive:—

TABLE V. BIRTHS, 1879. Number of the Child.

NUMB	EF	2 0	F	CHILD.	1879.	24 years, 1855-1878.	NUMBER OF CHILD. 1879. 24 years 1855-1878
First ch	ild	of	the	mother	561	11,263	Twelfth child of the moth'r 11 183
Second	**	46	44		476	9,419	Thirteenth " " " 10 69
Third	**	**	**	**	413	7,276	Fourteenth " " " 5 30
Fourth	**	**	**		346	5,436	Fifteenth " " " 2 14
Fifth	44	"	**	"	241	3,945	Sixteenth " " " " 3
Sixth	**	**	**		169	2,875	Seventeenth" " " " "
Seventh	**	44	**	**	118	2,016	Eighteenth " " " " 11
Eighth	**	**	**	15	89	1,276	Nineteenth " " " 4
Ninth	**	**	**	44	45	840	
Tenth	*	**	"	**	17	555	Whole number of chil-
Eleventh	**	**	**	**	19	288	dren

The 561 children who were the first children of their mothers were 22.24 per cent. of the whole number of children born in 1879. Of the 45,500 children born in the twenty-four years previous, 1855 to 1878 inclusive, 11,263 or 24.75 per cent., were the first children their mothers had borne. The percentage of first children in 1879 was considerably larger than in the preceding year.

Average number of children to each mother. The number of twin births in 1879 was 23; the number of mothers, therefore, was 2,499. Table VII. shows that these mothers had borne, in all, 9,039 children, giving an average of 3.62 children to each mother.

The next table shows the average number of children to each mother, in each of the twenty-five years, 1855 to 1879 inclusive. This table shows the number of children, the mothers included in the figures had borne previous to the end of the year 1879; but it does not show the average number of children the same mothers may have in their life-time.

Table VI. Births, 1855 to 1879.

Average number of children to each mother.

Years.	No. of Mothers.	No. of Children.	Aggregate No. of children. Average.
1855	1,568		5,2143.28 children to each mother.
1856	1,658	1,675	5,5038.85 children to each mother.
		1,688	
1858	1,706	1,724	5,685 8.33 children to each mother
1859	1,570	1,598	5.6183.58 children to each mother.
1880	1,627	1,648	5,8413.59 children to each mother.
1861	1,694	1,725	6,0178.55 children to each mother.
18 62	1,512	1,529	5,322
18 63	1,296	1,807	4,7178.64 children to each mother
1864	1,897	1,344	4,8873.68 children to each mother.
1865	1,279	1,299	4,6258.60 children to each mother
1866	1,614	1,632	5,744
1867	1,595	1,094	5,497 3.50 children to each mother
1868	1,841	1,886	6,3383.44 children to each mother
1869	1,797	1,819	5,954 3.30 children to each mother.
1870	1,859	1,889	6,1458.81 children to each mother.
1871	1,937	1,980	6,420
1873	2,174	2,206	7,8463.38 children to each mother.
1878	2,104	2,128	6,7318.20 children to each mother
1874	2,788	2,780	8,9808.27 children to each mother.
1875	2,634	2,664	8,9773.41 children to each mother
1876	2,604	2,631	8.9148.42 children to each mother
1877	2,516	2,548	8,705 8.46 children to each mother.
			9,4178.63 children to each mother
1879	2,499	2,5 2 2	9,039 3.62 children to each mother.
25 years	47.480	48,093	163,411 3.45 children to each mother.

Table VII. gives the number of the child in connection with the age of the mother; and shows how many first, second, etc., children were born in 1879 to mothers at the respective ages given, and the aggregate and average number of children they had borne.

Table VII. Births, 1879. Number of child, with age of mother.

		-	-	-	-	-	-	-	-	-	-	-	-	-			1	1	
				N	UMB	ER (OF C	HIL	D.						ı	20	E E		100
AGE OF THE	-	1					1	1		7						hers	ore	hey	the
MOTHERS													13	П		not	by b	ne.	ng ,
IN YEARS.				Ш							۰			H	۲	of I	they 1879.	ldre	H CH
IN IEARS.				1						И	ı			ı	ı	ach	ren	d children	Sept and a sept a sept a sept a sept a sept a sept a sept a sept a sept a sept a sept a sept a sept
				÷	4	2	1	-	-	i.	11th.	12th.	13th,	14th.	loth.	Number of mothers at each age.	Children they bore 1879,	Total children they had borne.	Lverage number of children to mothers at each age.
	1st.	2d.	3d.	4th	5th.	6th	7th.	8th.	9th.	10th.	111	12	13	14	15	Z	5	To	4
14	1															1	1	1	1.00
15	1										.,					. 1	1	1	1.00
16	3															3			1.00
17	9										• •	••		• •	• •	9	1	- 3	1.00
18	19		7000	••••							• •	• •	••	•	••	24	24		1.21
19	82	100	3	••••						••	••	••	••	•	•	38	38	-	1.18
20	36	1000		****							•			ш	•	59 83		1770	1.46
21	54	100	20	*2	2		****	****				•	•	М	**	110	110	1000	2100
23	61 *57	37		6	4	***										123	124	233	1.56
24	57	1		*14	3		****									146	147	297	1.88
25	50	000		21	3	1	2		1							155	155		2.39
26	39	1 30	1	22	11	1 7	100	1		ı						155	155		2.70
27	*36	14.54	1000	34	20	1	100	1	1			.,				185	189		2.99
28	*27	*34	36	*44	25	8	2	1								174	177	574	3.24
29	21	21	24	29	15	16	2	3		1						132	132	470	3.56
30	19	39	47	45	23	18	12	3	1	••				!		207	207	758	3.68
31	9	9	20	12	10	8		3		••	••				• •	74	74	280	3.78
82	7	15	10	24	24	11	100	8	100	••	•	••	••		•••	140	140	610	3.64
33	6		15	19	*16			- 5	3		121	• •	• •	••	•	98	100	457	4.57
34	3	50	*12	14	12	†14	17	1.757			1131		•	••	1	100	04	565	5,43
35	5	- 23	100	*20	†25	16	1 6	*15	100	100	01 0	1			••	124	128	688	5.87
36	4	6	15	8	*17	*15	6	6	1	1	- 1		100	90		30	82 56	409	4.50
38	2	6	5	7 5	3	8	11 8	8	4	1 0	2 01	1 2	3		-1	56	57	344	6.14
39	2	4	6	4	4	5	*9	3			4	9		4		49	50	323	6.46
40	1	1	3	9	8	6	5	6	9	100		2			2	53	53	345	6.51
41		1		2	3		1	1	1			1		-		11	11	75	6.82
220		1		9	1	3	1	2			n		4	1	ю.	20	20	171	8,55
43			2	1		2					1	1		2		9	9	73	8.11
		1		1		1		1		1	1		1			7	7	54	7.71
45		1			2	2		2			1					8	8	51	6.37
46								1		• •		1				2	2	20	10.00
48						1	1			-						2	2	13	6.50
50					1	1		1	••	• •						3	3	19	6.33
Age not stated			1													1	1	3	3.00
Totals	561	476	413	346	241	169	118	89	45	17	19	11	10	5	2	2,499	2.522	9,039	8-15

^{*}One of these mothers bore twins.

† Two of these mothers bore twins.

† Three of these mothers bore twins.

The reference marks in this table indicate the age of the mother and the number of the child in the cases of plurality births; thus at the age of 27 four mothers bore twins; in one case they were the first and second children; and in three cases, the second and third children.

There were three women who bore children in Providence in 1879 who were fifty years old; one of whom had borne four children before; one, five; and one seven. The number of women of forty years and over, who bore children in 1879, was 115; they had borne 821 children, or an average of 7.14 children to each mother. These facts concerning the age of the mother have been recorded since July 1, 1863, a period of sixteen years and six months; in that time there have been 1,669 women of forty years and over, who have borne children; the number of children borne by them being 12,702, there is an average of 7.61 children to each mother.

During this period, since July 1, 1863, the whole number of children borne by mothers whose age was stated, was 34,097. The number of children borne by mothers of each age during this time is as follows:

Age of mother. Children.	Age of mother. Children. 272,041	Age of mother. Children.
15 19	282.387	41 218
16 50	291,712	42 262
17 147	302,753	43 147
18 876	811.145	44 97
19 806	321,611	45 102
20 974	88 1,842	46 27
21 1,141	84	47 24
★21,65 5	851,718	48 11
23	36 1,019	49 3
242,028	87 830	50 8
252,288	88 903	
28 2,114	89 584	All known ages.84,097

The mothers of these children were divided as follows, as to age: —

Under twenty years	1,202
Twenty, and under twenty-five	7,627
Twenty-five, and under thirty1	0,542
Thirty, and under thirty-five	7,989
Thirty-five, and under forty	5,049
Forty, and under forty-five	1,513
Forty-five, and over	175
	10000
Total	34,097

Of these 34,697 children, 19,371 or 56.81 per cent. were borne by mothers less than thirty years old; 13,038 or 38.24 per cent. by mothers between thirty and forty years old; and 1,688 or 4,95 per cent. by mothers forty years old, or more.

PLURALITY BIRTHS.

The number of plurality births in Providence in 1879, was 23, which was seven less than in 1878; all were twin births.

The following particulars concerning them are reported:— Cases, 23; number of children, 46.

Locality. First ward, 3; second, none; third, 3; fourth, 1; fifth, 1; sixth, 1; seventh, 1; eighth, 4; ninth, 3; tenth, 6.

Season. January, 2: February, 2; March, none; April, 2; May, 2; June, 2; July, none; August, 4; September, 3; October, none; November, 2; December, 4.

Sex. In nine cases, both children were males; in nine both were females; in five cases there was one male and one female; making 23 children of each sex.

Color. In twenty cases the children were of white parents; in three of colored parents.

Number of the Children and Age of the Mother. From Table VII. may be shown the age of the mother in each case of plurality birth. In three instances they were the first and second children of mothers 23, 27 and 28 years old; in four instances they were the second and third children of mothers, three of whom were 27 years old; and one 28 years old: in one case, the third and fourth

children of a mother 34 years old; in four cases, the fourth and fifth children of mothers 21, 24, 28 and 35 years old; in four, the fifth and sixth children of mothers, two of whom were 35 years old; one, 33; and one 36; in four, the sixth and seventh children of mothers, two of whom were 34 years old; one, 33; and one, 36; in one case, the seventh and eighth children of a mother 39 years old; and in two cases, the eighth and ninth children of mothers 34 and 35 years old. In thirteen cases the mothers were more than 30 years old.

Parentage. In eleven cases, or nearly half, the parents were both American; in three of these cases the parents being colored; in four cases, both were Irish; in three cases, both were English; in two, both were Scotch; in two, both were of the Western Islands (or Portuguese); and in one both were German.

TWENTY-FIVE YEARS' SUMMARY OF PLURALITY BIRTHS.

In the twenty-five years, 1855-1879 inclusive, there have been in Providence 583 cases of plurality births; 574 of twins, and 9 triplets; the whole number of children being 1,175.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children, and one female; making 9 male and 18 female children in the nine cases.

In the 574 cases of twins, there were 186 in which both of the children were males; 198 in which both were females, and in 190 there was one child of each sex; making 562 male and 586 female children. Of the 1,175 children in all the cases of plurality births, there were 571 male, and 604 female children; this is in the proportion of 105.8 females to each 100 males; or 51.40 females and 48.60 males, in each 100 of the children, in the cases of plurality births.

Color. In all the cases of triplets the parents were white. Of the 574 cases of twins in 558 the parents were white; and in 21 they were colored.

Parentage. Of the triplets, in one case, the parents were both American; in five cases both were Irish; in one case, both German; in one, the father was English, and the mother Irish; and in one the father Scotch, and the mother Irish.

In the following table for the 574 cases of twin-births, the nativity of both parents is given; the first letter representing the the father's birth-place, and the second the mother's.

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British American; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish.

Parentage. No.	Parentage. No.	Parentage. No.	Parentage. No.
A. and A 191	I. and A 13	G. and I 1	A. and S 1
I. and I222	E. and A 12	B. A. and I 1	I. and S 3
E. and E 28	S. and A 2	Sw. and I 1	E. and S 1
S. and S 6	B. A. and A 6	It. and I 1	I. and W 1
G. and G 16	G. and A 1	P and I 1	A. and B. A 5
B. A. and B. A 10	Sw. and A 1	A. and E 3	I. and B. A 2
D. and D 1	A. and I 12	I. and E 6	E. and B. A 1
P. and P 7	E. and I 11	S. and E 1	S. and B. A 1
It. and It 1	8. and I 3	G. and E 1	Total574

Of the 574 cases of twin-births, there were 226. or 39.37 per cent. of all, where the mothers were natives of the United States; 253, or 44.08 per cent., where the mothers were natives of Ireland. Of the 583 cases of plurality births in the twenty-five years the American mothers numbered 227; the foreign mothers, 356.

Proportion of plurality births to whole number. The following numbers show how the year 1879 compares with the twenty-four years previous, 1855-1878 inclusive, as to the whole number of births and the proportion of plurality cases:—

24 years, 1855-1	Single births,		Triplet.	Whole No. Births, 44 931	Whole No. Children.
1879					
25 years' total	46,847	574	9	47,430	48,022

PLURALITY BIRTHS.

Proportion of cases of plurality births:-

*	Twins.	Triplets.
1855-1878	1 in 81.5	1 in 4,992.8
1879	1 in 108.7	•••••
1855–1879	1 in 82.6	1 in 5,270
Proportion of o	children of plurality births:-	_
	Twins.	Triplets.
		Triplets.
1855–1878	Twins.	Triplets

Proportion of plurality births by parentage. The number of births in the twenty-five years, 1855-1879 inclusive, was 47,430. The following is a statement of the number of births of American, foreign and mixed parentage; and the number and proportion of plurality births in each class:—

		Plurality cases. 1 in Per cer
	•	192 99.10, or 1.0
		335 67.82, or 1.0
American and Foreign	5,681	56101.45, or
Total	47,430	583 81.36, or 1.5

Of the 583 cases of plurality births, the 192 cases of American parentage were 32.93 per cent.; the 335 cases of foreign parentage, were 57.46 per cent.; and the 56 cases of mixed parentage were 9.61 per cent.

MARRIAGES.

During the year 1879, there were 1,071 marriages in the city of Providence.

In Table VIII. is given the number in each month and quarter of each of the seven years, 1873-1879 inclusive; also the aggregate and average number for the corresponding seasons in the period of twenty-five years, 1855-1879 inclusive:—

TABLE VIII. MARRIAGES. Season.

MONTHS.	1879.	1878.	1877.	1876.	1875.	1874.	1873.	25 years. 1855-1879.	25 years average.
January	102	76	68	79	75	93	105	1.801	79
February	- 85	78	65	93	79	87	111	1,611	64
March	47	73	38	44	71	56	49	1,057	42
First quarter	234	222	171	216	225	236	265	4.469	178
April	83	77	83	70	84	95	87	1,633	65
May	98	68	85	80	82	113	91	1,784	71
Jane	96	104	83	94	100	101	107	1,858	75
Second quarter	277	249	251	244	266	309	285	5,275	211
July	59	94	85	56	101	88	105	1,578	63
August	88	62	76	62	73	80	63	1,369	55
September	106	98	87	93	111	129	108	1,909	76
Third quarter	253	249	248	211	285	297	276	4,856	194
October	103	99	99	95	107	127	118	2,121	- 85
November	128	115	102	109	114	114	125	2,290	92
December	76	82	82	74	82	93	81	1,676	67
Fourth quarter	307	296	283	303	334	334	324	6,087	244
Yearly totals	1,071	1,016	953	949	1.079	1,176	1,150	20.687	827

The number of marriages in 1879 was 55 more than in 1878, and was larger than in any year since 1875; it was 244 more than the average number for twenty-five years. Having estimated the population of the city in the middle of the year 1879 at 103,000, we find the following ratios: Number of persons married, 2,142, which is one in each 48.08 of the population, or 20.79 in each thousand. The ratio of persons married to children born was 85 to 100; of persons married to deaths 105.7 to 100. In 1878 the ratio of persons married to children born was 77 to 100; of persons married to deaths 102.2 to 100.

Season. The distribution of these 1,071 marriages in the different seasons follows the usual rule; the larger part of them being found in the last six months of the year, and the largest quarterly total, 307, being that of the fourth quarter; the smallest quarterly total, 234, was in the first quarter.

Dividing the year by the calendar seasons, and taking the aggregate of marriages for twenty-five years, we find that of each 100 marriages in Providence about 25 occur in the winter months, December, January and February; 22, in the spring months; 23, in the summer, and 30 in the autumn. The largest monthly total in 1879 was 128, in November; the smallest, 47 in March.

The following figures show the number and percentage of marriages in each quarter of the year 1879, and for the corresponding seasons for the period of twenty-four years, 1855–1878, inclusive:—

	18	79.	1855—1	1878.
3	farriages.	Per Cent.	Marriages.	Per Cent.
January-March	284	21.85	4,285	21.59
April-June	277	25.86	4,998	25.48
July-September	253	23.62	4,603	23.46
October-December	807	98.67	5,780	29.47
Total	 1,071	100.00	19,616	100.00

Nativity. Table IX. shows the birth-places of both parties in each of the 1,071 marriages of the year 1879:—

TABLE IX. MARRIAGES, 1879. Nativity of the Parties.

	BIRTH-PLACES OF BRIDES:							rooms
BIRTH-PLACES OF GROOMS.	United States.	Ireland.	England, Scot- land and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number of Grooms
United States	653	44	26	1	20		7	751
Ireland	37	125	5		6		2	175
England, Scotland and Wales	36	10	27		3			76
Germany	7	3		5			2	17
British America	10	2	1		9			21
Portugal and Western Islands						5		2
Other Countries	13	3		2			7	25
Whole number of brides	756	187	59	8	38	5	18	1,071

Bride and Groom born in the same Country. In 653 of these marriages, or 60.97 per cent. of all, both parties were born in the United States; this is 52 more than the same class of marriages in the previous year, and the proportion to the whole number 1.82 per cent. more. In 125 marriages, or 11.67 per cent. of all, both bride and groom were born in Ireland; this is 15 less than in 1878, and the proportion 2.11 per cent. less. There were also 27 marriages in which both parties were born in England, Scotland, or Wales; in 5, both were born in Germany; in 9, both in British America; in 5, both in Portugal or Western Islands; in 7, both were born in other countries than those mentioned in the table. There were therefore 831 marriages in which the bride and groom were natives of the same country; this was 77.59 per cent. of all the marriages. In 1878 the number was 799, or 78.64 per cent. of all.

Bride and Groom born in different Countries. In 240 marriages, or 22.41 per cent. of the whole number, the bride and groom were born in different countries; this is 23 more than in 1878, and the proportion 1.04 per cent. more. Of the 751 grooms born in the United States, 98, or 13.04 per cent., married natives of other countries; the number and proportion are both rather larger than in the previous year; of the 175 born in Ireland, 50, or 28.57 per cent., married natives of other countries; of the grooms born in England, 64.47 per cent. married natives of other countries; of those born in Germany, 75.59 per cent.; of those born in British America, 59.09 per cent. Of the 756 brides born in the United States, 103, or 13.62 per cent., married natives of other countries; the number and proportion are larger than in 1878. Of the 187 brides born in Ireland, 62, or 33.16 per cent., married natives of other countries; of those born in England, 52.54 per cent.; of those born in Germany, 37.50 per cent.; of those born in British America, 76.32 per cent.

The nativity of the 25 grooms born in "other countries" was as follows: 8 born in Italy; 4 in Sweden; 3 in Switzerland; 2 each in France and Cuba; and 1 each in Algiers, Australia, Holland, Poland, Russia and Tuscany.

Of the 18 brides born in "other countries," 7 were born in Sweden, 3 in Italy, 2 each in France and Switzerland, and 1 each in Bohemia, Denmark, South America, and Spain.

The number and percentage of American, foreign and mixed marriages in each of the twenty-five years from 1855-1879 inclusive, are given in the following table. By mixed marriages are understood those where one of the parties was of American, and the other of foreign nativity:

	Ame		7.75	eign.	Mixed:
Year.	Number.	Per cent.		Per cent.	404000000000000000000000000000000000000
1855	. 308	48.73	265	41.93	59 9.
856	. 852	54.74	212	32.97	7912.
1857	. 384	62.54	175	28.50	55 8.
858	283	50.54	214	38.21	6311.
859	. 333	54.50	224	36.66	54 9.
860	. 336	54.19	225	36.29	59 9.
861	804	55.58	182	33.27	6111
862	. 288	52.65	179	32.72	8014
863	342	56.07	192	31.47	7612
864	427	57.78	211	28.55	10113
865	417	58.82	202	28.49	90 12
866	471	58.00	244	30.05	9711
867	485	56.73	269	31.45	10111
868	. 530	60.78	215	24.66	12714
869	. 530	56.99	286	30.75	11412
870	. 552	57.56	262	27.32	145
871	. 523	55.46	278	29.48	14215
872	. 591	54.52	305	28 14	188
873	652	56.69	317	27.57	18115
874	648	55.10	346	29.42	18215
875	. 586	54.31	320	29.66	173 16.
876	. 547	57.64	241	25.39	16116.
877	. 545	57.19	233	24.45	17518
878	601	59.15	245	24.12	17016.
879	653	60.97	217	20.26	20118.

The proportion of mixed marriages is larger than in any previous year; the proportion of American marriages is also larger than in any year except 1857; while that of foreign marriages is considerably less than ever before. The whole number of marriages as given here is 20,681; while in Table VIII. the total for the same period of twenty-five years is given as 20,687. This discrepancy is caused by the omission from Table VIII., in the early years of the records, of eleven marriages, because it was not stated in what month they occurred; and the omission from the foregoing table of seventeen marriages, because the nativity of the parties was not given. If these numbers are added to their respective tables, the total, in each case, will be the same (20,698). In many years some marriages have been handed in too late to be recorded in the report for the year; but they have

since been added in the General Summary at the end of this Report; so that the total (20,716) from which the average number (829) given in Table XXI. is obtained, is the true number of marriages for the period of twenty-five years.

Marriages of colored persons. The number of marriages of colored people in Providence in 1879 was 43; this was eleven less than in 1878, and three less than in 1877. The aggregate and average age of the parties in these marriages were as follows:—

45 males Aggregate age, 1,341 yearsAverage age, 31.19 years	ars.
& femalesAggregate age, 1,143 yearsAverage age, 26.58 years	ars.

The whole number of marriages of colored persons in the twenty-four years, 1856-1879, was 871; the aggregate and average age of the parties as follows:—

871 malesAggregate age, 25,566	yearsAverage age, 29.35 years.
871 femalesAggregate age, 23,855	yearsAverage age, 26.24 years.

Age of persons married. In Table X. are shown the number and percentage of brides and grooms in each division of ages:—

TABLE X. MARRIAGES, 1879. Age of the parties.

	AGES OF THE BRIDES.							9.
AGES OF THE GROOMS.	Under 20 years.	20 to 25.	25 to 30.	80 to 40.	40 to 50.	50 to 60.	Whole number of Grooms.	Percentage Grooms
Under 20 years	16	8		•••••			24	2.24
99 to 95	106	214	29	5	1	 •••••	855	83.15
25 to 30	85	161	127	26	1	 .	850	32.68
36 to 40	9	64	77	77	7	1	235	21.94
40 to 50	8	8	7	96	16	8	63	5.88
50 to 60	 .	8		7	9	4	23	2.15
8 0 to 70	 .		1	6	8	6	16	1.49
70 to 80		·····	 .		1	4	5	.47
Whole number of Brides	169	458	241	147	88	18	1,071	
Percentage of Brides	15.78	42.77	22.50	18.72	8.55	1.68		100.00

In 42.39 per cent. of these marriages, the bride and groom were in the same division of ages; of the whole number of grooms, 379 or 35.39 per cent. were less than 25 years old; of the brides, 627 or 58.54 per cent.

The aggregate and average age of all persons married during the year were as follows:—

1,071 males.......Aggregate age, 31,016 years.......Average age, 28.96 years.
1,071 females......Aggregate age, 27,103 years......Average age, 25.32 years.

The following is a statement of the same facts for each of the twenty-four years, 1856-1879 inclusive, and for the whole period:

	Males,	Females.
Years.	No. Aggregate age. Average age.	No. Aggregate age. Average age
1856	639 18,104 years28.33 years	637 15,559 years24.42 years
1857	607 16.753 years27.59 years	606 14,427 years 23.84 years.
1858	560 15.850 years28.29 years	560 13,702 years24.47 years
1859	611 16,989 years27.81 years	611 14,616 years23.92 years.
1860	620 17,135 years27.60 years	620 14,667 years23.66 years.
1861	547 15,448 years28.24 years	547 13,316 years22.33 years.
1862	547 15,773 years28.83 years	547 13,565 years24.80 years.
1863	610 17,208 years28.21 years	610 14,792 years24.24 years.
1864	739 21,361 years28.90 years	739 18,075 years24.45 years.
1865	709 20,539 years28.96 years	709 17.605 years24.83 years.
1866	812 22,801 years28.08 years	812 20,036 years 24 67 years.
1867	855 24,443 years28.58 years	855 21.078 years24.65 years.
1868	872 24,546 years28 15 years	872 21,440 years24.59 years.
1869	930 26,548 years28.55 years	930 23,135 years24.88 years.
1870	959 27,343 years28.51 years	959 23,773 years24.79 years.
1871	943 26,743 years28.37 years	943 23.290 years24.70 years.
1872	1.084 31,110 years28.70 years	1,084 26,835 years 24.76 years.
1873	1,150 32,879 years28.59 years	1,150 28,701 years 24.96 years.
1874	1,176 33.927 years28.85 years	1,176 29,510 years25 09 years.
1875	1,079 30,848 years28.59 years	1,079 27,090 years 25 11 years.
1876	949 27,496 years28.97 years	949 23,640 years24 91 years.
1877	953 27,904 years29.28 years	953 24,092 years25 28 years.
1878	1,016 29,460 years28.99 years	1,016 25,796 years25.40 years.
1879	1,071 31,016 years28.96 years	1,071 27,103 years25.32 years.
24 years.	20,038572,224 years28,56 years5	20,035495,843 years24.75 years.

The number of males and females does not agree in this table, because in 1856 the ages of 17 males and 19 females, and in 1857 the ages of 7 males and 8 females, were not given.

Number of the marriage. Table XI. shows how many of the persons married in Providence in 1879 were married for the first, second, or third time:—

TABLE XI. MARRIAGES, 1879. Number of times married.

GROOMS.	BRIDES. 1	Whole		
Number of the Marriage.	First.	Second.	Third.	number of Grooms.
First	823	57	2	882
Second	96	70	5	171
Third	10	8		18
Whole number of Brides	929	135	7	1,071

In 823 cases, or 76.84 per cent. of all, it was the first marriage of both parties, both the number and percentage being larger than in 1878; in 70 cases, or 6.54 per cent., it was the second marriage of both; of this class the number was a little larger, but the percentage a little smaller than in the preceding year.

Denominational. The 1,071 marriages of the year 1879 were performed by the clergymen of the various denominations of the city, and by civil authority as follows:—

Roman Catholic	Seamen's Mission Clergyman 9
Calvinistic Baptist148	Lutheran 4
Free Baptist 29	Advent Christian 2
Six Principle Baptist 2	Evangelical Advent 8
Methodist	New Jerusalem 1
Episcopalian112	Latter Day Saints 10
Congregational 89	Hebrew 2
Unitarian 87	Friend's Ceremony 1
Universalist 45	Denomination not specified 8
Presbyterian 24	Justice of Supreme Court 2
Christian 4	Total

Nativity and Residence. In 87 marriages, or 8.12 per cent. of the whole number, both parties were born in the city; there were also 118 marriages in which the groom alone was born here; and 192 in which the bride alone was born here; there were therefore 484 persons married who were natives of Providence; this is 22.60 per cent. of the whole number of persons married; the number and proportion being larger than in 1878. The number of marriages in which both parties were non-resident was 182, or 16.99 per cent. of all. This is a much larger number than usual.

Marriage and Education. There were 2,142 persons married in Providence in 1879; as regards sex and nativity they were divided as follows:—

Born in United StatesMales,	751Females,	756 Total, 1,507
Born in foreign countriesMales	, 320 Females	315 Total, 535

In each class, the number and proportion who signed the marriage certificate with a mark were as follows:—

1879.	Whole number married.	Signed with a mark.	In each
Males, born in the United States	751	32	4.25
Females, born in the United States	756	48	6.55
Total, born in the United States	1,507	80	
Males, born in foreign countries	320	66	20.62
Females, born in foreign countries	315	94	29.84
Total, born in foreign countries	635	160	
Totals	2,142	240	11 20

There were 80 persons, born in the United States, and married in 1879, who signed their marriage certificates with a mark; of this number, 27 were colored persons, (10 males and 17 females); and 39 were of foreign parentage, (14 males and 25 females); this leaves 14 persons, of white American parentage who signed the marriage certificate with a mark; this is 5 more than in 1878.

The number and proportion of those who signed with a mark is 1879 was larger than in 1878, in all classes, except "females born in foreign countries;" of those born in the United States, 2

more; 10 males and 15 females; of those born in foreign countries, the whole number was 6 less than in 1878; the number of females being 13 less, and of males 7 more. In 1878 the whole number of persons married was 2,032; of those who signed with a mark, 221, or 10.88 in each hundred.

The statistics on this subject for the twenty years, 1859-1878 inclusive, are as follows:—

1859–1878.	Whole number married.		
Males, born in United States	10,848	529	4.88
Females, born in United States	11,050	746	6.75
Total, born in the United States	21,898	1,275	5.89
Males, born in foreign countries	6,818	2,093	88.18
Females, born in foreign countries	6,111	8,011	
Total born in foreign countries	12,429	5,104	41.00
Totals		6.879	18.56

DEATHS.

During the year 1879 there were 2,026 deaths in the city of Providence; 37 more than in the year 1878, and more than in any previous year. The number of males was 35 more than in 1878; of females, 2 more; the number of whites 31 more; of colored, 6 more; of natives of the United States, 10 more; of persons born in foreign countries, 27 more; of American parentage, 52 more; of foreign parentage, 15 less.

The average death-rate in Providence for the twenty-five years 1855-1879 inclusive, was one in 50.78 of the population. In 1879 it was as follows in the various classes of the population:—

				Deaths to
			To population, one death in	
Whole number	100,675	2,026	49.69	20.12
Males	48,701	1,004	48.51	20.62
Females	51,974	1,022	50.86	19.66
Whites	97,188	1,914	50.78	19.69
Colored	3,487	112	31.13	32.12
Born in the United States	71,438	1,575	45.36	22.05
Born in Ireland	18,458	334	55.26	15.00
Born in England, Scotland and Wal	es. 6,110	63	96.98	10.31
Born in Germany	1,246	16	77.87	
Born in other foreign countries				
Total of foreign born	29,237			15.43
American parentage	47,816	941	50.28	19.89
Irish parentage	37,440	776	48.25	20.73
English, Scotch and Welsh parentag	ge. 9,636	169	57.02	
German parentage	2,516	49	51.35	19.47
Other foreign parentage	3,767	91	41.40	24.16
Total of foreign parentage	53,359	1,085	49.18	20.33

Table XII. gives the general facts concerning the deaths that occurred during the year 1879:—

Table XII. Deaths, 1879.

Sex, condition, color, nativity, parentage, locality, season.

				l			1		1				
1879.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Whole year.
Whole number of Deaths	200	151	163	167	152	120	187	173	134	167	200	213	2,026
Sex													
Males	111	77	80	80	68	53	89	92	63	81	98	109	1,004
Females	89	74	83	87	81	67	98	80	71	83	102	104	1,022
Condition.—										ĺ	1		
Married	58	51	47	57	47	46	57	48	31	46	50	57	593
Single	104	84	92	81	81	60	112	99	80	107	129	137	1,166
Widows	21	11	19				18	17			14	15	178
Widowers	17	5	5	10	8	8	5	8	10	4	7	4	86
Divorced	••••	••••	• • • •	1	••••	••••	••••	• • • •	••••	• • • •		••••	1
Color.—	'												
Whites	186	139	152	151	146	116	176	162	125	162	196	203	1,914
Colored	14	12	•11	16	6	4	11	10	9	5	4	10	112
Nativity.—													
United States.	151	114	134	116	116	84	154	131	107	138	161	169	1,575
reland.	36	23	28	35	25	26	20	32	20		30	36	331
England and Scotland	9	6	8	7	6	6	9	4	4	1	5	3	63
Germany	1	4	••••	4	1	1	1	1			2	1	16
British America	2	4	2	5	8	• • • •	2	2	3		1	3	27
Portugal & West'n Islands.		•••	1	•••	· • •	1	•••	••••		• • • •	1	••••	8
Other countries	1	••••	• • • •	••••	1	2	1	2	••••	¦••••		1	8
Parentage.—										ĺ			
American	107	75	89	72	74	48	90	79	57	75	82	93	941
Irish	63	54	51	65	58	53	67	70		68	92	87	776
English and Scotch	20	14	12	15	13	10	19	10		1	13	24	169
German	4	4	1	6	2	8	6	4	6		6	3	49
British American	4	8	5	4	6	1	3	4	5	ı	3	3	45
Portuguese.		1	3	2	2	1	1	2	1	5	1 -	1	22
Other foreign	2	••••	2	8	2	4	1	3	1	3	1	2	24
Locality, Wards.—													
I	22	20	28	20	20	11	25	80		14		31	279
II	10	8	5	6	6	4	9	6	_	8		5	68
IV	19 18	13 16	16 9	11 9	12 11	7	16 6	23 6			20	22	194
V	20	13	17	23		6 14	19	16	-		9 21	6 21	116
VI	6	10	7	9	5	5	14	7	8			13	269 106
VII	17	13	16			17	18	14		_		20	180
VIII	32	6	14			9	15	18				24	198
1X	29	28	24	30		13	28	21	16		1	2:	200
X	18	17	24		25	29	26	25				39	305
Public Institutions	9	7			8	6	11	6	5		8	6	81
Totals	200	151	163	167	152	120	187	172	134	167	200	213	2,026
				1 -0.	-02	120	201		107	101	200	213	2,020

Season. The year 1879, like the preceding year, forms an exception to the general rule in regard to the distribution of deaths in the different quarters of the year. The largest number of deaths, instead of being in the third quarter of the year, as is nearly always the case, is this year found in the last quarter; the number in the first quarter is also larger than that in the third thus putting the third quarter the third in order, which has only happened once before in the twenty-five years during which these reports have been published.

The following figures show the number of deaths in each quarter of the year 1879, and in each of the five years preceding; also the aggregate and average number in each quarter for the period twenty-four years, 1855–1878 inclusive:—

	1879.	1878.	1877.	1876.	1875.	2-	i years, 18	Ark—
January-March								
April-June								_
July-September								_
October-December	580	505	530	490	495	486	7,683	
Whole year	2,026	1,989	1,938 1	1,865	1.915	1,987	81,464	1,517

The number and percentage of deaths in each quarter of the year 1879, and in the aggregate for the twenty-four years, 1855-1878, are as follows:

	1879.		24 years,	1866-1878.
1	Deaths.	Per cent.	Deaths.	Per cest.
January-March	514	25.87	7,519	93.99
April-June	489	21.67	6,909	11.91
July-September	498	24.33	9,333	99.73
October-December	580	28.63	· 7,683	94.6
•				
Totals	.2,026	100.00	81,464	300.0

The number and percentage of deaths in each quarter of each year, during the period of twenty-five years, 1855-1879, were as follows:

	July-Sept.	OctDec.
April-June.	Deaths. Per cent.	Deaths. Per ct.
Deaths. Percent.	Dentum 1 ct out	\$31\$3.31
Descris. Percent.	55954.31	90.87
9.91	99697.80	
10.78		30231.24
#714.	95795.27	9.519.50
00.00	97030.04	311
9090.88	99799.67	95395.18
93693.45	91.07	930 93.74
93.45		954 97.79
319321.01.	97129.59	24 70
494890.43.	37031.14	301 34.17
ro 979 91.81	40881.46.	30523.50
m 94019.88	37330.72.	30034.77
.9791821.04	29821.63.	21223.36
.8890291.04	99499.79	21422.29
1.8121421.98	90 91 17	98096.04
1.8121421.98		919 9194
.6.3525019.90)28.91	
28.9021.4	5 .3 71 39.3 7	99093.97
31.8096390.9	838530.70	99598.51
90.50	1518	l9194.39
.34.84 407 23.6	B50499.3	3 381 33.16
.22.3550125.5	556 97.9	818591.46
.25.4840120.5	n. 101 97.7	z9595 85
.25.4840130.3	P}2912i.i	n 400 ad 87
	4757850.7	399030 36
21.5144022.	.7955728.7	427.35
	.579898	14\$65\$5.89
25.372921	.5ī 24.1	1858028.60

and percentage of deaths in the fourth quarter of were the largest that have ever been recorded, er for the third quarter was less than in any year of the percentage less than in any previous year.

This departure from the usual rule was the prevament of the deaths from that disease being in the control of the deaths from that disease being in the latter part of deaths from diarrhical disease the percentage from a large proportion of the deaths from diarrhical disease the percentage from a large proportion of the deaths from that disease in the figure of deaths from the figure of deaths from diarrhical disease the percentage in the figure of deaths from the figure of deaths from the figure of deaths from diarrhical disease in the figure of deaths from the figure

The number of deaths in the first quarter, though less than in 1878, was larger than in the first quarter of any other year; the percentage was less than in several other years, but yet above the average.

Sex and Parentage. The 2,026 decedents in 1879 were divided as follows as to sex and parentage:—

American parentage46	9 males472	femalesTotal, 901
Foreign parentage53	5 males550	femalesTotal, 1,085
_	_	_
Total1,00	4 males1,022	females Total, 2,036

The decedents were very evenly divided as to sex, the number of females being only 18 more than of males; of these, 3 were of American parentage, and 15 of foreign parentage. The whole number of foreign parentage was 144 more than of American parentage; of males 66 more, of females 78 more. The decedents of both classes during the twenty-five years, 1855–1879 inclusive, were as follows in regard to sex:

1855-1879.......Males, 16,427.......Females, 17,063..........Total, 33,490

Condition. The number of married persons who died in 1879 was 595; 283 of American, and 312 of foreign parentage; the whole number was 106 more than the decedents of this class in 1878. The number of decedents in 1879 who were single was 1,166; 512 of American, and 654 of foreign parentage; this was 84 less than in 1878. The number of widows who died in 1879 was 178; 100 of American and 78 of foreign parentage; 9 more than in the previous year. The number of widowers among the decedents of 1879 was 86; 46 of American and 40 of foreign parentage; 12 more than in 1878. There was but one divorced person among the decedents of 1879; in 1878 there were 7.

Color. There were 31 more deaths of white persons in 1879 than in 1878; of colored persons 6 more. The number of colored children born in Providence in 1879 was 104; the number of colored decedents, 112; the proportion of births to deaths, 100 to 107.7 During the twenty-five years, from 1855–1879 inclusive, the number of births among the colored people of the city has been 1,847; the number of deaths 1,953; the proportion of births to deaths 100 to 105.7.

Nativity and Parentage. From Table XII. important comparisons may be made between the number of decedents registered according to parentage and those registered according to nativity. The number of decedents in 1879 of American nativity was 1,575, or 77.74 per cent. of all; while the number of American parentage was only 941, or 46.45 per cent. of all. The number of American nativity was 10 more than in 1878; of American parentage 52 more.

The number of decedents of foreign birth in 1879 was 451, or 22.26 per cent. of all; the number of foreign parentage 1,085, or 53.55 per cent. of all. The number of decedents of foreign birth in 1879 was 27 more than in 1878; of foreign parentage 15 less.

The nativity of the 8 decedents born in "other countries" was as follows: 3 born in Italy, 1 each in France, Wales, Austria, Belgium and Norway.

Of the 24 of "other foreign" parentage, there were 7 of Italian, and 5 of Swedish parentage, 4 French, 4 Swiss, and 1 each Belgian, Norwegian, Polish, and Welsh.

Locality. Table XII. shows the number of deaths in each ward and in the Public Institutions of the city for each month and for the whole year. Comparing the numbers with those of the preceding year, we find in the first ward 15 more deaths in 1879 than in 1878; in the second, 6 more; in the third, 12 more; in the fourth, 7 more; in the fifth, 20 more; in the sixth, 2 less; in the seventh, 11 more; in the eighth, 11 more; in the ninth, 9 less; in the tenth, 30 less; in Public Institutions, 4 less. In Wards 1, 2, 3, 4, 5, 7 and 8 collectively, the number of deaths was 82 more than in 1878; in Wards 6, 9 and 10 and Public Institutions, the number was 45 less. The number of decedents in each ward, according to parentage, was as follows:—

Wards.	American.	Foreign.	Wards.	American.	Foreign.
I	127	152	VII	88	92
II	54	14	VIII	118	80
ш	73	121	ıx	138	152
IV	57	59 .	X	85	220
v	81	128	Public Ins	titutions 46	85
v1	74	32			
Who	ole city			941	1,065

Average Age. Table XIII. gives the number of deaths, and the aggregate and average age of the decedents, according to parentage, for each month and for the whole year, 1879:—

Table XIII. Deaths, 1879.

Aggregate and Average Age, and Parentage.

	Ame	rican Pa	rentage.	For	eign Par	entage.	Tota	Total American and Foreign.		
MONTHS.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	
January	107	3,883	36.20	98	2,829	30.32	200	6,712	83.56	
February	75	2,666	35.54	76	2,183	28.72	151	4,849	32.11	
March	89	3,188	35.82	74	1,852	25.03	163	5,040	80.92	
April	72	2,411	33.49	95	3,407	35.86	167	5,818	34.84	
May	74	2,983	40.31	78	1,964	25,18	152	4,947	32.51	
June	48	1,388	28.92	72	2,470	34.31	120	3,858	32.15	
July	90	2,684	29.62	97	2,236	23,05	187	4,920	26.31	
August	79	2,301	29.13	93	2,362	25.40	172	4,663	27.11	
September	57	1,762	30.91	77	1,834	23.82	134	3,596	26.87	
October	75	2,178	29.03	92	1,815	19.78	167	3,993	23.91	
November	82	2,215	27.01	118	2,588	21.93	200	4,803	24.00	
December	93	2,237	24.05	120	2,906	24.22	213	5,143	24.14	
Whole year	941	29,896	81.77	1,085	28,446	26.22	2,026	58,342	28.80	

The average age of the 2,026 decedents in 1879, was 28.80 years; 2.71 years more than the average age of all decedents in 1878. The average age of the 941 American decedents in 1879 was 31.77 years; 1.71 years more than in 1878; of the 1,085 decedents of foreign parentage, 26.22 years; 3.33 years more than 1878.

Comparing this table with the same for 1878, we find the following differences in the different months. Among the decedents of American parentage the average age was higher in each of the first five months of the year 1879 than in the corresponding

months of the previous year. In June the average in 1879 was 11.49 years less than in 1878. Of the remaining six months, the average age in 1879 was greater than in 1878, for the months of July, September and October, and less than in 1878, for the months of August, November and December. Among the decedents of foreign parentage the average age was more in 1879 than in 1878 in the months of January, February, March, April, June, July and August; and less in the months of May, September, October, November and December.

Table XIV. is a recapitulation of the same facts for the twenty-four years, 1856–1879 inclusive.

Table XIV. Deaths, Twenty-Four Years, 1856-1879.

Recapitulation. Aggregate and average age by parentage.

Twenty-Four	American Parentage.				gn Pare	ntage.	Total American and Foreign.		
YEARS, 1856-1879.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.
January	1,447	49,719	84.86	1,259	31,813	25.27	2,706	81,532	80.18
February	1,300	44,125	88.98	1,213	27,286	22.49	2,518	71,411	28.42
March	1,365	47,645	84.90	1,236	30,591	24.75	2,601	78,236	30.08
April	1,326	44,908	83.87	1,281	33, 876	26.44	2,607	78,784	30.22
Мау	1,239	45,270	36.54	1,159	29,835	25.74	2,398	75,105	81.89
June	1,033	85,437	84.80	1,102	27,143	24.63	2,135	62,580	29.31
July	1,447	87,390	25.84	1,721	81,838	18.50	8,168	69,228	21.8
August	1,578	43,816	27.45	1,826	33,627	18.42	3,404	76,918	22.60
September	1,898	41,089	29.89	1,537	29,857	19.43	2,935	70,946	24.17
October	1,260	41,186	32.69	1,395	31,549	22.62	2,655	72,735	27.8
November	1,297	42,806	88.00	1,836	83,340	24.96	2,633	76,146	28.9
December	1,407	49,702	85.32	1,337	84,033	25.47	2,744	83,735	80.5
Total	16,097	522,593	82.47	16,402	374,788	22.85	32,499	997,381	27.6

By examining these two tables we find many differences between the average age of American decedents and that of foreign decedents in the different months. In the statistics for the year 1879 (Table XIII.), we find the highest average age of American decedents to be 40.31 years in May; the lowest average age 24.05 years in December. Of the decedents of foreign parentage, the highest average age was 35.86 years in April; the lowest, 19.73 years in October. The greatest difference between the two classes was in May, when the average age of American decedents was 15.13 years greater than that of decedents of foreign parentage; the least difference was in December, when the average age of American decedents was 0.17 of a year less than that of the other class. The year is an exception to the general rule in that there were three months when the average age of foreign decedents was greater than that of American decedents; in April it was 2.37 years greater; in June, 5.39 years; in December, 0.17 of a year. For the whole year, the average age of the American decedents was 5.55 years the greatest.

In the statistics for the longer period of time, as given in Table XIV., we find similar contrasts. In this table we find that in every month the average age of American decedents was greater than that of the foreign class. The highest average age of American decedents was 86.54 years in May; and the lowest, 25.84 years in July. The highest average age of decedents of foreign parentage was 26.44 years in April; the lowest, 18.50 in July. The greatest difference between the two classes was 11.44 years in February; the least 7.34 years in July. The difference for the whole year in the long period of time was 9.62 years. It will be seen from the two Tables, XIII. and XIV., that in both classes of decedents in 1879, the average age in the months of October, November and December was much below the average for those months in the long term of years. This was owing to the epidemic of scarlatina, which prevailed, as already mentioned on page 31, in the last quarter of the year. It was fatal mostly among very young children, increasing the number of decedents of that class, and consequently reducing the average age of all decedents.

The following table shows the difference between the two classes of decedents in each month, for the year 1879, and for the period of twenty-four years, 1856-1879 inclusive. In the months of April, June and December, 1879, the average age of American decedents was less than that of decedents of foreign parentage;

this is indicated in the table by prefixing the — sign, in the column of "difference" to the figures for those months. In the other months of 1879, and in every month of the long term of years, the average age of American decedents was greater than that of foreign decedents.

Difference in years, between the average age of American and foreign decedents.

		1879.		24 years, 1856–1879.				
MORTHS.	Averag	e Age.	Difference.	Averag	Difference.			
	American.	Foreign.	Difference.	American.	Foreign.	Difference		
January	36.29	30.32	5.97	84.36	25.27	9.09		
February	85.54	28.72	6.82	83.93	22.49	11.44		
March	85.82	25.03	10.79	84.90	24.75	10.15		
April	33.49	85.86	-2.87	83.87	26.44	7.48		
Мау	40.81	25.18	15.13	86.54	25.74	10.80		
June	28.92	84.31	-5.39	84.30	24.68	9.67		
July	29.62	23.05	6.57	25.84	18.50	7.84		
August	29.13	25.40	3.78	27.45	18.42	9.03		
September	30.91	23.82	7.09	29.39	19.43	9.96		
October	29.03	19.78	9.30	32.69	22.62	10.07		
November	27.01	21.93	5.08	83.00	24.96	8.04		
December	24.05	24.22	-0.17	85.33	25.47	9.85		
Whole year.	31.77	26.22	5.55	82.47	22.85	9.62		

As a rule, the difference between the two classes of decedents is greater in the long period of years than in the single year. March and May are the only exceptions.

The largest monthly aggregate of deaths in 1879 (Table XIII.), was in January among the American decedents, and in December among those of foreign parentage; the smallest in September among the American, and in June among the foreign decedents.

For the twenty-four years, 1856-1879, the largest monthly aggregate among both classes of decedents was in August; the smallest in June. The percentages of the whole number of deaths in the different quarters for the period of twenty-four years, 1856-1879 inclusive, were as follows:

American parentage	1st quarter. 25.54	•	3d quarter. 27.48	4th quarter. 24.63
Foreign parentage	93.61	21.59	31.00	24.80

Table XV. gives the number of decedents, and the aggregate and average age, according to parentage, in each of the twenty-four years, 1856-1879 inclusive, and for the whole period:

TABLE XV. DEATHS, 1856-1879.

Recapitulation. Aggregate and average age by parentage.

>							<u>. </u>		
	Ameri	ican Pare	ntage	Foreign Parentage.			Total American and Foreign.		
YEARS.	Deaths.	Aggregate Tears.	Average Years.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregate I cars.	Average Years.
1856	541	14,554	26.90	524	8,958	15.75	1,085	22,812	21.49
1857	523	16,814	81.19	403	8,007	19.91	925	24,321	26.29
1858	519	15,916	31.07	505	8,478	16.78	1,017	24,388	23.98
1859	479	14,998	31.80	490	8,640	20.57	899	23,633	26.2 8
1800	512	16,660	83.54	489	9,751	19.94	1,001	26,411	26.88
1861	588	16,327	30.34	513	9,605	18.72	1,051	25,932	24.67
1862	460	14,589	81.71	454	10,994	24.21	914	25,583	28.00
1888	597	19,398	32.49	617	18,757	22.29	1,214	83.155	27.31
1864	649	21,782	33.4 8	633	11,865	17.98	1,281	33,097	25.83
1865	631	19,628	81,10	580	11,991	20.67	1,211	31,619	24.11
1886	513	18,906	86.85	523	12,762	24.40	1,036	81,668	80.57
1867	528	19,258	36.47	432	12,149	28.12	960	81,407	33.73
1868	596	21,779	36.54	514	12,539	24.89	1,110	34,318	80.92
1869	627	20,490	32.66	629	14,776	23.49	1,256	35, 26 6	28 08
1870	642	21,078	32.83	621	14,393	23.18	1,263	85,471	2 8.08
1871	643	22,725	85.84	611	17,005	27.83	1,954	89,730	81.69
1879	882	27,572	33.14	771	19,180	24.88	1,603	46,732	29.15
1878	862	28,675	88.27	857	19,061	22.24	1,719	47,736	27.77
1874	885	27,588	31.17	1,102	24,802	22.51	1,967	52,390	26.37
1875	914	30,419	88.28	1,001	24,802	24.28	1,915	54,721	28.57
1876	875	29,025	38.17	990	28,950	24.20	1,865	52,984	28.41
1877	908	28,858	81.22	1,030	25,410	24.67	1,988	58,763	27.74
1878	889	26,724	80.06	1,100	25,178	22.89	1,989	51,902	26.09
1879	941	29,896	81.77	1,085	28,446	26,22	2,096	68,342	28.80
24 years' totals	16,097	522,593	83.47	16,402	874,788	22.88	32,499	897,381	27 61

The greatest difference between the two classes was 15.50 years in 1864; the least difference, 5.55 years in 1879. The difference in the entire period is 9.62 years.

Number and percentage of decedents at different ages. Table XVI. shows the number and percentage of decedents, according to parentage, in each division of ages in 1879; also the same facts for the whole number of decedents in the twenty-four years, 1856-1879 inclusive:

TABLE XVI. DEATHS, 1879.

Percentage at different ages, according to parentage.

		American, Parentage 1879.		Foreign Parentage, 1879.		Total American and For- eign, 1879.		24 years, 1856 to 1879.	
Ages.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	
Under 1 year	140	14.88	176	16.22	316	15.60	6,328	19.47	
I and under 2	64	6.80	81	7.47	145	7.16	2,890	8.89	
2 and under 5	106	11.26	147	13.55	253	12.49	3,058	9.41	
Total under 5	310	32.94	404	37.24	714	35.25	12,276	37.77	
5 and under 10	86	9.14	89	8.20	175	8.64	1,676	5.16	
10 and under 15	22	2.34	21	1.94	43	2.12	672	2.07	
15 and under 20	15	1.59	33	3.04	48	2.36	1,087	3.34	
20 and under 30	79	8.40	114	10.51	193	9.53	3,272	10.07	
30 and under 40	74	7.86	90	8.29	164	8.10	2,950	9.08	
10 and under 50	53	5.63	70	6.45	123	6.07	2,470	7.60	
50 and under 60	60	6.38	92	8.48	152	7.50	2,333	7.18	
00 and under 70	95	10.10	77	7.10	172	8.49	2,290	7.05	
70 and under 80	83	8.82	66	6.08	149	7.35	2,143	6.59	
0 and under 90	59	6.27	24	2.21	83	4.10	1,105	3.40	
and over	5	.53	5	.46	10	.49	225	.69	
Total.	941	100.00	1,085	100.00	2,026	100.00	32.499	100.00	

The number and percentage of decedents under 5 years of age in 1879, were less in both classes than in 1878. Of American decedents, the number of this class in 1879 was 310, or 32.94 per cent. of all; this was 17 less than in 1878 and the per cent. 3.84

less. Of decedents of foreign parentage the number in 1879 was 404, or 37.24 per cent. of all; the number was 65 less and the per cent. 5.40 less than in 1878. The whole number of decedents under 5 years of age in 1879, was 714 or 35.25 per cent. of all; this was 82 less than in the previous year, and the per cent. 4.77 less. This accords with the fact shown by table XIII., that the average age of decedents in both classes in 1879, was higher than in 1878.

The following statement gives the percentage, according to parentage, of the decedents under 5 years old in each of the twenty-four years, 1856-1879 inclusive —

	American. Under five years.	Foreign. Under five years.	Total American and Foreign. Under five years.
856	37.15 per cent		nt
1857	32.31 per cent	51.74 per ce	nt40.75 per cent.
1858	35.35 per cent	55.44 per ce	nt
1859	28.60 per cent	48.34 per ce	nt37.82 per cent.
1860	32.81 per cent	48.87 per ce	nt
1861		52.63 per ce	nt44.05 per cent.
862	31.08 per cent	42.07 per ce	nt 36.54 per cent.
1863	28.47 per cent		nt
	STATE OF THE PARTY	A CONTRACTOR OF THE PROPERTY OF	nt39.82 per cent
	The state of the s	Control of the Contro	nt37.82 per cent
			nt32.24 per cent
	A CONTRACTOR OF THE PROPERTY O	TO US THE OWNER OF THE PARTY OF	nt29.59 per cent
	Contract of the Contract of th	and the same of the same of the	nt34.69 per cent
	The second secon	•	nt37.02 per cent
	Contract of the Contract of th	A STATE OF THE PARTY OF THE PAR	nt
	Action to the second se	Table 1 to the second second second	nt32.22 per cen
	and the second s	The second secon	nt35.27 per cen
			ont36.71 per cen
		A STATE OF THE PARTY OF THE PAR	nt39.31 per cea
	The state of the s	ATTACABLE STATES OF THE STATES	nt36.66 per cen
	The state of the s	A STATE OF THE PARTY OF THE PAR	
	THE RESERVE AND ADDRESS OF THE PERSON ADDRESS OF THE PERSON AND ADDRESS OF THE PERSON AND ADDRESS OF THE PERSON AND ADDRESS OF THE PERSON AND ADDRESS OF THE PERSON AND ADDRESS OF THE PERSON AND ADDRESS OF THE PERSON ADDRESS OF THE PERSON ADDRESS OF THE PERSON ADDRESS OF THE PERSON ADDRESS OF THE PERSON ADDRESS OF THE PERSON ADDRESS OF THE PERSON ADDRESS	O. O. C. Strategy and Strategy	ent38.98 per cen
			ent37.36 per cen
	Control of the Samuel of Samuel		nt40.02 per cem
1879	32.94 per cent	37.24 per ce	nt35.25 per cem

The percentage of decedents of this class in 1879 was smaller than in any year since 1871.

In the 40 years, 1840-1879 inclusive, the number of decedents in Providence was 45,644; of whom 18,011 were less than five years old; the number and percentage at each age under 5 years were as follows:—

Whole number of decedents	45,644
Under one year	. 8,833, or 19.35 per cent. of all.
One, and under two years	. 4,678, or 10.24 per cent. of all.
Two, and under five years	. 4,505, or 9.87 per cent. of all.
Total, under five years	.18.011, or 39.46 per cent. of all.

Table XVII. is a recapitulation, and shows the number and percentage of decedents at different ages, according to parentage, in the period of twenty-four years, 1856-1879:—

Table XVII. Deaths, 1856-1879.

Number and percentage at different ages according to parentage.

	American 24 years,	Parentage, 1856 to 1879.	Foreign I 24 years,	Parentage, 1856–1879.	American and Foreign Parentage, 24 years, 1856–1879.		
Age.	Number of Deaths.	Percentage in each divi- sion of age.	Number of Deaths.	Percentage in each divi- sion of age.	Number of Deaths.	Percentage in each divi- sion of age.	
Under one year.	2,778	17.26	8,550	21.64	6,328	19.47	
l and under 2	1,056	6.56	1,834	11.18	2,890	8.89	
2 and under 5	1,241	7.71	1,817	11.08	8,058	9.41	
Total under 5	5,075	81.58	7,201	43.90	12,276	87.77	
5 and under 10.	839	5.21	837	5.10	1,676	5.16	
10 and under 15.	86 3	2.26	809	1.89	672	2.07	
15 and under 20.	575	8.57	512	3.12	1,087	8 .84	
90 and under 30.	1,500	9.32	1,772	10.80	8,272	10.07	
30 and under 40.	1,392	8.65	1,558	9.50	2,950	9.08	
40 and under 50.	1,235	7.67	1,235	7.58	2,470	7.60	
50 and under 60.	1,311	8.14	1,022	6.23	2,333	7.18	
60 and under 70.	1,384	8.60	906	5.52	2,290	7.05	
70 and under 80.	1,441	8.95	702	4.29	2,143	6.59	
80 and under 90.	833	5.17	272	1.66	1,105	3.40	
90 and over	149	.93	76	.46	225	.69	
Totals	16,097	100.00	16,409	100.00	32,499	100.00	

From Tables XVI. and XVII. we see that of the decedents under five years of age, the number and proportion of those of foreign parentage are much larger than of those of American parentage. In the twenty-four years 1856–1879 inclusive; the number of decedents under five years of age were: American parentage, 31.53; foreign parentage, 43.90. This large proportion of young decedents among the foreign class explains the fact already shown in Table XIV., that the average age of decedents of this class is much less than that of American decedents. The average age of American decedents for the same period of twenty-four years, was 32.47 years; of foreign decedents, 22.85 years. During this period the proportion of American decedents living beyond the age of fifty years was 31.78 per cent. of all; while the proportion of foreign decedents of that age was only 18.16 per cent. of all.

Also by comparing Tables XVI. and XVII. we see that the percentage of decedents under five years of age in 1879, was 2.52 per cent. less than the average for the period of twenty-four years; the percentage of American decedents of this class in 1879, was 1.41 above the average; and the percentage of those of foreign parentage, 6.66 less than the average. This again corresponds with the facts of tables XIII. and XIV., where it may be seen that the average age of all decedents in 1879 was 1.19 years above the average for the period of twenty-four years; the average age of American decedents in 1879, being 0.70 years less than the average; and the average age of the foreign class being 3.37 years above the average for 24 years.

Deaths by age and parentage in each month and quarter of the year. In Table XVIII. will be found the number of deaths, according to parentage, in each division of ages, and the number of all ages in each month and quarter of the year; the table thus shows the comparative mortality among the two classes at different ages and seasons of the year. The whole number of deaths in 1879 was 2,026; of American parentage, 941; of foreign parentage, 1,085; the percentages of each in the different seasons of the year were as follows:—

	1st quarter.	2d quarter.	3d quarter.	4th quarter.
American parentage	28.80	20.61	24.02	26.57
Foreign parentage	22.39	22.58	24.62	30.41

In the remarks upon Table XII., were given the number and percentage of all deaths in each quarter of the year, and the largest percentage was in the fourth quarter; the next in the first quarter; the third quarter being third in order, instead of first as is more generally the case. Considering the deaths now in the two classes of parentage, we see that among the decedents of American parentage, the largest percentage was in the first quarter of the year; the next in the fourth quarter; among those of foreign parentage the largest percentage was in the fourth quarter; and the next in the third quarter.

The mortality at different ages and seasons may be seen from Table XVIII. The number of young children who died in 1879, was 57 less than in the previous year; of American parentage, 28 less; of foreign parentage, 29 less; as usual the largest percentage of decedents of this class was in the third quarter of the year, both among the American and foreign; of American parentage the number of children under one year old who died during the year was 140, of whom 57, or 40.71 per cent., died in the third quarter.

Of foreign parentage, the number of decedents under one year old was 176; of whom 79, or 44.89 per cent. died in the third quarter.

Causes of death; sex, age, parentage, and percentage from each cause. Table XIX. gives the number of deaths in 1879 from each cause, according to sex and parentage, and in each division of age; also the percentage from each cause of all from known causes.

Season and Disease. Table XX. gives the number of deaths from each cause in each month and quarter of the year, showing the comparative mortality from different causes at different seasons of the year.

TABLE XVIII. Deaths by age and parentage, in each month and quarter, 1879.

_	_				Div	1510	y al	. Pı	ESO	D6 0	T L	IFE.				, K
I	DIVISIONS OF THE YEAR.	Under 1.	1 to 2.	# to &	\$ to 10.	10 to 18.	16 to 20.	-	30 to 40.	40 to 50.	30 to 60.		70 to 80.	80 to 90.	90 to 100.	ALL AUKS
	January February March	17 15			6 3	1	1	7	15		5	10 10 8	7	9 6 13		10
	ist quarter	41	16	29	15	4	7	20			16		24	27	2	7
	April May June	9 4 5	, į	6	5	2	1	7 12 7		3	5 3 2	8	7			
	24 quarter	19		13	13	8	2		21	11	10	26	19	10		1
	July August Heptember	25 20 11	4	7 9 6		1 3	••••	5 3	•	5	8 8 5		, 9	.2 2 4	 I	
	3d quarter	57	- 1	22	18	4		16	11	15	21	17	25	8		
	October November December	9	7	10 12 21		 1 5	1 8 2	10 4		4 8 5	5 4	9 3 10	5. 6	4 6 4	1	
l	4th quarter	23	19	43	42	9	6	17	14	12	18	22	15	14	<u>i</u>	1
	Whole year	140	64	106	86	22	15	79	74	53	60	96	83	59	5	8
	January February March	18 0	2 3 8	11 12 13	10° 5	1	8	11 13 7	5	6 8 7	8 11 6	10 4 8	9 3 8	2 1	i	
l	1st quarter	80	13	85	20	-6	4	81	20	21	25	17	15	5	1	9
	April May June	18 11: 13		10 5	5 5 4	3 1 1	2 4 2	10 11		6 3 2	15 8 8	6 4 9	11 5	1	2	
1	2d quarter	87	17	21	- 14	5	-8	29	19	11	81	19	23	9	2	2
ı	July August September	81 25 22	4 5 5	5 8 9	6 2 4	2 3	4 3 2	18 8 9	14	5 5 8	4 5 5	5 9 5	8 6 4	1 1 2	····i	
	\$4 quarter	79	14	22	12	4	8	30	20	18	14	19	18	4	1	1
ı	October November December	10 11	15 10, 12,	25 21 23	17 17	1 5	2 6 5	19 10	9 7 6	18 7	4 5 18	4 7 11	4 2 4	 8 	1	1
	4th quarter	80	87	69	48	-6	18	24	22	25	22	22	10	6	1	1
I	Whole year	176	81	147	89	21	88	114	90	70'	92	77	86	21	5	1.0

Table No. XIX. Deaths, 1879. Disease, sex, age, parentage and percentage from each cause.

		SEX.				4	11118	IONA	DIVISIONAL PERIODS OF LIFE.	RIOD	8 or	Lir				P.	PARRNTAGE.	AGE.	-WD:	u.m.o
CAUSES OF DEATH.	Males.	Females.	.latoT	Under I Year.	I and I and under 2.	under 5.	under 10. 10 and	onder 15. Is and See 20.	20 and 20. 30.	30 and 30.	40 and under 50. 50 and	.00 тэрад Бая 00	.0% rebau bas 0%	.08 Taban bas 08	.00 rebau baa 00	over. American.	Foreign.	Total	Ratio of des from each to whole m	ber from kn. causea.
Accidents, (various)	12	00	8	100	1:	04	05	<u> </u>	1	07	-	 	1 :	-	<u> :</u>	 :	-	12	18	gi
" Asphyxla	63	:	00	01	÷	+	-	•	<u>:</u>	i	÷		<u>:</u>	\div	- <u>÷</u>		6 9	_	83	31.
" Burns and Scalds	æ	•	a		-	03	÷	<u>:</u>	64	:	=	$\frac{\cdot}{\cdot}$	<u>:</u>	+	÷	•	-65		<u>a</u>	ś
" Drowning		:	60	:	÷	:		:		*	-	=	÷	\div	÷		-	6	66	ŝ
" Falls	_	**	7	i	:	:	<u>:</u>	<u>:</u>	:	÷	:	60	61	_	=	:		8	-	矣
" Poisoning	64	69	*	÷	:	01	<u>:</u>	:	<u>:</u>	÷	÷	$\overline{}$	-	:	÷		61	81	4	8
" Ballroad	-	:	-6	i	- <u>÷</u>	;	<u>:</u>			=	oi		\div	<u>-</u> :-	<u>:</u>	— <u>;</u> -		4	-6	켨
A Decesses		•		=	÷	÷	÷		8	:	\vdots	\div	:	÷	$\frac{\cdot}{\cdot}$			_	10	Si
Ansemis	<u>:</u>	- 	69	i	:	-	:	<u>:</u>	-	:	÷	:	÷	\div	÷	•			GA .	9.
Aneurism	_	i		i	\vdots	÷	÷	<u>:</u>	<u>:</u>		- i	$\dot{}$	÷	\div	÷	<u>:</u>		_		ફ
Aorta, Disease of	_	:	_	i	÷	:	÷	<u>:</u>		÷	•	÷	÷	÷	<u>:</u>		<u>.</u>		_	ŝ
Apoplexy	8	\$	28	÷	÷	:	<u>:</u>	<u>:</u>	*	4	10	Ξ	6	71	6		88	88	28	2.7
Asthma	i	61	61	÷	\div	<u>:</u> -	÷	<u>:</u>	:	:	÷	:	- 64	÷	÷	<u>:</u>	<u>.</u>	69	69	e.
Bladder, Diseases of		-	90	÷	÷	-	÷	<u>:</u>	:	:	69	-	_	- ; -	÷	<u>.</u>	- 69		60	ş
Bones, Diseases of	_	_	61	÷	:	:	÷	<u>:</u>	:	-	÷	÷	.	- : -	÷	- :		—	69	Š.
Brain, Congestion of	·	7	13	_	7	8	-	÷	69	91	-i	÷	<u>:</u>	<u>-</u> :-	÷	<u>.</u>	10	-	81	ş
" Diseases of	_ _	15	83	69	_	<u>:</u>	_		-	8	_	4	F	- ;	_	_	<u> 66</u>	-	- 18	1.14

TABLE NO. XIX.—CONTINUED.	1	DEATHS, 1879.	187	٠.	S.	80	8	8 8	8	pare	a !	8	Pag (ξ,	\$	2007	Disease, sea, age, parentage and percentage from each	96	\ <u>\text{8}</u>	CHASE.
		SEX.	<u> </u>	! :		֡֞֞֞֞֞֞֞֞֞֞֞֞֞֓֓֓֞֞֞֞֞֞֞֞֓֓֓֡֞֜֞֡	ž	DIVISIONAL PERSONS OF LIFE.	<u> </u>	. Š	à					<u> </u>	PARKNTAU	L'AUM.	-ग्यूज	TERIO 1 TRANS
CAUSES OF DEATH.	Males.	Females.	Total.	Teatbas t	S s-sbau bas 2	dander 5. Dan 3. Ol 19dar	bas 01 Al rebas	ban či under 20.	So and ander 30.	bas 96 .02 19bas	bas 98 .08 mbus bas 98	.00 T-buss bass 40	Of when bas Vi	.08 raban bas 08	Das VV	American.		Total	Estiv of des	COMMENT POLICIANT FOR EAST COUNTY OF
Brain, Inflammation of	2	क्ष	=	12		1 20		1:	1:	ΪĒ	 	╁╾	100	:		<u> </u>	1 =	3	<u>.</u>	3
Bronchitis	3	81	3	2	-	-		:		<u>:</u>	:	:	-	•	4	.	*	£	1	=
Cancer, (various)	2	2	\$		÷	<u>:</u>	:		:	4	8	•	•		~		-	=	5	=
" of Breast	:	-	66	:	<u>.</u>	÷	<u>:</u>	:	:	7	-	-	=	:	÷	-	4	ट	-	÷
" of Stomach	•	66	Ė	-	÷	<u>:</u>	<u>:</u>	:	:	64	-	-	-		$\frac{\cdot}{\cdot}$	- :	t -	2	<u></u>	I
of Uterus		*	2	÷	<u>:</u>	<u>:</u>	•	:	:	*	4	•	÷	<u>:</u>			€	=	=	ş
Cancrum Orls	:		<u> </u>	$\frac{\cdot}{\cdot}$		-	<u>:</u>	:	÷	_:	÷	$\frac{\cdot}{\cdot}$	\div	<u>:</u>	<u>:</u>		-	-:	-	3
Catarrh		_	-	-	:	<u>:</u>	_ :	:	<u>:</u>	- :-	:	<u>:</u>	<u>:</u>		<u>:</u>	<u>.</u>	<u>:</u>	-:	-	S
Cellulitis Pelvic	i		=	<u>:</u>	<u>:</u>	<u>:</u>	:			:	÷	÷	:	:	:	-	-:	-	_	3
Childbirth		.	_ <u>.</u> -	-:	<u>:</u>	<u>:</u>	:	:	-	_ _	÷	÷	:	÷	<u>:</u>	<u> </u>	4	-	5	ş
" Puerperal Convulsions		-6	-6	:	<u>:</u>		_:	_	45	:-	÷			-	<u>.</u>	_			4	3
" Puerperal Fover	:	-	6	:	<u>:</u>	<u>:</u>	_ :	_ :	•	-		$\frac{\cdot}{\cdot}$	<u>:</u>	- :	<u>:</u>		•		•	Ħ
Cholera Infantum	8	8	2	8	-6-		_:_		÷	÷	:	÷	÷	<u>:</u>	<u>:</u>		3	=	æ	2
" Morbus	_	-	40	-	<u>:</u>			:	-	÷	:	-	_	_	<u>:</u>	-	-	4	4	ş
Colfe	:	a	60	=	<u>:</u>	<u>:</u>	_	İ	Ė	÷	÷	\div	:		<u>:</u>		:	-:	-	.16
Consumption	171	158	ğ	_	-	_	후	22	8	8	Z	£	2	-	<u> </u>		=	=	8	14.00
Convulsions	5	8	28	-	_	_	-	:	⋮	-	-		∹	_ <u>:</u>	_:	- <u>.</u>	_	-	2	8.8

each cause.	-10 D	Ratio of de from each c to whole u ber from kn causes.	8.13	1.14	97	8	1.49	98	5.80	8 ;	4.	£.	8	9	.15	1.98	4.	8	6.31
	. 1	Total.	3	88	2	4	2	-	101	4	6	9	•	2	8	\$	90	_	300
Disease, sex, aye, parenings, and percentage from	PARKNTAGE.	Foreign.	18	16	65	**	18	:	23	8	•	80	•	-	64	7	64	:	3
lage.	PAB	American.	83	-	۲	-	25	-i-	\$	_	65	65	A	65		8	ά	7	8
cen		90 and over.	1	•	•	:	:	:	:	7	:	•	:	:	:	:	:	:	
ž.	٠	.08 and .08	:	:			:	:			:	•	:	i		<u>:</u>	01	:	<u> </u>
nd		70 and .05	:	*	:		-	1		•	_	:	:	_	_	61	*	:	2
8, 0	LIFE.	.05 as 08 .05 19bau	1	@1			_	:	<u>:</u>		OR	69			<u>:</u>	_	~	<u>:</u>	ਡ
jug.	OF L	50 and 00.	<u>:</u>	91		_	_	:		01			_				<u>:</u>		8
re	28	- bna 04 .05 19bnu	<u> </u>	_	<u>:</u>	<u>:</u>		<u>:</u>	:			_	:	<u>:</u>	<u>:</u>	_	<u>:</u>	:	72
ď	DIVISIONAL PERIODS	30 and .04 rebau	<u> </u>	•			_		:	<u>:</u>	_		O4	_	<u>:</u>	<u> </u>	<u>:</u>	<u>:</u>	- -
rije,	. P.	20 and 02.	Ŀ	_	:	_	:	i	•		_			<u>:</u>	:	=	<u>:</u>	<u>:</u>	
8	OMA	da and st. 30.	<u> </u>			<u>:</u>			83	<u>:</u>	<u>:</u>	_	_		<u>:</u>	4	<u>:</u>	<u>:</u>	
	V181	bas 01 .31 19bau	_	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>		<u>:</u>	<u>:</u>		<u>:</u>		<u>:</u>	•	<u>:</u>		_
ass,	DI	5 and under 10.	R		:	:	<u>.</u>	<u>:</u>	8	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	_	<u>:</u>	-0	<u>:</u>	<u>:</u>	•
Stae		2 and 3. Tabau	12	_	<u>:</u>	<u>:</u>	:	<u> </u>	3	<u>:</u>	<u>:</u>	:	Ė	_	:	:	<u>:</u>	:	<u> </u>
T		bas l Laban			<u>:</u>	<u>:</u>	_	<u>:</u>	ह	i	_	10	<u>:</u>	<u>:</u>	<u>:</u>		:	:	-
79.		Under 1 Year.	_	81	10		91		-	<u>:</u>			<u>:</u>	•	09		:	:	
, 18		Total.	3	8	21	•	8	_	107	_	.	16	•	2	∞	\$	α0	_	100
DEATHS, 1879.	BEX.	Females.	11	15	•	OR	14		6	•	•	۴	60	•	•	#		-	28
DE		Males.	8	60		a	18	i	\$		10	a	65	4	1	19	7	:	8
TABLE NO. XIX.—CONTINUED.		CAUSES OF DEATH.	Croup	Debility	" Premature Birth	Diabetes	Diarrhos	" Chronic	Diphtheria	Dropsy	Dysentery	Entertite	Epilepsy	Erysipelas	Exposure and Neglect	Fever, Typhold	Gangrene, Senile	Glottle, Spasm of	Heart, Diseases of

Ratio of deaths
from each cause
to whole number from known 8 4 6 8 8 8 8 5 2 3 Total. PARKNTAGE. Foreign. American bna 00 ban 08 .09 19bau han 07 .08 тэһап 60 and 05 Tebra LIFE. under 60. ban 00 OF 20 and 30 and 30 and 10 and 10 and 10 and 10 and 10 and PKRIOD8 DIVISIONAL under 20. bas di at rebau **ban** 01 6 and under 10, g Japan yan 2 under 2. bna 1 Under 1 Year. Total. Lemulca. Males. Leucocythæmia Heart, Hypertrophy of..... Hip Joint, Diseases of...... Intemperance Hemorrhage Hooping Cough..... Insanity Laryngitis..... from Lungs..... Delirium Tremens Kidneys, Diseases of...... Hernia OF DEATH. Hydrocephalus..... Icterus Neonatorum..... Bright's Disease of from Stomach. Influenza CAUSES :

Disease, sex, age, parentage and percentage from each cause. DEATHS, 1879. TABLE NO. XIX.-CONTINUED.

cause.
e from each c
percentag
ser men, age, parentage and
DEATHS, 1013. Discuss, week, uge,
ABLE No. MINI-
TABLE No.

		SEX.				-	IVE	DIVISIONAL	LP	PERIODS	40 St	F LI	LIFE.				A	PARENTAGE	NOB.	9sm
GAUSES OF DEATH.	Males,	Females,	Total.	Year.	S and S and	dader 5. band	10 and Il	15 and under 15. under 20.	20 and under 30.	30 and under 40.	40 and of under 50.	50 and under 60.	60 and under 70,	70 and under 80.	80 and .00 rebnu	90 and over.	American.	Foreign.	Total	Ratio of des from each ca to whole m ber from kno causes.
Liver, Diseases of	13	to	8	1:	1:	1:	H	1:	-	104	18	00	7	100	01	1	1	13	80	66.
" Inflammation of		1	-	:	+	+	÷		:	:	:		1	-	1	:	-	1		90.
Lungs, Diseases of	*	-	4	-	:	-:				-	1	1	0.5	1	1	i	60			1.20
Malformations	6	01	11	6.	O.	÷	- [:	1	:	1	į	:	1	:	1	•		-	19.
Marasmus	13	15	88	19	10			:			-	:	1	1	i	i	1	11	O.	1.39
Meningitis, Cerebro-Spinal	9	1	-	÷	÷	4	-	•	:	1		ì	:	i	1	1	_	~		28.
Murder and Homicide	F	-	F	1	:	:	1	:	1	1	:	-	:	:	i	:		-		-
Old Age	22	37	. 19	:	:	:	-	:	:	1	1	:	1	8	33	0	*	64	9	3.17
Paralysis	90	10	30	÷	:	:	-	1	1		*	4	7	6	10		80	ĭ	85	30 1.49
Peritonitis	9	00	7	:	:	:	-	_	-	-	0)	:	01	1	1	1	~	_		697
Pharyngitis	F	:	-	ř	-	÷	-	:			-	i	1	:	1	:	_	1		10.05
Phlegmasia Dolens	1	-	-	i	-	:	-	1	_		i		1	•	:	1	į	_	+	10.05
Pleurisy	-	60	*	Ť	1	:	-	:	-		:	-	ÇS	1	:	1	1			.20
Pneumonia	69	92	145	124	16	21	-		9	60	15	19	15	15	60		64	8	_	45 7.19
" Congestion of Lungs	-	1-	14	9	÷	÷	+	1		-	61	1	:	0.5	01	1			-	.69
Prostate, Disease of	-	:	1	:	:	:	:	-		:	i	1	1	1	-			1		1 .05
Pymmia		.04	69	i	÷	:	-	:	-	1	i	i	1		:	1		_	01	01.
Rhenmatism	50	×	11			9	-						-							1

Ratio of deaths from each cause ber from known causes. 100.00 19.40 S 2 ġ Deaths, 1879. Disease, sex, age, parentage and percentage from each cause. 20.0 .latoT PARENTAGE. 8 Foreign. Ē 3 American. under 80. 80 and udder 90. 90 and over. 2 8 : 2 baa us रता <u>क्या</u> स्था .07 19bnu LIPE. baa 08 ban 08 .00 1∍ban DIVISIONAL PERIODS OF od 19bau 40 under 50. of Tabun pun og ğ under 30. pau oz **\$** under 20. ban 61 3 10 and In. 13. band under 10. 8 3 under b pun 2 3 8 under 2. pun t 2.036 316 Year. Under 1 .faioT 8 SKX. Females. 8 Males. TABLE No. XIX.—CONTINUED. Tabes Mesenterica..... Tecthing Tetanus and Trismus Nascentlum..... Urinary Organs, Disease of...... Soarlatina Spine, Diseases of...... Stomach, Diseases of...... Sulcide Syphilis, Congenital..... Thrombosis..... Tamor Uraemia...... Inflammation of...... Scrofula..... Tabercalosis..... CAUSES OF DEATH. Skin, Diseases of...... Unknown. Totals

	TABLE	XX.		DEATHS, 1879.	18, 1	879.	Sea	Season and Diseass.	nd	Disea	.96						
Caubes of Death.	·Transat	February.	March.	First Quarter.	April.	.ysM	упле.	Second Quarter.	July.	August.	September.	Third Quarter.	October.	.тэфшэхоИ	December.	Fourth Quarter.	Whole Year.
Accidents, (all kinds)	85	-	8	2	*	8	-	2	1-	-	8	2	*	2	•	æ	8
Apoplexy	-	æ	61	=	۲	60	•	a	8	61	٦	15	Ø1	64	-6	a	28
Brain, Diseases of	OI.	-	60	8	٦	10		14	4	4	Ø	07	2	-	-	ä	76
Bronohitis	•	•	10	16	4	69	**	6	:	61		*	:	4	60	7	z
Cancer, (all kinds)	_	94	*	a	4	6 6	60	8	1-	-	16	19	10	76	10	18	8
Childbirth	*	_		10	÷	:		61	8	7	н	i	:	4	4	a	8
Cholera Infantum		i	÷	:	- <u>:</u>	:	-	~	2	81	•	6	÷	i	:	_	2
Cholera Morbus		i	i	-	<u> </u>	-	<u>:</u>	:	-	8		ö	:	i		:	ю
Consumption	8	12	8	28	8	8	7	8	3	8	8	8	8	81	8	67	888
Convulsions	æ	6	65	11	۵-	-	61	Ξ	80	=	10	*		П	10	-	8
Croup	a	•	80	83	4	94	69	*	<u>:</u>		-	<u>:</u>	4	7	69	82	\$
Diarrhoa	_	÷	:	<i>9</i> 4		-	4	6	<u>a</u>	7	10	12	:	01	:	09	81
Diphtheria	2	-	65	8	16	8	9	8	ю	7	0 0	8	81	82	0	\$	101
Dysentery	:	i	:	:	-	÷	:	61	•	69	-	•	÷	i	:	_	.
Enteritis	:	-	:	_	64		F	4	4	લ		-	oi.		æ	4	16
Erysipelas		:	ðī	60	8 1	-i	:	•	÷	:	:	П	-	F	_	60	10
Fever, Typhoid	_	91	10	60	81	-	6	2	80	69	-	6	A	ON.	۲	Ħ	\$
Heart, Diseases of	=	F	2	88	2	=	₹	8	80	81	-60	8	20	Ξ	65	ब्र	н

TABLE XX.—CONTINUED. DEATHS, 1879. Season and Disease.

CAUSES OF DEATH.	January.	February.	March.	First Quarter.	April.	May.	Aune.	Second Quarter,	July.	August.	September	Third Quarter.	October.	Мочетрет	December	Fourth Quarter.	Whole Year.
Hooping Cough	24	-	04	10	-	-	O.S	7	***	01	-	01	Ot.	-	1	7	15
Hydrocephalus	C)	*	9	11	10	69	co	10	60	*	1	90	-	4	C4	9	35
nsanity	O1	-	G4	শা		4	-	4	-	1	64	00	-	1	-	-	12
ntemperance and Delirium Tremens	-	7		04	-	1	-	01	-			1	-	:	03	0)	-
Kidneys, Diseases of	90	4	4	16	4	7	80	11	4	04	00	8	60	01	01	-	48
Liver, Diseases of	1	-	00	10	00	-	00	1-	-	01	1	60	CI	04	01	9	12
Marasmus	1	01	-	00	10	O.s.	C3	6	20	01	co	10	01	00	-	9	88
Meningitis, Cerebro-Spinal	-	1	1	01	-		1	61	01		1	60		-			-
Old Age	111	00	t-	160	1-	g1	0	14	20	-	9	18	-	9	-	14	72
Paralysis	-	4		*	80	0.5	-	9	60	5	4	12	01	24	4	90	30
Peritonitis	-	-	64	60	C4	61	1	10	G#	1		20	04	1	-	00	14
Pneumonia and Congestion of Lungs	38	18	18	7.4	18	11	10	36	1	4	9	11	00	6	21	38	159
Scarlatina	13	12	6	34	0	10	7	233	20	10	19	34	36	10	T	161	252
Suicide	1	-	-	1	-	-	-	60						1	1	CI	9
Ceething	-	-	1	1	-	-			4	CN	1	-	7	-	1,4	. 5	13
Inknown	01		-	OI.	01	-		25	-	01		20	*****	******	1	1	0
Other causes as given	17	10	10	48	18	93	14	8	27	18	-	20	15	30	15	50	212
Totals	900	151	163	514	147	150	VIII	100	1	P	1	1		1		1	1

NOTES ON THE CAUSES OF DEATH,

DURING THE YEAR 1879.

In these notes attention is called to some of the facts given in Tables XIX. and XX., and fuller explanation made of them. There are also given additional statistics in regard to many of the causes of death; and the facts are compared with the statistics for the twenty-four years, 1856–1879 inclusive.

The percentage of deaths from the different causes is always reckoned on the number from known causes.

There were 2,026 deaths in Providence during the year 1879; of 9 of which the cause was not stated, this was 0.44 per cent. of the whole number; it was 7 less than in 1878, and the per cent., 0.36 less. The number from known causes, and on which, therefore, all the percentages are reckoned, was 2,017. During the twenty-four years, 1856–1879, the whole number of deaths was 32,499, of which 31,966 were from known causes; and 533 or 1.64 per cent. were from unknown causes. In the forty years, 1840–1879 inclusive, the number of deaths in Providence, was 45,644; of these 44,109 were from known causes; and 1,535 or 3.36 per cent. from unknown causes.

From Table XX., which gives the number of deaths from different causes in each month and quarter of the year, we may see the comparative prevalence of different diseases in 1879. Of the deaths from bronchitis, croup and pneumonia a large number occurred in the first quarter of the year; of the deaths from diphtheria, 56 per cent. were in the third and fourth quarters; while of the deaths from scarlatina, 161, or 63.9 per cent. were in the fourth quarter.

Abscesses. There were 5 deaths reported from abscesses in 1879, or 0.25 per cent. of all from known causes; this is the same number and per cent. as in 1878. One was abscess of the pelvis;

one, abdominal; one, pharyngeal; and one retro-pharyngeal. The decedents were, 1 male and 4 females; all of American parentage; in 1878, the 5 decedents were all of foreign parentage. The following are the statistics for the 24 years, 1856-1879 inclusive:—

Number of deaths from all known causes, 31,966.

From abscesses: males, 48; females, 43; total, 91, or 0.28 per cent, of all American parentage, 41; foreign parentage, 50.

Accidents. The number of deaths reported as the result of accidents of different kinds, was 56; 6 more than in 1878, and 2.78 per cent. of all from known causes. Of these deaths, the table specifies 3 from asphyxia; 9 from burns and scalds; 8 from drowning; 7 from falls; 4 from poisoning, and 5 from railroad accidents; there were also 20 deaths from "various" accidents not named in the table, of which some were as follows: two were "injured at birth;" one was killed while "coasting;" one by the "kick of a horse;" one by the "falling of lumber;" one by "swallowing a shawl-pin;" one "caught in gearing;" one "hit by a ball bat;" one "run over by a horse;" one died from a "pistol-shot in the abdomen;" one from a "penetrating wound of the eye;" one from "fracture of the leg;" one from "injury to the ankle-joint," and one of "rupture of the intestines."

Of the deaths from "burns," the only one which was described more definitely, was from the explosion of a kerosene lamp. Of the four deaths by poisoning, one was from a long-continued habit of opium eating; one from an over-dose of opium drops as a cough medicine; and two were of children three years old; one of whom was poisoned by eating orange peel, and the other by whiskey, given her by her mother.

The statistics as to sex and parentage are as follows :-

Males, 37; females, 19; total, 56, or 2.78 per cent. of all.
American parentage, 19; foreign parentage, 37.

The statistics for the twenty-four years, 1856-1879 :-

Males, 692; females, 229; total, 921, or 2.88 per cent. of all.

Aneurism. There was one foreign male who died from the cause in 1879. The statistics for the twenty-four years, 1856-18 are:

Males, 21; females, 3; total, 24, or 0.08 per cent. of all.

American parentage, 8; foreign parentage, 16.

Apoplexy. The number of deaths reported from apoplexy in 1879, was 56, or 2.78 per cent. of all; this was 14 more deaths and 0.65 per cent. more than in 1878. The number of males was 32; of females, 24; American parentage, 28; foreign parentage, 28. (See "Paralysis.")

The following statistics for apoplexy are for the twenty-four years, 1856-1879:—

Males, 333; females, 343; total, 676, or 2.11 per cent. of all.

American parentage, 474; foreign parentage, 202.

Brain, Diseases of. There were 76 deaths from diseases of the brain in 1879, including 12 from "congestion," and 41 from "inflammation." The percentage of all from known causes was 3.77; the number was 9 more, and the per cent. 0.37 more than in the year previous.

The following are the statistics of sex and parentage:-

Males, 35; females, 41; total, 76, or 3.77 per cent. of all.
American parentage, 41; foreign parentage, 35.

The statistics for the twenty-four years, 1856-1879:—

Males, 673; females, 567; total, 1,240, or 3.88 per cent. of all.

American parentage, 654: foreign parentage, 586.

Bronchitis. The number of deaths from this cause in 1879, was 34, or 1.68 per cent. of all from known causes; this was 13 less, and 0.70 per cent. less than in 1878. The particulars are as follows:—

Males, 12; females, 22; total, 34, or 1.68 per cent. of all.

American parentage, 18; foreign parentage, 16.

The statistics for the twenty-four years, 1856-1879 are:-

Males, 172; females, 231; total, 403, or 1.26 per cent. of all.

American parentage, 182; foreign parentage, 221.

Cancers. There were 66 deaths from cancers in 1879; this was 18 more than in 1878, and 8.27 per cent. of all. The decedents were: males, 21; females, 45; American parentage, 25; foreign, 41. Of these, 8 had cancer of the breast; 17, cancer of the stomach; and 14, cancer of the uterus. Of the 27 reported as "various," the following were specified: liver, 5; rectum, 8; face, tongue and bowels, 2 each; and one each of the æsophagus, abdomen, neck, orbit of the eye, pelvis, penis, and lymphatics.

The statistics for the twenty-four years, 1856-1879, are:-

Males, 200; females, 561; total, 761, or 2.88 per cent. of all.

American parentage, 486; foreign parentage, 275.

Child-birth. The number of deaths from child-birth in 1879, was 20, which was one less than in 1878; this number was about one per cent. of all the deaths, and 0.80 per cent. of the 2,499 mothers bearing children in 1879. Five of these deaths were from puerperal convulsions, and six from puerperal fever; 11 of the decedents were of American parentage; and 9 of foreign parentage. For the twenty-four years, 1856–1879, the statistics are:—

American parentage, 161; foreign parentage, 218.

Total, 379, or 1.19 per cent. of all deaths from known causes; and 0.88 per cent. of the 45,842 mothers who have borne children during that time.

Cholera Infuntum. This was reported as the cause of 59 deaths in 1879, or 2.92 per cent. of all. This was 11 less than in 1878, and the per cent. 0.68 less.

Males, 30; females, 29; total, 59, or 2.92 per cent. of all.
American parentage, 28; foreign parentage, 31.

The following table gives the statistics for each of the 24 years, 1856-1879, and for the whole period:—

	SEC	K.	Parentage	. '	Whole
	Males.	Females.	American. Fo	oreign. N	umber. Percentage
856	85	23	16	41	575.54
857	27	23	26	24	505.56
i 858	\$7	23	18	86	494.93
.659	14	19	18	90	88
:8 00	85	88	96	43	687.09
361	29	82	25	36	. 616.03
	25	19	17	27	. 444.98
	36	80	19	47	. 665.56
864	29	29	18	40	. 584.68
.865	20	80	13	87	. 504.91
886	80	17	15	82	474.64
	21	95	98	26	. 495.19
86 8	88	82	22	48	. 706.44
	35	80	27	38	. 655.34
870	48	45	87	56	. 987.44
871	80	32	26	36	625.00
873	73	79	58	98	. 1519.56
878	43	80	89	58	. 925.41
874	72	62	44	90	. 1846.89
875	62	74	50	86	. 1367.16
876	66	56	48	79	. 1226.59
877	70	52	48	79	. 122 6.81
878	85	35	26	44	. 708.55
879	80	29		81	. 592.92

The number for 1879, is less than in any year since 1867, and the percentage the lowest recorded.

Consumption. The number of deaths from this cause in 1879, was 293, or 14.52 per cent. of all; this number was 12 less, and the per cent. 0.94 less than in 1878. The decedents were divided as follows in regard to sex and parentage:—

American parentage males,	58females,	58Total, 116
Foreign parentagemales,	83females,	94 Total, 177
Totalmales,	141females,	152Total, 293

The statistics for the twenty-four years, 1856-1879, are as follows:—

	mber of deaths from known causes, 31,966; deaths from consumption, 16.79 per cent. of all. Annual average, 224.
	parentage males, 1,114females, 1,450 Total, 2,56
Foreign p	arentagemales, 1,323females, 1,481Total, 2.80
Total	
Males,	45.40 per centfemales, 54.60 per cent.
Ameri	can parentage, 47.76 per centforeign parentage, 52.24 per cent.

Having estimated the population of the city in the middle of the year 1879, at 103,000, the death-rate from consumption would be one in 351 of the inhabitants, or 2.84 deaths in each thousand persons. The average annual rate for the fifteen years, 1840–1854 inclusive, was one in 209.9, and for the twenty-five years, 1855–1879, it was one in 304.6.

By periods of five years from 1840 to 1879 inclusive, and for the whole period of forty years, the whole number of deaths from known causes in Providence, and the number and percentage from consumption, were as follows:—

	Number of deaths from known causes.	Number from Consumption.	Percentage of deaths from Consumption
1840-1844	2,824	A STATE OF THE PARTY OF THE PAR	And the second second
1845-1849	3,893	761	19.55
1850-1854	4,465	958	
1855-1859	4,753	930	19.57
1860-1864	5,298	997	18.82
1865-1869	5,474	1,009	18.43
1870-1874	7,784		
1875-1879	9,668		15 26
1840-1879, 40 year	s44,109	8,049	18.25

The number and percentage of deaths in each of the last ten years have been:—

		Per cent
1870 .	. · · · · · · · · · · · 238 · · · · · · · · · · · · · · · · · · ·	
1871	194	15.66
1872	241	15.26
1873		
1874	270	18.76
1875	299	
1876	284	
1877		15.22
1878	305	15.46
1879	293	14 59

The number and percentage of the decedents from consumption in each division of age, were as follows for the year 1879, and for the period of 24 years, 1856 to 1879 inclusive:—

Periods.	Under 1	and	under 2.	2 and under 5.	6 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 60.	50 and under 60.	60 and under 70.	70 and under 80.	80 and under 90.	TOTAL.
1879. Number		1	8	8	4	10	23	96	65	84	26	200	7	1	295
Per cent	.8	41	.02	1.02	1.87	8.41	7.85	82.77	22.19	11.60	8.87	6.83	2.89	.84	100.00
1856–1879. Number	6	4	63	57	57	109	490	1,652	1,185	748	501	846	134	19	5,869
Per cent	1.1	91	. 17	1.08	1.06	2.08	9.18	80.78	21.14	18.94	9.83	6.45	2.50	.22	100.00

Of the 293 decedents from consumption, in 1879, 66.56 per cent. were between twenty and fifty years of age; 52.22 per cent. lived beyond the age of thirty. The average age of decedents from all causes in 1879, was 28.80 years. Of the 5,368 decedents in 24 years, 65.86 per cent. were between the ages of twenty and fifty; 53.58 per cent. lived beyond the age of thirty.

Convulsions. There were 59 deaths reported from convulsions 1879, or 2.92 per cent. of all, and 3 more than in 1878. Two 'the decedents were between 5 and 10 years old; the others all sthan five years.



The statistics of sex and parentage are :-

Males, 33; females, 26; total, 59, or 2.92 per cent. of all.
American parentage, 22; foreign parentage, 37.

For the 24 years, 1856-1879 :-

Males, 471; females, 377; total, 848, or 2.65 per cent. of all. American parentage, 348; foreign parentage, 500.

Croup. This was the cause of 43 deaths, in 1879; 5 more than in 1878; 22 of these deaths occurred in the first quarter of the year. Four of the decedents were under 1 year of age; 9, between 1 and 2 years; 19, between 2 and 5 years; 10, between 5 and 10; and 1, between 10 and 15.

The statistics of sex and parentage for the year are:-

Males, 26; females, 17; total 43, or 2 13 per cent. of all. American parentage, 22; foreign parentage, 21.

For the twenty-four years, 1856-1879, the statistics are :-

Whole number of deaths, 739; annual average, 31.

Males, 376; females, 363; total, 739, or 2.31 per cent. of all.

American parentage, 307; foreign parentage, 432.

Debility. This was returned as the cause of 33 deaths in 1879, including 10 from premature birth. The particulars as to sex and parentage are as follows:—

Males, 14; females, 19; total, 33, or 1.63 per cent. of all.

American parentage, 14; foreign parentage, 19.

For the 24 years, 1856-1879 :-

Whole number of deaths, 573; annual average, 24.

Males, 301; females, 272; total, 573, or 1.79 per cent. of all.

American parentage, 238; foreign parentage, 335.

Diarrhea and Dysentery. The number of deaths from these causes in 1879, was 40, which was one more than in 1878. The number from each was as follows:—

Acute Diarrhœa				-	Total. Percentage.
Chronic Diarrhœa		1	1		1
Dysentery	5	4	8	6	9
Total	 21	_ 19		_ 24	401.99

The following are the statistics of deaths from other diseases of the same class, in the year 1879:—

Cholera Infautum	Males30			•	Total. Percentage
Cholera Morbus	1	4	····-	5	525
Enteritis	9	7	8	8	16
Total	 40	40	— 36	 4	80 8.96

Combining the figures for the five diseases, we find the total, as follows:—

Males, 61; females, 59; total, 120, or 5.95 per cent. of all.

American parentage, 52; foreign parentage, 68.

The number of deaths is 10 less, and the per cent. 0.64 less than in 1878; of males, 1 less; of females, 9 less; and 5 less in each class of parentage.

The statistics of the five diseases for the twenty-four years, 1856-1879, are as follows:—

Number of deaths, 3,615; annual average, 151.

Males, 1,811; females, 1,804; total, 3,615, or 11.31 per cent. of all.

American parentage, 1,468; foreign parentage, 2,147.

Of these decedents, the proportion of those of foreign parentage to those of American parentage in 1879, was 131 to 100; or 56.67 of foreign and 43.33 of American parentage in each 100. Of all decedents in 1879, the proportion was 115 of foreign to 100 of American parentage, or 53.55 foreign, and 46.45 American in each 100. The proportion of the foreign class is considerably larger in this particular class of diseases than in the number of decedents from all causes. A large number of the decedents from these diseases are young children; 75.83 per cent. of those in 1879 being

less than two years old. Of the 2,522 children born in Providence in 1879, 51.31 per cent. were of foreign parentage, and 49.69 per cent. of American parentage; or 105 children of foreign to each 100 of American parentage.

The same proportions are true in regard to the figures for the period of 24 years, 1856-1879 inclusive.

The following table shows the number of deaths from each of these causes, with the number from the whole class of causes, and their percentage of all from known causes, reported in Providence during the period from 1840 to 1854 inclusive; during each year, from 1855 to 1879 inclusive, and in the aggregate during the whole period of 40 years, 1840 to 1879 inclusive:—

,	Cholera Cl	holone			All of all from
		0.3.000,00	ambaa Da	contour Entantile C	All of all from ombined. known cause
1840-1854		. 63			.1,87016.7
1855			37	38	The state of the s
1856	. 57	5	29	35 6	. 13211.15
1857	. 50	2	38	29 5	. 12413.40
1858	. 49	2	27	12 8	98 9.60
1859	. 33	1	28	13 4	
1860	. 68	2	21	18 9	. 11811.78
1861	. 61	5	22	32 6	12611.90
1862	. 44	1	31	22 9	10711.70
1863	. 66	5	35	77 9	192 15.81
1864	. 58	4	45	40 13	16012.49
1865	. 50	7	34	62 4	15712.96
1866	. 47	17	28	44 9	14513.99
1867	. 49	2	15	43 10	11912.39
1868	. 70	3	24	15 8	12010.81
1869	65	6	24	23 6	124 9.87
1870	93	3	22	16 15	14911.92
1871	62	6	24	16 18	12610.15
1872	151	10	52	35 14	26216.58
1873	92	4	37	7 25	165 9.68
1874	134	8	50	17 25	22911.67
1875	136	6	36	14 31	22311.71
1876	122	4	40	22 18	20611.13
1877	122	6	34	21 21	20410.56
1878	70	3	28	11 18	130 6.59
1879	59	5	31	9 16	120 5.95

The percentage in 1879, is lower than in any previous year; and the number of deaths from the five diseases combined, less than in any year since 1868.

Deaths from Asiatic cholera are not included in these statistics; they have been as follows: in 1849, 158; in 1854, 159; in 1866, 14; total, 331.

Diphtheria. The number of deaths from this cause in 1879, was 107; which was less than half of the number in the preceding year; yet it prevailed throughout the year to an extent to be considered epidemic; there were 26 deaths in the first quarter of the year; 21, in the second quarter; 20, in the third quarter; and 40, in the fourth. Of the 107 decedents, 100, or 98.46 per cent., were children under 10 years old.

The statistics of sex and parentage were as follows:-

Males, 46; females, 61; total, 107, or 5.30 per cent. of all. American parentage, 54; foreign parentage, 53.

The first cases of this disease reported in Providence were in 1858; the following are the statistics in regard to it, for the 22 years, 1858-1879:—

Deaths from known causes in 22 years, 30,038.

From diphtheria, 1,093; annual average, 50.

Males, 543; females, 550; total, 1,093, or 3.64 per cent. of all.

American parentage, 527; foreign parentage, 566.

In the following table are given the statistics as to sex, parentage, season and locality, for each of the twenty-two years, 1858– 1879, and for the whole period:—

DEATHS FROM DIPHTHERIA IN PROVIDENCE, TWENTY-TWO YEARS.

1858, 1859	1859, 1860.	0. 1861.	11. 186	1862, 1863.	3. 1864	1865	1886.	. 1867.	1864, 1865, 1865, 1867, 1868, 1870, 1871, 1872, 1873, 1874, 1875, 1876, 1877, 1878, 1879,	1869.	1870.	1871.	1872.	1873.	1874.	1876.	1876. 1	1877.	1878.	1879.	Total.
171		2.60	22 1	17 42 .92 8.54	8.97	1.85	1.08	0.53	0.46	1.05	0.80	1.69	1.71	1.35	1.02	0.74	5.951	296	246	107	1.093
00 00		22	100	481	18 20 21	SI 4		- 10 to	50 04 50 00	2010	10.10	120	10	14	99	© 00	55.55	146	136	46	548
E 9		69	13	111 3	98		00	14	41.0	D1-	4.9	47.	28	15	11	10	88	129	105	25.52	527
: :01 :		:01-	HH01:	00 : : =	101-1010		H .00 .	91.10	01	00 00			02 44 42 64	2044	01 CH :		10 .410	18031	4523	01-010	25.85
:		:0100 00	11 1				11 11			- ! ! ! !	11		- 11	4	910	1 1	1000000	25°25	13.82	00000	82448
.400		0 50 60 60	0 04 04 4	- A D D4	1400		4 40 00	0100	10	-01		00 to 01	001 :4		- 00 C0 F	-01-4-1	1333	8228	888	888	071
:-=-		00000	0-0:	HH046	81-10 to -			F - 69 69 6		00000	111		20 - 00 - 1	00 - 01 :	H : 010	H 01 01 :0	00124	20.8.8.05	8448	F-4640	24 2 2 2 3
0101		004::	0014	111	111				N	101 : 4				-0000	-		S-Jares	8=82×	22225	5.2.2	8185

The connection of season with the mortality from diphtheria is a subject which has received considerable attention in these reports, since the disease has prevailed so extensively in our city; and facts of importance and interest are shown by the study, particularly when compared with similar statistics in regard to scarlatina and croup.

The following table shows the number and the percentage of deaths in Providence, from diphtheria, scarlatina, and croup, in each month and quarter of the year. The figures for scarlatina and croup include the aggregate mortality from these diseases during the twenty-five years, from 1855 to 1879 inclusive. The figures for diphtheria are for twenty-two years, from 1858 to 1879 inclusive.

		HBRIA. -1879.		ATINA. -1879.		OUP. -18 79.
Months.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
January	101	9.24	207	11.49	101	18.17
February	71	6.49	184	10.22	89	11.60
March	86	7.87	159	8.88	86	11.91
lst quarter	258	23.60	560	80.54	276	85.98
April	72	6.59	151	8.88	64	8.34
May	68	6.33	157	8.72	42	5.48
June	67	6.13	145	8.05	27	8.52
id quarter	207	18.94	458	25.15	183	17.84
July	41	8.75	96	5.88	27	8.52
August	44	4.08	75	4.16	11	1.48
September	98	8.51	88	4.89	40	5.99
3d quarter	178	16.29	259	14.38	78	10.17
October	170	15.55	128	7.11	78	9.52
November	158.	14.46	178	9.61	99	12.91
December	122	11.16	238	18.91	108	14.08
th quarter	450	41.17	539	29.98	280	86.51
Totals.	1,098	100.00	1,801	100.00	767	100.00

Dividing the year into periods of four months, the numbers and proportions are as follows:-

	DIPHT	HERIA.	SCARL	ATINA.	CR	OUP.
SEASONS.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.
1st 4 months	330	30.19	701	38.92	340	44.32
2d 4 months	220	20.13	473	26.26	107	13.95
3d 4 months	543	49.68	627	34.82	320	41.73

Dividing the year into two periods, the numbers and proportions are:—

The state of the s	DIPHT	HERIA.	SCARL	ATINA.	CRO	our.
SEASONS.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.
1st 6 months	465 628	42.54 57.46	1,003	55.69 44.31	400	53.32

From these figures we find that the smallest percentage from each disease is in the second four months of the year; but in respect to the prevalence in the first third of the year, and the last, there are marked differences.

Diphtheria shows a large excess in the last four months of the year; and 57.46 per cent. of all the deaths were in the last six months.

In Scarlatina and Oroup, it is exactly the reverse of this; in both cases the figures for the first four months are considerably larger than those for the last four; 55.69 per cent. of the deaths from scarlatina were in the first six months of the year; of those from croup, 53.32 per cent.

In the whole period, the smallest percentage of mortality in any month was for diphtheria 3.75 in July; for scarlatina 4.16, and for croup 1.43, both in August. The highest monthly percentage from diphtheria, was 15.55 in October; from scarlatina, 13.21, and from croup 14.08; both in December.

It has been stated that the first cases of diphtheria reported in Providence were in 1858; and there have been some cases each year since then; but the table on page 64, shows that for a long time the number in each year was small; the whole number in the eighteen years, from 1858 to 1875 inclusive, being only 334, or an average of 18.5 annually. But since 1876 it has prevailed as a severe epidemic, and in the four years, 1876 to 1879 inclusive, the number of deaths from this cause was 759, more than twice as many as in the eighteen years previous, and giving an annual average of 189.7 deaths during the epidemic; the annual average during the whole period of twenty-two years, 1858 to 1879, was nearly 50.

To show the differences between the statistics of the disease as epidemic or not epidemic, the following table has been prepared, which shows, side by side, certain statistics of diphtheria in Providence, first, for eighteen years, from 1858 to 1875 inclusive, when the disease was not at any time epidemic; second, for the four years, from 1876 to 1879 inclusive, when the disease prevailed as a very severe epidemic, and third, for the two periods combined:—

Dan S		HERIA. -1875.		HERIA. -1879.	DIPHT 1858-	HERIA. -1870.
MONTHS.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths,	Per cent.
January	31	9.28	70	9.22	101	9.94
February	32	9.58	39	5.14	71	6.49
March	32	9.58	54	7.11	86	7.87
1st quarter	95	28.44	163	21.47	258	23.00
April	19	5.69	.53	6.98	79	6,50
Мау	15	4.49	53	6.98	68	6.22
June	23	6.89	44	5.80	67	6.15
2d quarter	57	17.07	150	19.76	207	18,96
July	19	5.69	22	2.90	41	3.75
August	10	2.99	84	4.48	44	4.03
September	25	7.49	68	8.96	93	8.51
3d quarter	54	16.17	124	16.34	178	16,29
October	40	11.98	130	17.13	170	15.55
November	45	13.47	113	14.89	158	14.40
December	43	12.87	79	10.41	122	11.10
4th quarter	128	38.32	322	42.43	450	41.17
Totals	834	100.00	759	100.00	1,093	100.00
Males	162	48.50	361	50.20	543	49.0
Females	172	51.50	378	49.80	550	50.3
American	196	58.68	331	43 61	827	48.9
Foreign	138	41.32	428	56.39	566	517

There is quite a difference in the percentages of deaths in the different months, as given in the first two columns. January shows a slight reduction of mortality when the disease was epidemic, as compared with the earlier period; February and March show a much greater reduction of mortality, during the epidemic; April and May show an increased mortality, and June and July a largely decreased mortality when the disease is epidemic. After July there is an increased proportional mortality, when the disease is epidemic, in each month until December. By

quarters of the year there is a decreased comparative mortality when the disease is epidemic, in the first quarter, and an increased mortality in the other seasons of the year.

By periods of four months the percentages of mortality were as follows:—

DIPHTHERIA.

•	1868	-1875.	1876-	-1879.	1858	1879.
.skosaes	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.
lst 4 months	114	34.18	216	28.45	330	80.19
td 4 months	67	20.06	158	20.16	230	90.18
3d 4 months	158	45.81	390	51.39	543	49.68

By periods of six months, the percentages of mortality were as follows:—

DIPHTHERIA.

	1858	-1875.	1876-	-1879.	1858–1879.		
Seasons.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	
lst 6 months	159	45.51	818	41.23	465	49 54	
2d 6 months	189	54.49	446	58.77	628	57.46	

At all times, we see that the disease is more prevalent in the latter part of the year, but the difference between the earlier and later parts of the year is much greater when the disease is epidemic.

There are also very great differences in regard to sex and parentage in the mortality from diphtheria, when epidemic or not epidemic. In the first period, 1858–1875, the decedents were 48.50 males, and 51.50 females in each 100; but from 1876–1879, when the disease was epidemic, the proportions were 50.20 males and 49.80 females in each 100. The same change is true in regard to parentage; during the first period, there were 58.68 of American and 41.32 of foreign parentage in each 100 decedents; but from 1876 to 1879, there were 48.61 of American, and 56.39 of foreign parentage in each 100.

Erysipelas. This was returned as the cause of 10 deaths in 1879; males, 4; females, 6; total, 10, or 0.49 per cent. of all from known causes; American parentage, 3; foreign, 7.

The statistics for the twenty-three years, 1856-1879, are:-

Males, 82; females, 88; total, 170, or 0.53 per cent. of all.

American parentage, 99; foreign parentage, 71.

Exposure and Neglect. The number of deaths in 1879, ascribed to exposure or neglect, was 3; two of the decedents were less than a year old, and one was between 70 and 80.

Males, 1; females, 2; total, 3, or 0.15 per cent. of all.

American parentage, 1; foreign parentage, 2.

Fever, Typhoid. There were 40 deaths from this cause in 1879; 7 less than in 1878.

Males, 19; females, 21; total, 40, or 1.98 per cent. of all. American parentage, 26; foreign parentage, 14.

In the following statistics for the twenty-four years, 1856-1879, besides the deaths from typhoid, are a few that have been reported in some years from "fever" and "bilious fever," and about 30 from "typhus fever":—

Whole number, 1,022; annual average, 42.

Males, 527; females, 495; total, 1,022, or 3.20 per cent. of all.

American parentage, 570; foreign parentage, 452.

Heart, Diseases of. The number of deaths in 1879 from diseases of the heart was 111, including 6 from "hypertrophy;" the whole number was 28 more than in 1878.

Males, 57; females, 54; total, 111, or 5.51 per cent. of all.

American parentage, 63; foreign parentage, 48.

In the twenty-four years, 1856-1879, the number of deaths from diseases of the heart, was 1,478, or 4.62 per cent. of all from known causes; annual average, 61.

The particulars for each year are as follows :-

	SEX		Par	RENTAGE.	Whole			
	Males.	Females.	American.	Foreign.	Number. Percentage			
856	. 8	18	16	5	212.03			
857	. 13	18	22	9	81			
858	. 18	25	28	8	36			
.859	. 18	11	17	7	242.76			
860	. 15	16	91	10	81			
961	. 23	18	29	19	414.05			
802	. 26	28	88	16	495.54			
968	23	19	80	19	42			
864	. 26	39	44	14	584.68			
.865	. 18	29	88	14	478.95			
866	. 19	21	27	18	408.95			
.867	96	93	35	14	495.19			
868	25	27	84	18	524.79			
.869	83	99	85	27	625.00			
1870	28	81	88	26	594.72			
871	40	37	46	81	776.22			
1872	43	49	68	29	925.82			
1873	46	54	59	41	1005.88			
1874	48	58	62	49	1045.29			
875								
876	41	47	51	87	884.75			
877		•						
878								
					1115.51			
14 years' total.	711	767	996	552	1,4784.62			

The number of deaths from heart disease in 1879 was greater than in any previous year, though there are four years when the percentage was greater: 1862, 1871, 1872 and 1873. Of the 1,478 decedents 62.65 per cent. were of American, and 37.35 per cent. of foreign parentage.

Hemorrhage. Ten deaths from this cause were reported in 1879; this was 4 less than in 1878, and 0.50 per cent. of all from known causes.

Males, 4; females, 6.
American parentage, 5; foreign parentage, 5.

In the table, 3 are specified as "hemorrhage from the lungs," and 1 "from the stomach;" there were also 2 cases of "uterine hemorrhage," and one of "cerebral."

Hernia was given as the cause of 5 deaths in 1879; the same number as in 1878, and 0.25 per cent. of all from known causes; 2 males, 3 females: 2 of American, and 3 of foreign parentage.

Hooping Cough. There were 15 deaths from this cause in 1879, which was 28 less than in 1878. Though the number was so small, the disease continued throughout the year, some deaths occurring in each month except July and September.

Males, 5; females, 10; total, 15, or 0.74 per cent. of all. American parentage, 8; foreign parentage, 7.

In a majority of cases this disease is accompanied or followed by some other, which however is not accounted as the cause of death, but merely as a consequence of the hooping cough, that being the primary cause. Thus in 1879, there were only 4 cases where hooping cough alone was reported as the cause of death; in 5 cases it was followed by convulsions; in 4, by pneumonia; in 1 each, by congestion of the lungs, and catarrh.

The statistics for the twenty-four years, 1856-1879, are :-

Whole number, 386; annual average, 16.

Males, 164; females, 222; total, 386, or 1.21 per cent. of all.

American parentage, 161; foreign parentage, 225.

Hydrocephalus. The number of deaths from hydrocephalus in 1879 was 35, or 4 less than in the year previous.

Males, 21; females, 14; total, 35, or 1.73 per cent. of all American parentage, 17; foreign parentage, 18.

For the twenty-four years, 1856-1879, the statistics are :-

Whole number 733; annual average, 30.

Males, 400; females, 333; total, 733, or 2.29 per cent. of all.

American parentage, 370; foreign parentage, 363.

Insanity. There were 12 deaths reported in 1879 from insanity; 5 less than in 1878.

Males, 8; females, 4: total, 12, or 0.59 per cent. of all.

American parentage, 7; foreign parentage, 5.

The statistics for the twenty-four years, 1856-1879, are as follows:-

Whole number, 248: annual average, 10.

Males, 120; females, 128; total, 2.48, or 0.78 per cent. of all.

American parentage, 177; foreign parentage, 71.

Many of these deaths occurred at Butler Hospital for the insane, which is within the limits of the city, but where the patients, of course, are from many other places than Providence.

Intemperance. This was the cause of 7 deaths in 1879, including 2 from delirium tremens; the whole number was two less than in 1878. All the decedents were males; 2 of American, and 5 of foreign parentage; 0.35 per cent. of all from known causes.

The following statistics are for twenty-three years, 1857-1879:-

Whole number from known causes, 30,938.

From intemperance, 229: annual average, 10.

Males, 168; females, 61; total, 229, or 0.74 per cent. of all.

American parentage, 66; foreign parentage, 163.

Of the 229 decedents, 73.36 per cent. were males; and 71.18 per cent. of foreign parentage.

Kidneys, Diseases of, caused 43 deaths in 1879; 4 less than in 1878. Of these deaths, 30 were from "Bright's Disease."

The statistics for the year 1879 are:-

Males, 26; females, 17; total, 48, or 2.18 per cent. of all.

American parentage, 20; foreign parentage, 28.

For the twenty-four years, 1856-1879:-

Whole number, 480; annual average, 20.

Males, 286; females, 194; total, 480, or 1.50 per cent. of all.

American parentage, 274; foreign parentage, 206.

Liver, Diseases of. The number of deaths from diseases of the liver in 1879 was 21; 3 more than in 1878, and 1.04 per cent. of all from known causes.

Males, 13; females, 8; total, 21, or 1.04 per cent. of all.

American parentage, 9; foreign parentage, 12.

For the twenty-four years, 1856-1879, the numbers are:—
Whole number, 858; annual average, 15.

Males, 181; females, 177; total, 358, or 1.12 per cent. of all.

American parentage, 175; foreign parentage, 183.

Malformations. Eleven deaths were reported from this cause in 1879, which was just half of the number in the preceding year. Nine cases were malformation of the heart, the others were not specified.

Males, 9; females, 2; total, 11, or 0.54 per cent. of all. American parentage, 4; foreign parentage, 7.

The statistics for the twenty-four years, 1856-1879, are :-

Males, 119; females, 66; total, 185, or 0.58 per cent. of all.

American parentage, 118; foreign parentage, 67.

Marasmus was stated as the cause of 28 deaths in 1879; 13 less than in 1878; all but one of the decedents were children under 5 years old.

> Males, 13; females, 15; total, 28, or 1.39 per cent. of all. American parentage, 15; foreign parentage, 13.

The following are the statistics for the twenty-four years, 1856-1879:-

Whole number, 618; annual average, 26.

Males, 329; females, 289; total, 618, or 1.93 per cent. of all.

American parentage, 311; foreign parentage, 307.

Measles. There were no deaths from measles in Providence in 1879; this was the more remarkable, as in 1878, the number from this cause, 50, was the largest ever recorded; all the deaths in 1878 occurred in the first six months of the year; so that there have been no deaths from measles since June, 1878. This irregularity of appearance has always characterized this disease in Providence. The following table shows the number in each year since 1856:—

1856 1	1862 1	1968 4	1874
1857 2	186316	1869 6	1875
185840	1864 7	187016	1876 0
1859 0	1865 7	1871 1	1877 9
1860 0	1866 9	1872 7	1878
1861 6	1867 0	187328	1879 0
			Total, 24 years207

The statistics for the twenty-four years, 1856-1879, are :-

Males, 100; females, 107; total, 207, or 0.65 per cent. of all.

American parentage, 60; foreign parentage, 147.

Of the 207 decedents, 71.01 per cent. were of foreign parentage.

Meningitis, Cerebro-Spinal. There were 7 deaths from this cause in 1879, the same number as in each of the two preceding years.

Males, 6; female, 1; total, 7, or 0.35 per cent. of all. American parentage, 4; foreign parentage, 3.

The first cases of this disease in Providence were reported in 1864, since when the number in each year has been as follows:—

1864 2	1870 0	1876 4
1865 2	187112	1877 7
1866 1	187210	1878 7
1867 0	187339	1879 7
1868 1	187410	
1869 0	1875 6	Total, 16 years 108

Whole number from known causes, in 16 years, 24,129.

From cerebro-spinal meningitis, 108.

Males, 61; females, 47; total, 108, or 0.45 per cent. of all.

American parentage, 52; foreign parentage, 56.

The number and percentage of deaths in each division of age and according to sex and parentage, were as follows for the sixteen years, 1864-1879 inclusive:—

AGES.	Under 1 Year.	1 and under 2.	2 and under 5.	5 and under 10,	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	Males.	Females.	American.	Foreign.	Total.
No.	9	ю	14	32	13	14	13	10	1	1	1	19	1.7	53	99	108
PER CENT.	5.65	4.63	19.96	23.15	пп	13.96	19.04	9.36	87.9	.98	98.	26.48	43.52	48.15	51.85	100.00

Sixty-two of these decedents, or 57.40 per cent., were less than 15 years old.

Murder and Homicide. One death only was reported under this head in 1879; that of a foreign male between 50 and 60 years old.

Old Age. There were 64 deaths in 1879 reported from this cause, 3 more than in 1878, and 3.17 per cent. of all from known causes. Five of the decedents were more than 90 years old; 9 less than in 1878. The statistics are as follows:—

Males, 27; females, 37; total, 64, or 3.17 per cent. of all.

American parentage, 40; foreign parentage, 24.

For the twenty-four years, 1856-1879 :-

Whole number, 1,254; annual, average, 52.

Males, 416; females, 838; total, 1,254, or 3.92 per cent. of all.

American parentage, 775; foreign parentage, 479.

Of the 1,254 decedents, 66.83 per cent. were females, and 61.80 per cent. of American parentage.

Only a small proportion of the aged decedents in Providence are reported under the head of old age. Thus in 1879, there were 242 decedents, 70 years old and over; 147 of American, and 95 of foreign parentage; this was 12.00 per cent. of all the decedents from known causes in 1879.

In the twenty-four years, 1856-1879, the decedents of 70 years and over, numbered 3,473; 2,423 of American, and 1,050 of foreign parentage; 10.86 per cent. of all from known causes.

Paralysis. This is generally a symptom rather than a distinct disease, and the cause in most instances ought to be given more fully than it is. It was given in 1879 as the cause of 30 deaths; 6 more than in 1878, and 1.49 per cent. of all.

Males, 20; females, 10; total, 30, or 1.49 per cent. of all.

American parentage, 20; foreign parentage, 10.

The statistics for the twenty-four years, 1856-1879, are :-

Whole number, 408; annual average, 17.

Males, 213; females, 195; total, 408, or 1.28 per cent. of all.

American parentage, 306; foreign parentage, 102.

The statistics for apoplexy and paralysis are generally given together in this connection; the following are the figures for each of the 24 years, 1856-1879:—

Years.	Apoplexy.	Paralysis.	Total.	Years.	Apoplexy.	Paralysis.	Total
1856	21	2	28	1869	82	16	48
1857		9	27	1870	28	24	52
1858	15	8	18	1871	83	28	61
1859	16	6	22	1872	81	20	51
1860	17	11	28	1878	86	21	57
1861	14	11	25	1874	86	23	59
1862	19	5	24	1875	41	35	76
1863	20	8	28	1876	44	24	68
1864		16	29	1877	50	24	74
1865	20	18	88	1878	49	24	66
1866		8	29	1879	56	30	86
1867	81	18	49				
1868	22		46 7	otals. 2	t v'rs. 676	408	1.084

The statistics of both causes for the 24 years, as regards sex and parentage are:—

Whole number, 1,084; annual average, 45.

Males, 546; females, 538; total, 1,084, or 3.39 per cent. of all.

American parentage, 780; foreign parentage, 304.

The decedents were 71.96 per cent. of American parentage, and 28.04 of foreign parentage.

Peritonitis. The number of deaths from this cause in 1879, was 14; 3 less than in the previous year.

Males, 6; females, 8; total, 14, or 0.69 per cent. of all.

American parentage, 8; foreign parentage, 6;

The statistics for the 24 years, 1856-1879, are as follows:—

Whole number, 215; annual average, 9.

Males, 96; females, 119; total, 215, or 0.67 per cent. of all.

American parentage, 122; foreign parentage, 93.

Pneumonia. There were 145 deaths from pneumonia and 14 from congestion of the lungs, in 1879; total, 159; this was 19 more than in 1878, and 61 more than in 1877. The deaths in the different seasons of the year were:—

1st quarter74	8d quarter11
2d quarter	4th quarter

Of these decedents, 68, or 42.77 per cent. were under 10 years of age; and 57, or 85.85 per cent. were over 50.

Males, 76; females, 88; total, 159, or 7.88 per cent. of all.

American parentage, 73; foreign parentage, 86.

The statistics for the 24 years, 1856-1879, are:

Whole number, 2,316; annual average, 96.

Males, 1,192; females, 1,124; total, 2,816, or 7.25 per cent. of all. American parentage, 1,167; foreign parentage, 1,149.

Scarlatina. This disease prevailed in Providence in 1879, as a very severe epidemic, especially in the latter part of the year. The number of deaths caused by it was 252, a larger number than ever before in a single year, except in the epidemic of 1874, when there were 268 deaths. The decedents were divided as to age as follows: under 1 year, 9; between 1 and 2 years, 29; between 2 and 5, 113, or 44.84 per cent. of all the decedents from this cause; between 5 and 10 years, 85; between 10 and 15, 12; between 20 and 80, 3; between 30 and 40, 1.

The deaths were distributed through the year as follows:-

1st quarter84	8d quarter 84
2d quarter28	4th quarter161

The 161 deaths in the fourth quarter of the year were 68.84 per cent. of the whole number; the largest number in any one month was 71 in December, a larger number than ever occurred in the city before in any one month. The largest number in one ward was 54, in the ninth ward. The statistics of sex and parentage are:—

Males, 135; females, 117; total, 252, or 12.49 per cent. of all.

American parentage, 101; foreign parentage, 151.

The statistics for the 24 years, 1856-1879, are-

Whole number, 1,759; annual average, 78.

Males, 856; females, 903; total, 1,759, or 5.50 per cent. of all.

American parentage, 784; foreign parentage, 1,025.

The following table gives the particulars of sex, parentage, season, and locality for each of the 25 years, 1855-1879, and for the whole period:—

SCARLATINA IN PROVIDENCE, TWENTY-FIVE YEARS.

: <u>-</u> i	125	878 829	85. 84.	24355545 24355545 24355545 2435545 2435545 2435545 2435545 2435545 2435545 243554 2435	200 200 201 201 201 201 201 201 201 201
Total	1,801		<u>ج</u> ۾		
1679.	13.45	185	<u> </u>	230000400227	8-8525878
B78.	75.89	88	23	988780000000000000000000000000000000000	
1977.	0.63	42	87.0 3	60 - 61	
676.	1.87	20	*#	400 40	4
1875.	86.	\$3	83	ფენტენტაფას	≅ 201-∞≅450€
1874.	88	23	F 2	822222222	2 8 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
1878.	35.7 26.	85	25	<u> </u>	8 * 2 * 0 0 1 2 8
873.	28.	* =	*=		D 4 . H4D
1870. 1871. 1872.	28.	55 gp	0.5	854A8A	~84~××
.870.	88	28	28	. most en :	
1860.	138	43	28	\$12500441014w	2 0 2 0 2 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
1968.	5.9	83	22		©
1866. 1867.	1.08	-3 00			48 4
	. B.		86 14		
1865.	5.00	32	33	0 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	න සැති සැට සිද්
1864.	17.11	85	82		10110888
1963.	20 eq	28	136		@- -
	23	80 80	8 3		3 4 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
1861.	85	22	22	444688	Mag-190-1
1880	EF.	2	II.	8 -	484 968
1869.	5.18	22	#	700004-04-4-0	⊕ ∟64 ≈ 50 æ : :
1868.	£ 8.	28	22	82865	
1857. 1858. 1859.	22	20	84	<u> </u>	5884484
.999	128	58	45	- 228214400-00	· 8585518 ·
1886.	25	88	28	::	6 − 8 − 8 − 8 − 8 − 8 − 8 − 8 − 8 − 8 −
BCARLATIKA. 1	Phole number ercentage	ex: Males Females	ARENTAGE: American Foreign	BASONS: Pantary February March May June July Beyten Gebreen Gebreen December	Mard II. Mard II. III. IV. V V. V VI.

A comparison of the statistics of scarlatina with those of diphtheria and croup, is given under the head of "diphtheria."

Scrofula. The number of deaths from scrofula in 1879 was 5; 2 less than in 1878.

Male, 1; females, 4; total, 5, or 0.25 per cent. of all. American parentage, 8; foreign parentage, 2.

Small-pox. There have been no deaths from small-pox in Providence, since 1875. For the twenty-four years, 1856-1879, the statistics are as follows:—

Males, 58; females, 21; total, 79, or 0.25 per cent of all. American parentage, 48; foreign parentage, 31.

The particulars for each year are as follows:-

1856	2 4 2 7	2	American	3	5 5 4	0.57
1857	2 4 2 7	2	5	3	5 5 4	0.57
858	2 4 2 7	2	5	3 3	5 5 4	0.57
.859	2 2 7	2	5	8 8	5 5 4	0.58
	4 2 7	2	1	3 8	4	0.58
1861	2 7	2	1	8 4	4 7	0.45
1863	. 2 . 7	2	1 8	8 4	7	0.45
.863	7	2	8	4	7	0.56
1864 1865	. 7	2	8	4	7	0.56
1865	. 9	2		•••••		•••••
1866			8	8	11	n est
	•••••					
1007		•••••••			•••••	
100/	1	• • • • • • • • • • • • • • • • • • • •	1	•••••	1	0.11
1968	1	•••••	1		1	0.00
1869	8	•••••			8	
1870		2	• • • • • • • • • • • • • • • • • • • •	2	2	0.16
1871		8	1	2	3	0.94
1872	9	4		4	18	0.85
1873	12	3	8	6	14	0.85
1874						•••••
1875	1			1	1	
1876			•••••			
1877	•••••		•••••		*********	
1878	••••		••••••		*********	
1879						
	_	_	_ 48	_	_	

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Skin, Diseases of. There were 2 deaths from diseases of the skin in 1879, one male, and one female; one of American, and one of foreign parentage; 0.10 per cent. of all from known causes.

Spine, Diseases of. The number of deaths from diseases of the spine was 6; 2 more than in 1878, and 0.30 per cent. of all. Males, 4; females, 2; American parentage, 3; foreign, 3.

Stomach, Diseases of. The deaths from diseases of the stomach in 1879, numbered 15; 11 of which were called "inflammation;" the number was one more than in the previous year, and 0.74 per cent. of all. Males, 6; females, 9; American parentage, 7; foreign parentage, 8.

Suicide. There were 6 cases of suicide in 1879; one less than in 1878. In one case it was by laudanum; in one, by Paris green; in one, by hanging; in one, by prusic acid; in two by shooting.

Males, 5; females, 1; total 6, or 0.30 per cent. of all. American parentage, 2; foreign parentage, 4.

The statistics for the 24 years, 1856-1879, are:

Males, 103; females, 42; total, 145, or 0.45 per cent of all.

American parentage, 81; foreign parentage, 64.

Teething. The number of deaths ascribed to this cause in 1879, was 13; 6 more than in 1878.

Males, 4; females, 9; total, 18, or 0.64 per cent. of all. American parentage, 5; foreign parentage, 8.

Tuberculosis. There were 20 deaths reported from tuberculosis in 1879; 3 more than in 1878.

Males, 9; females, 11; total, 20, or 0.99 per cent. of all. American parentage, 10; foreign parentage, 10.

This number is very small in comparison with the number of diseases that might properly be classed under this head. There

are many tubercular diseases, as consumption, hydrocephelus and others, which if united in this general class, would show a larger percentage than any other cause.

Tumors. There were 7 deaths caused by tumors in 1879; 1 more than in 1878, and 0.35 per cent. of all from known causes; all the decedents were females; 4 of American, and 3 of foreign parentage. The tumors were specified as follows: 2, abdominal; 2, ovarian; 3, uterine.

Unknown Causes. The number of deaths in 1879, for which no cause was reported was 9; this was 7 less than in 1878.

Males, 4; females, 5; total 9, or 0.44 per cent. of all deaths in 1879. American parentage, 6; foreign, 3. Five of the decedents were less than one year old.

The statistics for the twenty-four years, 1856-1879, are:-

Whole number, 533; annual average, 22.

Males, 277; females, 256; total, 533, or 1.64 per cent. of all.

44

٠.

American parentage, 176; foreign parentage, 357.

Of the 533 decedents, 66.98 per cent. were of foreign parentage.

STILL-BORN.

The statistics of still-born children in Providence, in 1879, were as follows:—

Number. The number of this class was 122; 19 less than in 1878, and 4.84 per cent. of the 2,522 children born in 1879, or one in 20.67. The number in 1878, should have been 141 instead of 139 as given in the report for that year.

Sex. The number of males was 63, or 19 less than in 1878; the number of females was 59, the same as in the previous year.

Color. White, 112; colored, 10.

Locality. First ward, 19; second, 2; third, 9; fourth, 2; fifth, 11; sixth, 7; seventh, 8; eighth, 15; ninth, 22; tenth, 27.

Season. January, 12; February, March, and April, 7 each; May, 11; June, 12; July, 14; August, 11; September, 10; October, 14; November, 7; December, 10.

The number and percentage in each quarter of the year were as follows:—

	Number.	Per cent.		Number.	Per cent.
lst quarter	26	21.81	8d quarter	85	98.69
2d quarter	80	24.59	4th quarter	81	25.41

In the following table, the first letter indicates the birth-place of the father, the second, that of the mother.

A. and A 60	E. and A 8	A. and E 8
	G. and A 1	
E. and E 7	B. A. and A 1	S. and I 1
G. and G 1	8. and A 1	1. and 8 1
B. A. and B. A 9	P. and A 1	E. and S 1
It. and It 1	A. and I 9	A. and B. A 1
L and A 3	B. A. and I 1	Total123

A.—American; I.—Irish; E.—English; S.—Scotch; G.—German; B. A.

—British American; It.—Italian; P.—Portuguese

The number of children born in Providence in 1879, was 2,522; of these, 122, or 4.84 per cent., were still-born. Of the 2,522 children born, 1,288, or 51.07 per cent., had American mothers; 778, or 30.85 per cent., had Irish mothers; and 456, or 18.08 per cent., had mothers who were natives of other foreign countries. Of the 122 still-born children, 70, or 57.38 per cent., were children of American mothers; 34, or 27.87 per cent., were children of Irish mothers; and 18, or 14.75 per cent., were children whose mothers were natives of other foreign countries.

Of the 1,288 children born of American mothers, 70, or 5.44 per cent., were still-born; of the 778 with Irish mothers, 34, or 4.37 per cent., were still-born; of the 456 children of mothers born in other foreign countries, 18, or 3.95 per cent., were still-born. The proportions are; to American mothers, one still-born in 18.40; to Irish mothers, one in 22.88; to mothers of other nativities, one in 25.33.

These statistics may be compared with those of 1878 as follows: The whole number and proportion of children born to American mothers in 1879 was considerably larger than in the previous year; the number and proportion of still-born children of this class were also somewhat larger in 1879. Of children of Irish mothers in 1879, the number and proportion were less than in the previous year; the number and proportion of still-born among them, very much less. Of the children of other foreign mothers, the number in 1879, was rather less than in 1878, but the proportion a little larger; of still-born children in this class, the number and proportion were considerably larger in 1879 than in 1878.

For the twenty-four years, 1856-1879, the statistics are as follows:—

Number of children born, 46,422.

Number of still-born, 2,727, or 5.87 per cent. of all; or one in 17.02. Of these 2,727 still-born children, 2,557 were white; 170 were colored; 1,565, or 57.43 per cent. were males; and 1,162, or 42.57 per cent. females.

Reckoning the parentage according to the father's birth-place, 1,157 were American; 1,217 were Irish; and 346 were of other foreign fathers; of 7 the parentage was unknown.

STILL-BORN.

Reckoning the parentage according to the mother's birth-place, the still-born children were in 24 years, as follows:—

American parentage	1,159
Irish	1,272
English, Scotch or Welch	161
German	28
British American	63
Portuguese	15
Other foreign	22
Unknown	
Total still born	

GENERAL SUMMARY.

The following recapitulatory table will give the general results of registration in Providence during the 25 years, 1855 to 1879 inclusive:—

TABLE XXI. BIRTHS, MARRIAGES, AND DEATHS.

YEAR.	Population.	Births.	To population, one birth in	Marriages.	Of population, one person married in	Deaths.	Of population, one death in
1855	47,785	1,600	29.86	636	37.56	991	48.22
1856	48,727	1,675	29.09	656	37.14	1,065	45.73
1857	49,457	1,688	29.30	614	40.27	925	53.46
1858	49,457	1,724	28.68	560	44.15	1,017	48.63
1859	50,100	1,593	31.45	616	40.66	899	55.72
1860	50,666	1,648	30.74	633	40.02	1,001	50 61
1861	50,666	1,725	29.37	547	46.31	1,051	48.21
1862	52,238	1,529	84.16	547	47.75	914	57.35
1863	53,024	1.307	40.56	610	43.46	1,214	45.67
1864	53,810	1.344	40.04	789	36.40	1,281	42 01
1865	54,595	1,298	42.06	709	38.50	1,211	45.08
1866	55,709	1,632	34.13	812	34.30	1,036	53.77
1867	56,824	1,624	84.99	855	33.23	960	59.19
1868	64,138	1,866	34.37	872	36.77	1,110	57.78
1869	66,522	1,819	36,57	930	35.76	1,256	52.96
1870	68,904	1,889	36.48	959	35,93	1,263	54.55
1871	72,011	1,960	86.74	943	38.13	1,254	57.42
1872	75,118	2,206	34.05	1,084	34.63	1,603	46.86
1873	78,225	2,128	36.76	1,150	34 01	1,719	45.51
1874	96,452	2,780	84.69	1,176	41.01	1,987	48.54
1875	100,675	2,664	87.79	1,079	46.65	1,915	53.57
1876	101,500	2,631	38.58	949	53.48	1,865	54.42
1877	100,675	2,543	39.58	953	52.82	1,938	51.95
1878	101,000	2,627	38.45	1,016	49.70	1,989	50.78
1879	103,000	2,522	40.84	1,071	48.08	2,026	50.84
Average	68,051	1,921	35.42	829	41.04	1,340	50.78

BIRTHS AND DEATHS BY PARENTAGE.

The following recapitulatory table shows the number of births and deaths, by parentage, in Providence, in each of the twenty-four years, from 1856-1879 inclusive. In the three years, 1863, 1864, and 1865, there was an excess of deaths over the births, in the population of American parentage:—

TABLE XXII.

				<u> </u>			
	AMERIC	DAN PARE	NTAGE.	Foreign Parentage.			
YEAR.	Deaths.	Births.	Excess of Births.	Deaths.	Births.	Excess of Births	
1856	541	759	218	524	916	392	
1857	523	701	178	402	987	585	
1856	512	771	259	505	953	448	
1859	479	659	180	420	934	514	
1880	512	781	219	489	917	428	
1861	538	740	202	513	985	472	
1862	460	663	203	454	866	412	
1863	597	5 2 8	89	617	779	162	
1864	649	579	—70	632	765	183	
1865	631	- 546	85	580	752	172	
1866	513	726	213	523	906	883	
1967	528	725	197	432	899	467	
1968	596	882	296	514	981	470	
1869	627	926	299	629	898	264	
1870	642	890	248	621	999	878	
1871	643	980	887	611	980	369	
1879	832	1,085	908	771	1,171	400	
1873	862	1,008	141	. 857	1,125	268	
1874	885	1,217	882	1,102	1,563	461	
1875	914	1,180	266	1,001	1,484	488	
1876	875	1,169	294	990	1,462	479	
1877	908	1,238	830	1,030	1,305	275	
1878	889	1,215	826	1,100	1,412	819	
579	941	1,928	287	1,085	1,294	209	
24 years	16,097	21,091	4,994	16,402	25,331	8,929	

CITY ORDINANCES .- CHAPTER XLVII.

REGISTRATION OF DEATHS.

SECTION 1. There shall be appointed by the City Council a sufficient number of persons to act as undertakers, removable at the pleasure of the City Council.

Sec. 2. Whenever any person shall die in the city, the physician attending in his or her last sickness shall furnish to the undertaker attending the funeral, or to the city registrar, a certificate, giving the name of the person, date of death, and the disease or cause of his or her death.

SEC. 3. No person shall bury, or place in the tomb, or remove from the city, or otherwise dispose of the body of any human being who shall die in the city, without first reporting the death to the city registrar, and obtaining a permit from him.

Sec. 4. No permit shall be given as provided in section three, until the city registrar is furnished with the information in relation to the deceased person, required by the laws of the State, for record, so far as the same can be ascertained, together with the physician's certificate of the cause of death, whenever a physician has been in attendance, or a coroner's certificate, whenever a coroner's inquest has been held.

Whenever a permit for burial is applied for, in a case of death without the attendance of a physician, or if it is impossible to obtain the physician's certificate, the city registrar shall investigate the case so far as may be necessary, and when he has obtained satisfactory evidence in relation to the cause and circumstances of the death, he shall sign the certificate and give the required permit. If not satisfied in relation to the cause and circumstances of the death, or if, in his opinion, the public good requires it, he shall report the case to a coroner for investigation.


Sec. 5. Whenever the body of a human being who has died out of the city shall be brought here for burial, the undertaker, or other person attending the funeral, shall furnish the report required in sections three and four, with the exception of the physician's certificate.


SEC. 6. All funerals shall take place between sunrise and sunset, unless otherwise permitted or directed by the board of health. The top of every coffin deposited in the ground shall be at least three feet below the usual surface thereof.

Sec. 7. The city registrar shall cause an abstract of the returns of deaths made to him to be published in all the newspapers which contract to do the city printing, monthly, and oftener if required by the board of health. He shall also, on or before the first day of April, annually, prepare and present to the city council a statement of the number of births, marriages and deaths which occurred in the city during the year ending with the thirty-first day of December next preceding, with such other formation and suggestions in relation thereto as he may deem useful for the promotion of the public health, and other interests of the city.

SEC 8. No undertaker or other person shall bury or cause to be buried the body of any deceased person in the city, except in such grounds as are or may be designated as burying-grounds, and authorized to be used as such.

SEC. 9. Every person violating any provision of this chapter shall pay a fine of not less than five and not more than twenty dollars for each offence.





1881.]

CITY DOCUMENT.

N

TWENTY-SIXTH ANNUAL REPORT

UPON THE

BIRTHS, MARRIAGES, AND DEAT

IN THE

CITY OF PROVIDENCE:

FOR THE YEAR

1880,

By EDWIN M. SNOW M. D.,

SUPERINTENDENT OF HEALTH AND CITY REGISTRAR.



PROVIDENCE PRESS COMPANY, PRINTERS TO THE CITY. 1881.

PROV. PUB. LIB'

THE NEW YORK PUBLICLIBRARY

118672

ASTOR, LENOX AND TRIDEN FOUNDATIONS.
1892

THE CITY OF PROVIDENCE.

JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved May 20, 1881.]

RESOLVED, That the twenty-sixth annual report of the City Registrar, for the year ending December 31, 1880, be received, and that six hundred copies thereof, and of the communication accompanying the same, be printed in pamphlet form for the use of the City Council, under direction of the joint committee on printing.

A true copy: witness,

HENRY V. A. JOSLIN, CITY CLERK.

INTRODUCTION.

CITY REGISTRAR'S OFFICE, PROVIDENCE, May 16, 1881.

To the Honorable the City Council:

The Twenty-Sixth Annual Report of the City Registrar is herewith presented, being for the year ending December 31, 1880. During that year, there were 2,627 children born in Providence, 2,464 persons were married, and 2,080 persons died. The children born were 105 more, the persons married, 322 more, and the decedents, 54 more than in the year 1879.

According to the national census of 1880, the population of the city was 104,857, of whom 101,211 were whites, and 3,646 were colored. The ratio of births, marriages, and deaths, to population, for the year 1880, was therefore as follows:—

- 2,627 births, or one in 39.91; or 25.05 in each 1.000.
- 2,464 persons married, or one in 42.55; or 23.50 in each 1,000.
- 2,080 decedents, or one in 50.41; or 19.84 in each 1,000.

Among the white population of the city, there was, in 1880, one death in 51.01, or 19.60 in each 1,000; among the colored population there was one death in 37.98, or 26.33 in each 1,000.

The present report contains all the usual, and some additional annual and recapitulatory tables; and also, a table showing the population of Providence according to the census of 1880, by wards, and by nativity, sex, and color.

EDWIN M. SNOW, M. D.,

Superintendent of Health and City Registrar.

CONTENTS.

	A A
Wards months say and salor (Table I)	1-
Wards, months, sex and color, (Table I.)	. 1
Proportion of the sexes born at different seasons. Number and proportion of the sexes, 27 years	
Color:—Births, marriages and deaths in 25 years Parentage of children born in each ward, (Table III.) Number and percentage of American and of foreign children, 25 years.	
Number and percentage of American and of foreign children, 25 years.	. 8
Parentage:—Mixture of different nations, (Table IV.). Number of 1st, 2d, etc. children in 1880, and 1855-1879, (Table V.). Average number of children to cach mother, 26 years, (Table VI.). Number of the child, with age of the mother, (Table VII.)	
Average number of children to each mother, 26 years, (Table VI.)	12-
Number of the child, with age of the mother, (Table VII.)	14-
Plurality-births, statistics for 1880	16-
	204
MARKIAGES. Marriages at different seasons, 26 years. (Table VIII.)	20-0
Bride and groom born in same country. &c	22-
Proportions of American, foreign and mixed marriages, 26 years Marriages of colored persons. Age of persons married in 180,	-
(Table X.) Average age of persons married, 25 years, (Table) Number of times married, (Table Xl.) Denominational	3
Number of times married, (Table XI.) Denominational Nativity and residence	- 2
Nativity and residence	28-5
DEATHS	30-9
Deaths to population in 1880; proportion in each class	- 3
Deaths by sex, condition, color, nativity, parentage and locality, (Table XII.)	3
Deaths by season; numbers and proportions; recapitulation	32-3
1880	34-3
Aggregate and average age by parentage, 1880. (Table XIII.)	3
XV.)	384
(Table).	2
Percentages at different ages by parentage, 1880, (Table XVI.)	4
Percentages under five years of age by parentage, 25 years, (Table)	44-6
Percentages at different ages by parentage, 1880, (Table XVI.)	
Causes of death, sex, age, parentage and percentages, (Table XIX.)	45-6
Causes of death in connection with season, (Table XX.)	35-31
NOTES ON THE CAUSES OF DEATH IN 1880	57.8
Cholera Infantum; decedents by sex and parentage, 25 years, (Table) Consumption; sex and parentage; number of deaths at each age	61-6
Centry etatistics for 1980	6
Croup; deaths in 26 years, (Table) Diarrhoea and dysentery; recapitulation for 40 years Diphtheria; statistics for 1880	60-00
Diphtheria; statistics for 1880	- 0
Diphtheria in Providence, 23 years, (Table). Diphtheria; season and mortality compared with scarlatina and croup (Tables)	2
croup (Tables)	71-3
Heart, diseases of, decedents by sex and parentage, 25 years. (Table) Meningitis, cerebro-spinal; deaths at each age. 17 years	78-7
Pneumonia; statistics for 1880. Pneumonia; statistics for 25 years, (Table)	- 8
Pneumonia; statistics for 25 years, (Table)	83-8
Senriatina in Providence, 26 years, (Table)	8
Small-pox; recapitulatory table 25 years	8
Still-born children in 1880	89-9
Still-born children, 1856-1880	9
GENERAL SUMMARY	
(Table XXI.). Births and Deaths by parentage; excess of births, 25 years, (Table	95
	9
Census of Providence, 1880, sex, nativity, and color	9
Over Oppositions Des terro no Decrepo terror	16

CITY REGISTRAR'S REPORT.

1880.

BIRTHS.

In the year 1880, there were 2,627 children born in the city of Providence.

The following table shows how many children were born in each month and quarter of the year, in each ward of the city, and for the whole city the number of each sex, and the number of colored children born in each mouth and quarter of the year:—

TABLE I. BIRTHS, 1880. Wards, Months, Sex, and Color.

PROVIDENCE.					WA	RDS	3.				Wb	ole C	ity.	chil clud	olore dren ed in cedi:	in-
1880.	ī.	n.	ш.	ıv.	v.	VJ.	vп.	VIII	ıx.	x.	Majes.	Females.	Total.	M.	F.	Total.
January February March	25 24 31	3 4 3	13 20 23	18 7 19	94	6	19 19 21	19 22 22	43 40 42	53 81 50	104 98 130	109 99 101	218 197 281	9	8 2 2	5 4 7
lst Quarter	80	10	55	89	59	19	59	63	125	189	382	809	641	9	7	16
April	33 84 33	10 7	16 31 20	7	12 23 22	7 4 8	19	22	82 28 40	84 41 47	91 94 127	94 114 114	185 208 241		2 2 5	4 5 14
3d Quarter	100	21	57	25	56	19	63	71	100	122	812	822	634	14	9	28
July August September	27 28 32	7 3 3	11 27 29	11 10 11	28 16 15	8 8 12	24 23 22	30 17 25	31 37 28	64 49 40	128 128 107	90	218	3	8 7	7 11 11
3d Quarter	87	13	67	82	54	28	69	72	96	153	363	808	671	11	18	29
October November December	29 21 94	5 6 · 7	21 18 21	9	18 21 27	6 11 6	90	30	30 43 41	49 54 49		120		i 5	1 2 8	12 12
#h Quarter	74	18	60	82	66	28	60	82	114	152	856	825	681	17	6	22
Whole Year	841	62	289	121	285	89	251	288	485	566	1,868	1 264	2,627	51	40	91

The number of children born in Providence, in 1880, was the same as in the year 1878, and 105 more than in 1879; males, 66 more; females, 39 more. As to season, there was an increase in each quarter of the year 1880, as compared with 1879, except in the second, when the number of births was 14 less. The second, sixth and eighth wards show respectively a loss of 12, 14 and 9 in the number of children born; total loss, 35. In the other seven wards, collectively, there was an increase of 140; the greatest gain in my one ward being 50 in the seventh ward. By the United States Census, taken in June, 1880, the population of Providence was 104,857. The ratio of births for the year, therefore, was one in 39.91 of the population; or 25.05 births in each thousand inhabitants.

Season. The largest number of births in any one month of 1880 was 241 in June; the smallest, 185 in April. The largest quarterly total was 681, in the fourth quarter; the smallest, 634, in the second. The number of births in the first six months of the year was 1,275; an increase of 40 over the number in the same period of 1879; the number in the last six months of the year was 1,352; an increase of 65 over the corresponding period of the year previous. In 1880 the difference between the number in the two divisions of the year was 77; in 1879 it was 52.

The number of births in each quarter of the year 1880, and the aggregate number in the corresponding seasons, for the twenty-four years preceding, 1856–1879, and the number for the twenty-five years, 1856–1880, inclusive, were as follows:—

The following figures give the aggregate monthly totals of births in Providence for the twenty-six years, 1855–1880, inclusive; also the daily average for each month, and for the whole period:—

Births in	January.	February.	March.	April.	May.	June.
25 years, 1855-1880	8,995	8,280	4,4\$2	8,903	4,181	4,467
Dally average	4.96	5.28	5.50	5.00	5.19	5.78
Births in	July.	August.	Sept.	Oct.	Nov.	Dec.
26 years, 1855-1880	4,177	4,255	4,182	4,372	4,279	4,526
Daily average	5.18	5.28	5.36	5.42	5.49	5.62

The largest aggregate number of births in any month was in December; the smallest, in February; the highest daily average was in June; the lowest, in January. There were six months in which the daily average was above the general average of the whole period, namely: March, June, September, October, November and December; in the other months the daily average was less than the general average.

Sex and Locality. The following table will show the number of children, and the number of each sex, born in each ward, in each quarter and during the whole year, 1880:—

TABLE II. BIRTHS, 1880. Wards, Sex, and Season.

	lst	Qua	rler.	2 d	Qua	rter.	84	Qua	rter.	4th	Qua	rter.	Whole Year.		
WARDS.	М.	F.	Tota)	M.	F.	Total	M.	F.	Total	M.	F.	Total	M.	F.	Tota
I	47	83	80	54	46	100	43	44	87	43	81	74	187	154	841
II	6	4	10	8	13	21	6	7	13	9	9	18	29	83	63
ni	31	24	55	96	81	57	82	85	67	84	26	60	123	116	231
IV	19	13	82	9	16	25	21	11	82	13	19	82	63	59	12
v	28	31	59	25	81	56	84	90	54	35	81	66	122	113	233
۲I	8	11	19	11	8	19	16	12	28	18	5	23	53	36	89
VI(30	29	59	80	83	63	82	87	69	28	82	60	120	181	251
VIII	27	36	68	81	40	71	42	30	72	44	88	82	144	144	26
1 3	71	54	125	55	45	100	53	43	96	54	60	114	233	202	430
X	65	74	139	68	59	129	84	69	153	78	74	152	290	276	560
Whole City	332	809	641	812	322	634	868	308	671	856	325	681	1.863	1,264	2,627

Proportions of the Sexes. In the year 1880, there were, as is the general rule, more male than female children born in Providence. But there were six months of the year when the number of females exceeded that of the males, as may be seen by referring to Table I.; the excess in each case, however, was very small, and the excess of males in the other six months was much greater. In the eighth ward of the city the number of male and female children was equal; in the second and seventh wards there was an excess of females; in all the other wards an excess of males. The whole number of children was 2,627; 1,363 males, and 1,264 females, a proportion of 107.8 boys to 100 girls; or 51.88 boys and 48.12 girls in each 100 children born.

The number and proportion of each sex in the different quarters of the year 1880 were as follows:—

Males. Females. Males. Females. Males. Females.

Jan.—March........332 and 309, or 107.4 to 100, or 51.79 and 48.21 in each 100 born.

April—June.......312 and 322, or 96.8 to 100, or 49.21 and 50.79 in each 100 born.

July.—Sept........363 and 308, or 117.9 to 100, or 54.10 and 45.90 in each 100 born.

Oct.—Dec........356 and 325, or 109.5 to 100, or 52.28 and 47.72 in each 100 born.

Whole year.......1,363 and 1,264, or 107.8 to 100, or 51.88 and 48.12 in each 100 born-

In the aggregate for the twenty-five years, 1856–1880, inclusive the number and proportion of each sex in each quarter of the year were as follows. The whole number of children born during this period was 49,049; males, 25,213; females, 23,836:—

Males. Females. Males. Females. Males. Females.

Whole period...25,213 and 23,836, or 105.8 to 100, or 51.40 and 48.60 in each 100 born-

In the twenty-seven years, 1854-1880, inclusive, the number of children born in Providence was 52,018; males, 26,731; females, 25,287. The number and proportion of each sex in each year and in the whole period were as follows:

```
Males.
                    Females. Males. Females. Males. Females.
1854.... 711 and
                     658; or 108.0 to 100; or 51.91 and 48.09 in each 100 born.
                     793; or 101.7 to 100; or 50.41 and 49.56 in each 100 born.
1353..... 807 and
1856..... 891 and
                     784; or 113.6 to 100; or 53.19 and 46.81 in each 100 born.
1857..... 873 and
                     816; or 106.8 to 100; or 51.66 and 48.84 in each 100 born.
1858...... 890 and
                     834; or 106 7 to 100; or 51.62 and 48,88 in each 100 born.
1859..... 825 and
                     768; or 107.4 to 100; or 51.79 and 48.21 in each 100 born.
1880. ..... 809 and
                     889; or 98.4 to 100; or 49.09 and 50.91 in each 100 born.
                     789; or 118.6 to 100; or 54.26 and 45.74 in each 100 born.
1861..... 985 and
                     738; or 107.2 to 100; or 51.78 and 48.27 in each 100 born.
1862..... 791 and
1863..... 699 and
                     618; or 111.4 to 100; or 52.71 and 47.39 in each 100 born.
1864..... 663 and
                     681; or 97.8 to 100; or 49.83 and 50.67 in each 100 born.
1865...... 691 and
                     607; or 113.8 to 100; or 53.24 and 46.76 in each 100 born.
1894..... 849 and
                     783; or 108.4 to 100; or 52.02 and 47.98 in each 100 born.
1867..... 830 and
                     794; or 104.5 to 100; or 51.11 and 48.89 in each 100 born.
1868...... 944 and
                     922; or 102.4 to 100; or 50.59 and 49.41 in each 100 born.
                     874; or 108.1 to 100; or 51.95 and 48.05 in each 100 born.
1869..... 945 and
1870....... 987 and 922; or 104.8 to 100; or 51.19 and 48.81 in each 100 born.
1271...... 956 and 1,004; or 95.6 to 100; or 48.78 and 51.22 in each 100 born.
1872............1,079 and 1 127; or 95.7 to 100; or 48.91 and 51.09 in each 100 born.
1874............1,408 and 1,372; or 102.6 to 100; or 50.65 and 49.35 in each 100 born.
1876................1,359 and 1,272; or 106.8 to 100; or 51 65 and 48.85 in each 100 born.
1878.......1,356 and 1,269; or 107.0 to 100; or 51.69 and 48.51 in each 100 born.
27 years .. .....26,731 and 25,287; or 105.7 to 100; or 51.89 and 48.61 in each 100 born.
```

Color. There were 91 colored children born in the city, during the year 1880, which was 13 less than in 1879, and is less than in any year since 1871, when the number was the same as in 1880. By the United States Census of June, 1880, the colored population of Providence numbered 3,646; the ratio of births to population in this class was, therefore, one birth in 40.07 of the colored inhabitants of the city; or 24.96 in each thousand persons. On page 2 it was stated that the ratio of the whole number of births to the total population of the city was one in 39.91; or 25.05 births in each thousand. The white population of the city was 101,211; the number of white children born in 1880 was 2,536; the ratio for this class, therefore,

was one birth in 39.91 of the white inhabitants; or 25.06 births in each thousand. It will be seen that the difference between the two classes is very slight; but it is quite probable that the number of the colored population should be considerably larger than it is given, owing to omissions in marking the color on the census returns, and this would reduce somewhat the ratio of births in that class. The births of colored children in 1880 were in the different wards as follows: First ward, 8; Second, 13; Third, 8; Fourth, 4; Fifth, 2; Sixth, 9; Seventh, 12; Eighth, 8; Ninth, 27; Tenth, 0. The number in each month was: January, 5; February, 4; March, 7; April, 4; May, 5; June, 14; July, 7; August, 11; September, 11; October, 4; November, 7; December, 12; Total, 91.

Of these 91 colored children, 51 were males and 40 females; a proportion of 127.5 males to 100 females; or 56.04 males and 43.96 females in each 100 children born.

The following table shows the number of colored children, and the number and proportion of each sex, born in each of the 25 years, 1856-1880, inclusive, and the number for the whole period:—

Year.	Total. M	fales. Fer	nales. Males. Females. Males. Females.
1856	50	26 and	24; or 108.3 to 100; or 52.00 and 48.00 in each 100 born-
1857	31	17 and	14; or 121.4 to 100; or 54.84 and 45.16 in each 100 born.
1858	51	25 and	26; or 96.2 to 100; or 49 02 and 50.98 in each 100 born.
1859	36	20 and	16; or 125.0 to 100; or 55.56 and 44.44 in each 100 born.
1860	51	31 and	23; or 134.8 to 100; or 57.41 and 42.59 in each 100 born.
1861	51	28 and	23; or 121.7 to 100; or 54.90 and 45.10 in each 100 born.
1862	61	26 and	35; or 74 3 to 100; or 42.62 and 57.38 in each 100 born.
1863	32	16 and	16; or 100.0 to 100; or 50.00 and 50 00 in each 100 born.
1864	38	25 and	13; or 192.3 to 100; or 65.79 and 34.21 in each 100 born
1865	46	27 and	19; or 142.1 to 100; or 58.70 and 41.30 in each 100 born.
1866	67	39 and	28; or 139.3 to 100; or 58.21 and 41.79 in each 100 born.
1867	75	38 and	37; or 102.7 to 100; or 50.67 and 49.33 in each 100 borns
1868	90	49 and	41; or 119.5 to 100; or 54.44 and 45.56 in each 100 born.
1869	85	51 and	34; or 150.0 to 100; or 60.00 and 40.00 in each 100 born.
1870	104	56 and	48; or 116.7 to 100; or 53.85 and 46.15 in each 100 born.
1871	91	50 and	41; or 121.9 to 100; or 54.95 and 45.05 in each 100 born.
1872	91	51 and	48; or 106.2 to 100; or 51.52 and 48.48 in each 100 bors.
1873	112	62 and	50; or 124.0 to 100; or 55.36 and 44.64 in each 100 born.
1874	109	55 and	54; or 101.9 to 100; or 50.46 and 49.54 in each 100 bors.
1875	101	49 and	52; or 94.2 to 100; or 48.51 and 51.49 in each 100 born.
1876	108	48 and	60; or 80.0 to 100; or 44.44 and 55.56 in each 100 born-
1877	104	60 and	44; or 136.4 to 100; or 57.69 and 42.31 in each 100 born.
1878	104	52 and	52; or 100.0 to 100; or 50.00 and 50.00 in each 100 born.
1879	104	58 and	46; or 126.1 to 100; or 55.77 and 44.23 in each 100 born
1880	91	51 and	40; or 127.5 to 100; or 56.04 and 43.96 in each 100 born.
25 years1	.8941.	010 and 8	84; or 114.3 to 100; or 53.33 and 46.67 in each 100 born.

The births, marriages and deaths of colored people in Providence in the 25 years, 1856-1880, inclusive, have been as follows:—

Twenty-four years, 1856-18791	,803 births	871 marriages	1,899 deaths.
1880	91 births	57 marriages	96 deaths.
Twenty-five years, 1856-18801	,894 births	928 marriages	 1.995 deaths.

In 1880, the proportion of deaths to births in this class was 105.5 to 100; in the 25 years, 1856-1880, it was 105.3 deaths to 100 births.

PARENTAGE.

The variety of nationalities represented in the population of the city makes the subject of the parentage of the children born, one of interest and importance.

Table III. gives the number of children, according to parentage, born in each ward of the city; and the number and percentage of each nationality in the whole city. Where the parents of a child were of different foreign nations, the parentage is given according to the birth-place of the father:—

TABLE III. BIRTHS, 1880. Wards and Parentage.

PARENTAGE.				1	VAI	RDS					City.	h 100
PARENTAGE.	ı.	11.	ш.	ıv.	v.	VI.	VII.	viii	ıx.	x.	Whole City	In each 10 born ther were:-
American	130	44	66	52	78	56	126	146	210	140	1,048	39.90
Irish	105	3	71	26	71	7	55	58	104	187	687	26,15
English, Scotch and Welsh	21		7	5	20	5	9	20	26	84	197	7.50
German	5	2	4	1	6		1	3	12	10	44	1.67
British American	14	1	6	2	20	1	6	10	13	13	86	3.27
Portuguese			36		2		1	1	1	1	42	1.60
Other Foreign	6			7	5	3	4	1	4	5	35	1.33
Total Foreign	151	6	124	41	124	16	76	93	160	300	1,091	41.52
American father and Foreign mother	32	6	16	10	16	6	15	25	26	52	204	7.77
Foreign father and American mother	28	6	33	18	17	11	34	24	39	74	284	10.81
Ward totals	341	62	239	121	235	86	251	288	435	566	2,627	100.00

The "other foreign" in this table were as follows: Italian, 13; Swedish, 8; French and Polish, 4 each; Russian, Swiss, Belgian, Austrian, Norwegian, and San Domingo, 1 each.

The foreign element seems to be in excess in the First, Third, Fifth and Tenth Wards; the number of children born of foreign parents in those wards being larger than the number born of American parents; and in the Third and Tenth Wards the number of children of Irish parents alone exceeded the number born of American parents. The number of children of American fathers was 1,252, or 47.67 per cent. of the whole number born; this was 24 more children, but 1.02 per cent. less than the number of this class in 1879.

The following recapitulatory table gives the number and percentage of children of American, foreign and mixed parentage, in each of the twenty-five years, 1856-1880, inclusive; and the number for the whole period:—

	American.	Foreign.	Amer. father. For, mother.	For. father.
Year.		Tumber. Per cent. N		
856	70542.09	. 85751.17	553 22	588.5
857	65038.51	. 93755.51	513 02	
858	71241 30	. 88651.39	583.36	683.
859	599 37.60	. 88755.69	593.70	
860	66140.11	. 85752.00	694.19	
861	65437.91	91653.10	864.99	
862	58938.52	. 79952.26	744.84	
863	46735.73	. 71154.40	624.74	
864	52038.69	. 69952.00	604.47	65(=
865	47636.67	68452.70	705.39	68
866	65640.20	. 82150.30	704 29	955.=
867	64539.72	. 79148.70	804 93	1086.
868	77941.75	. 86846.51	1035.52	116 8.=
869	82245.19	. 77642.16	1045.72	1176.
870	80042.35	. 862 45.64	904.76	1377.=
871	87944.85	. 82842.24	1015.15	1527.
872	90741.12	.1,00245.42	1285.80	1697.
873	869 40.84	. 94144.22	1346.29	
874	1,06838.42	47.70	1495,36	
875	99937.50	,25647.15	1816.79	. 2288
876	1,00738.27	1,23046.75	1626.16	2328
877	1,05441.45	.1,06741.96	1817.23	2389_
878	1,03539.40	1,17444.69	1806.85	2389_
	1,02940.80			
880	1.04839.90	,09141.52	2047.77	28410
	ds.19,63040.02			

The number of children in each class is larger than in 1879; but of the children of American parentage the percentage is slightly less; while of those of foreign parentage it is a little larger than in the previous year. Of the children of American fathers and foreign mothers the percentage was less in 1880 than in 1879, but larger than in any other year; and of children of foreign fathers and American mothers the percentage was larger than in any previous year of the record.

The following table shows the percentage of children born in Providence, of parents of each nation, for each of the twenty-five years, 1856-1880:—

	American.	Irish.	English,	German.	British	Other Portuguese. for'n.
	Per cent.	Per cent.				Per cent. Per ct.
×56						
						1.54
						1.89
						1.5
						1.6
						1.9
						1.04
						1.9
						1.8
						1.90
						2.8
						22
						1.0
						1.8
						9
						1.78
						2.8
						2.10
						1.221.19
						1.221 24
						1.101.80
						1.181.71
						1.601.8

The British-American were included in the "other foreign" until 1868, and the Portuguese until 1874. The highest percentage of

American parentage in any year was 45.19 in 1869; the lowest, 35.73 in 1863. The highest percentage of Irish parentage was 45.38 in 1859; the lowest, 26.15 in 1880. The percentage of English and Scotch parentage in 1880 was larger than in any year since 1876; that of German parentage was smaller than in any year since 1871; the percentages of British-American and Portuguese parentage were larger than in any previous year.

More exact statistics as to the parentage of the children bon in Providence in 1880, are given in Table IV. which shows the birth-places of both parents:—

	BI	RTH-	PLAC	es o	F MC	THE	ts.	4
BIRTH-PLACES OF FATHERS.	United States.	Ireland.	England, Scotland and Wales,	Germany.	British America.	Portugal and Western Islands.	Other Countries.	to the father's birt
United States	1,048	107	52	4	37		4	1,990
Ireland	147	629	37		18		3	884
England, Scotland and Wales	68	38	144		12		3	265
Germany	15	7	2	32			3	.50
British America	36	15	5		66			122
Portugal and Western Islands	4	1				41		40
Other Countries	14	7		1	2		25	0
Whole number according to the mother's birth-place	1,332	804	240	87	135	41	38	2.02

TABLE IV. BIRTHS, 1880. Parentage.

Of the 2,627 children born in Providence in 1880, 1,048, or 39.90 per cent. of all, were of purely American parentage; this was 19 more children, but 0.90 per cent. less than in the previous year. There were also 204 children with American fathers and foreign mothers, which made 1,252 children, or 47.67 per cent. of all, who were children of American fathers. There were 284 children of

American mothers and foreign fathers; this gives a total of 1,332 children of American mothers, which was 50.71 per cent. of the whole number. Of the two classes of children of mixed parentage (that is where one parent was American and one foreign) the number was 488, or 18.58 per cent. of all. Of children of purely foreign parentage the number was 1,091, or 41.52 per cent. of all; this was 56 more children, and 0.48 per cent. more than in 1879. Of these 1,091 children there were 937 whose parents were natives of the same foreign country; this was 85.88 per cent. of the number of children of foreign parentage, and 35.66 per cent. of the whole number of children born during the year. There were 154 children whose parents were natives of different foreign countries; this was 14.12 per cent. of the 1,091 children of foreign parentage, and 5.86 per cent. of the 2,627 children born during the year.

NUMBER OF THE CHILD.

Table V. shows how many of the 2,627 children born during the year 1880 were the first, second, third, fourth, &c., children of their mothers; also the same statistics as to the 48,022 children born in the twenty-five years preceding, 1855–1879, inclusive:—

TABLE V.	Births,	1880.	Number	of	the	Child.
----------	---------	-------	--------	----	-----	--------

NUME	BER	of	CHILD.	1880.	25 years, 1855–1879.	NUME	ER	of	спі	LD.	1880.	25 years. 1855–1879.
First ch	dld	of th	e mother	612	11,824	Troclfth	chil	d of t	he n	otb'r	6	194
Second	44	44	**	467	9,895	T hirtees	ith '	• "		**	8	79
Third	**	64	66	424	7,689	Fourteen	th '	46		•4	5	85
Fourth	••	**	44	887	5,782	Fifteenti		• ••		44		16
Fifth	66	40	••	254	4,186	Sixteent	h '	• ••		44	8	a
Sixth	**	44	46	180	8,044	Seventee	n/h '	4 66		44		7
Sevensk	14	••	**	181	2,134	Eighteen	th '	• ••		66		1
Eighth	••	46	44	82	1,365	Nineteen	th '	• ••		44		4
Ninth	"	**	64	56	885		—.					
Tenth	**	"	••	39	572	Whole	nur	nber	of	chil-		4
Eleventi		14	44	96	307	dren.	•••	• • • • •	• • • • •	•••••	2,627	48,022

Of the 2,627 children born in 1880, 612 or 23.30 per cent, were the first children their mothers had borne; this was 1.06 per cent, more than the same class of children in 1879. Of the 48.022 children born in the twenty-five years, 1855–1879, 11.824 or 24.62 per cent, were the first children of their mothers.

Average number of children to each mother. The number of twin births in 1880 was 27; the whole number of mothers in 1880, was, therefore, 2,600. Table VII. shows that these mothers had borne, in all, 9,386 children, giving an average of 3.61 children to each mother.

The next table shows the average number of children to each mother, in each of the twenty-six years, 1855 to 1880, inclusive. This table shows the number of children the mothers included in the figures had borne, previous to the end of the year 1880; but it does not show the average number of children the same mothers may have in their life-time:

Table VI. Births, 1855 to 1880.

Average number of children to each mother.

	No. of	No. of	Aggregate
Years.	Mothers.	Children. No	o. of children. Average.
1855	1,588	1,600	5,2143.28 children to each mothe
856	1,658	1,675	5,569
857	1,664	, 1,688	5,7363.44 children to each mothe
1858	1,706	1,794	5,6853.83 children to each mothe
is 59	1,570	1,593	5,6153.58 children to each mothe
1880	1,627	1,648	5,8418.59 children to each mother
861	1,694	1,725	6,0173.55 children to each mother
1863	1,512	1.529	5,322 8.52 children to each mothe
1963	1 296	1,807	4,717 8.64 children to each mothe
1864	1,327	1,344	4,887 8.68 children to each mothe
1865	1,279	1,298	4,6258.60 children to each mothe
1996	1,614	1,682	5,7448.56 children to each mothe
1867	1,595	1.624	5,4978.50 children to each mothe
1868	1,841	1.866	6,338 3.44 children to each mothe
1969	1,797	1,819	5,954 8.30 children to each mothe
1870	1.859	1,889	6,1458.31 children to each mothe
1871	1,937	1.960	6,420
1872	2,174	2,206	7,346 8.38 children to each mothe
1873	2,104	2,128	6,7818.20 children to each mothe
1874	2,788	2,780	8,960 8.27 children to each mothe
1875	2.634	2,064	8,9778.41 children to each mothe
1876	2,604	9,631	8.9143.42 children to each mothe
1877	2,516	2,543	8,7058.46 children to each mothe
1878	2,597	2,627	9,417 8.63 children to each mothe
1879	2,499	2,523	8,943
***	9 200	9 697	9,386 3.61 children to each mothe

Table VII. gives the number of the child, in connection with the age of the mother; shows how many first, second, &c., children were born in 1880 to mothers at the respective ages given; the number of mothers at each age; and the aggregate and average number of children they had borne:—

Table VII. Births, 1880. Number of child with age of mother.

				N	UMB	ER (OF C	HIL	D,							TS BE	ui e	y had	of series
MOTHERS IN YEARS.	lst.	gd.	3d.	4th.	5th.	6th.	7th.	sth.	9th.	10th.	lith.	19th.	lsth.	14tb.	16th.	Number of mothers each age.	Children they bore 1880.	Fotal children they had borne,	Average number of children to mothers at each age.
ŧ	1	24	80	-		9	_	- 00	6	=	-		=	=		1	1	2	1.00
6	9			•											ij	2		4	
7	3	1														4	4	1	
8	18	3														21	21	24	
9	21	12	199											ii		35		1 0	
0	*46		4)												73	75	1 25	
1	41	13	3								00					58	100	1	
2	66			5	3									-		128		1 . 3	
3	65		19	9												132	100	236	3
4	60		29	8	2	1	1				M			ij		158	159		100
5	61	55	100	*16		2	1									176			100
6	*47	35	*27	22	7	2	1	1								140	1000	344	
7	48	40	*27	32	11	6	4			1						168	169	46	100
8	32	*28	162	46	22	9										197	201	625	
9	19		25	23	1000	14	7	3	1							134	135	499	100
0	27	30	43	40	1000	*14	7	4	3	1881						209	210		
1	8	17	1 20	19		6	6	3		li	1					90	90		
2	13		0.00	18	22	11	15	000	100	3	100		1			123	124	569	
3	*9	11	*24	23	16	6	9	5	100	100						107	109	100	
4	5	7	18	17	14	21	13			183	ï	200				101	101	481	
5	5	7	15	18	15	+27	*20	12			9					127	129	620	
6	4	5		13	14	7	*11	*10		-	3					84	87	467	
7	2	5	6	6	9	15	10000	*9			9					76		457	
8	1	5	100	7	9	11	*5	14		*8	3	1		1		76			
9	4	4	4	5	4	4	6	3		*8	3			ı		50		320	
0	1	9	1 90	4	*11	14	5	6	m		4			9		64	65	415	
1	1	-	1	1	3	1	1	9		103	3		-	-		15		112	
			1	2	9	3	2	2			3	1	1	2		26	15 26		
3	****			1	2	9	3		9		1	-	1	-		10	10	71	7.10
4				1		1	1		1				1		1	5	5	49	
5					1	1	10		1	1		,			-	4		33	
6	1					1						1				2	4	7	3.50
8										1	-					1			10.130
0						1				1						1	1	8	60
ge not						1			1				1	1		-	1	1	.0
stated	2			1,3	100											2	2	2	-
Statedinii		***		-						-		-	-			-			7
otals	612	467	424	337	254	180	134	82	55	39	26	6	3	5	3	2,600	2,627	9,386	-
	1		200			-	1	‡ T					100				-		1 3

The reference marks in this table show the age of the mother and the number of children in each case of plurality-birth; thus at the age of 28, four mothers bore twins; in one case they were the second and third children, and in three cases they were the third and fourth children; at the age of 38, two mothers bore twins; in one case, the seventh and eighth children; in one, the tenth and eleventh.

In Table VII. for 1879, as printed in the report for that year, there were several errors; thus the total number was wrong of children borne by mothers at the following ages: 21 years, 23, 24, 27, 28, 33, 34, 35, 36, and 39; consequently the average number of children, borne by mothers at these ages, was also wrong. The whole number of children borne by mothers of all ages should have been 8,943, instead of 9,039; but the general average of the whole, 3.58, was right. There was also a typographical error in the number of children borne, in 1879, by mothers at the age of 34; instead of 04, it should have been 104.

There was one woman of 50 years of age who bore a child in 1880, having had five children before. The number of women 40 years old and over who bore children in 1880 was 128, the total number of children borne by them was 930, an average of 7.27 children to each mother. In the seventeen years and six months during which the record has been kept of the ages of the mothers, there have been 1,797 mothers of 40 years and over who have borne 13,632 children; an average of 7.59 children to each mother.

During this period, from July 1, 1863, to December 31, 1880, the number of children borne by mothers whose age was given was 36,722, the number of children borne by mothers of each age during this time is given in the following table:—

Age of mother.	Children.	Age of mother.	Children.	Age of mother. (Children.
14	5	27	2,210	40	854
15	19	28	2,588	41	238
16	59	29	1,847	42	288
17	151	80	2,963	43	157
18	897	81	1,285	44	102
39	641	32	1,785	45	106
20	1,049	88	1,451	46	. 29
21	1,199	84	1,239	47	24
23	1,788	3 5	1,842	48	12
23	1,961	36	1,106	49	. 8
24	2,187	87	907	5 0	. 9
35	2,466	3 8	981		
26,	2,256	89	625	All known ages,	86,722

The mothers of these 36,722 children were divided as follows as to age:—

Under twenty years	
Twenty, and under twenty-five	
Twenty-five, and under thirty	
Thirty, and under thirty-five	
Thirty-five, and under forty	,
Forty, and under forty-five	1,684
Forty-five, and over	
Total	

Of these 36,722 children, 20,811, or 56.67 per cent., were borned by mothers less than thirty years of age; 14,094, or 38.38 per cent., by mothers between thirty and forty years of age; and 1,817, or 4.95 per cent., by mothers forty years old and over.

PLURALITY BIRTHS.

There were, in 1880, 27 cases of plurality births in Providence: this was four more than in 1879.

The following are the statistics of these cases:-

Cases, 27; number of children, 54.

Locality. First ward, 5; second, 1; third, 2; fourth, none; 2; sixth, 1; seventh, 3; eighth, none; ninth, 4; tenth, 9.

Season. January, 3; February, 1; March, 3; April, 2; May. 2; June, 5; July, 3; August, 1; September, 4; October, none; November, 2; December, 1.

Sex. In eleven cases both children were males; in eight both were females; in eight there was one child of each sex; in all, thirty male and twenty-four female children.

Color. The twins were children of white parents in every case but one.

Number of the Children and Age of the Mother. By means of the reference marks in Table VII. may be found the age of the mother and number of the children in each case. In three cases, they were the first and second children, of mothers respectively 20, 26, and

33 years old; in four cases, the second and third children of mothers, 20, 24, 28, and 32 years old; in seven cases, the third and fourth children of mothers, three of whom were 28 years old, and the others 25, 26, 27, and 33; in one case the fourth and fifth children of a mother 25 years old; in two cases, the fifth and sixth children of mothers 29 and 40 years old; in two, the sixth and seventh children of mothers 30 and 35 years old; in three, the seventh and eighth children of mothers 35, 36, and 38 years old; in two, the eighth and ninth children of mothers 36 and 37 years old; in one, the ninth and tenth children of a mother 36 years old; in two cases, the tenth and eleventh children of mothers 38 and 39 years old. There were thirteen cases where the mothers were less than thirty years old.

Parentage. In eight cases both parents were American; in fourteen both were Irish; in one both were English; in two both were Portuguese; in one the father was English and the mother American; and in one the father American and the mother British American.

TWENTY-SIX YEARS' SUMMARY OF PLURALITY BIRTHS.

In twenty-six years, 1855–1880 inclusive, the number of cases of plurality births in Providence has been 610; 601 of twins, and 9 of triplets; the whole number of children in these cases being 1,229.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children and one female; making 9 male and 18 female children in the nine cases.

Of the 601 cases of twins, there were 197 in which both of the children were males; 206 in which both were females; and 198 in which there was one male and one female child; total, 592 males and 610 females. Of the 1,229 children of all the cases of plurality births, there were 601 males, and 628 females; a proportion of 104.5 females to 100 males; or 51.10 girls and 48.90 boys in each 100.

Colon In all the cases of triplets the parents were white. Of the 601 cases of twins, in 579 the parents were white, and in 22 they were colored.

Parentage. Of the triplets, in one case, the parents were both American; in five cases, both were Irish; in one case, both German;

in one, the father was English, and the mother Irish; and in one the father Scotch, and the mother Irish.

In the following table for the 601 cases of twin-births, the nativity of both parents is given; the first letter representing the father's birth-place, and the second the mother's.

In this table, some errors found in the table given in the report for 1879, have been corrected in adding the figures for 1880, so that the totals are correct for the twenty-six years.

A., American; I., Irish; E., English; S., Scotch; W., Welsh: G., German: B. A., British American; D., Dutch; P., Portugues: It., Italian; Sw., Swedish.

Parentage. No.	Parentage. No.	Parentage. No.	Parentage. Na.
	I. and A14		l
I. and I 235 E. and E 29	1	B. A. and I1 Sw. and I1	
	B. A. and A 6 G. and A 3		
B. A. and B. A 10	Sw. and A 1	A. and E	I. and B.A
D. and D 1 P. and P 9	E. and I11	I. and E	F. and B. A 1 S. and B. A 1
It. and It 1	S. and I 3	G. and E1	Total

From these figures we see that of the 601 cases of twin-birds, there were 236 in which the mothers were natives of the United States; this was 39.27 per cent. of the whole number; in 267 cases, or 44.43 per cent., the mothers were Irish.

Of the 610 cases of plurality births in the twenty-six years, the American mothers numbered 237; the foreign mothers, 373. The whole number of American mothers bearing children during this period was 23,496; the 237 mothers in cases of plurality births were, therefore. 1.01 per cent. of this number; the whole number of foreign mothers who bore children during the same period was 26,534: so that the 373 foreign mothers in cases of plurality births were 1.41 per cent. of all the foreign mothers.

Proportion of plurality births to whole number. The following figures show the number and proportion of plurality births, as compared with the whole number of births, for the year 1880, and for the twenty-five years previous, 1855–1879; also for the whole period of twenty-six years, 1855–1880, inclusive:—

Single Births. Twin. Triplet.	Whole No. Whole No.
25 years, 1855-187946,8475749	47,43048,022
1880 2 573 270	2,600 2,627
26 years' totals	50,03050,649
Proportion of cases of plurality births:-	
Twins.	· Triplets.
1855-18791 in 82.6	1 in 5,270.0
1880 1 in 96.8	•••••
1855–1880 1 in 83.2	1 in 5,558.8
Proportion of children of plurality births	;
Twins.	Triplets.
1855–1879 1 in 41.8	
18801 in 48.6	•••••
1855–18801 in 42.1	1 in 1,875.9
Proportion of plurality births by parentage.	. The number of births
in the twenty-six years, 1855-1880, inclusiv	
lowing figures show the number of births o	•
mixed parentage, and the number and pro	
in each class:—	F,
Whole No. Births. Plu	•
American parents20,069	199100.8 or 0.99

Of the 610 cases, the 199 of purely American parentage were 32.62 per cent.; the 350 of foreign parentage were 57.38 per cent.; the 23 with American father and foreign mother, 3.77 per cent.; and the 38 with foreign father and American mother, 6.23 per cent.

 Foreign parents
 23,798
 350
 67.9 or 1.47

 Am. father, foreign mother
 2,736
 23
 118.9 or 0.84

 Foreign father, Am. mother
 3.427
 38
 90.2 or 1.11

 Total
 50,030
 610
 82.0 or 1.22

WARRIA EN.

There were 1.272 marriages in Providence in the year 1889).

Table VIII. shows the number in each month and quarter of the year 1989, and in each of the six years preceding; also the aggregate and average number for each month and quarter of the twenty-six years, 1955–1989, helicite:—

TABLE VIII. MARRIAGES. SHEEDIR.

MOSTHS.	1986	1473	L:3	W	1578.	D575.	MAN.	M veer. Diss-Diss.	
Jacany)45	303	70	•	3	3	*6	1,500	=
February	300	85	73	65	-	; 3	8.	rin	
March	58	47	3	38	-	า	56	1,110	•
st Quarter	25-8	234	223	177	236	225	234	4.797	1%
A pril	98	83		83	78	84	95	1,731	•
Kay)48	96	68	85	80	82	113	1,700	1 7
June	111	96	104	83	94](C	101	1,960	
d Quarter	\$18	211	243	251	244	200	300	5,500	21
Saly	97	50	94	85	56	101	88	1,675	. 6
August	75	86	•	76	62	73	80	1.444	. 5
September	115	105	95	87	96	111.	129	2,024	. 7
d Quarter	267	253	249	248	211	285	297	5,143	19
October	106	108	99	99	95	107	127	2,239	8
November	148	128	115	102	100	114	114	2,438	. 9
December	113	76	83	83	74	89	98	1,789	
th Quarter	369	307	296	283	305		334	6,456	34
Tearly totals	1 989	1 071	1.016	953			1,176	21,919	84

The number of marriages in 1880 is the largest ever recorded in the city for one year; it is 161 more than in 1879, and is 389 more than the average number for twenty-six years. As the population of the city by the Census in 1880 was 104,857, the 2,464 persons married during the year give a ratio of one in each 42.55 of the population; or 23.50 persons married in each thousand inhabitants. The ratio of persons married to births was 93.80 to 100; of persons married to deaths 118.5 to 100. In 1879 the ratio of persons married to children born was 85 to 100; of persons married to deaths, 105.7 to 100.

Season. According to the rule which has generally prevailed, the larger number of these marriages occurred in the last six months of the year, the largest quarterly total being 369 in the fourth quarter; the smallest quarterly total was 258 in the first quarter. The largest number in any one month was 148 in November; the smallest, 53, in March. From the aggregate number of marriages for twenty-six years, as given in Table VIII., it seems that of each 100 marriages about 25 occur in the winter months of December, January and February; 22 in the spring months; 23 in the summer; and 30 in the autumn.

The number and percentage of marriages in each quarter of the year 1880 are given below; also the corresponding figures for the period of twenty-five years, 1855-1879, inclusive:—

18	80.	1855-1879.			
Marriages.	Per Cent.	Marriages.	Per Cent		
258	20.94	4,469	21.60		
318	25.81	5,275	25.50		
287	23.80	4,856	23.47		
389	29.95	6,087	29.48		
1,282	100.00		100.00		
	Marriages 258 318 287 389				

In Table IX. are given the birth-places of the brides and bride-grooms, in each of the 1,232 marriages of the year 1880:—

1

TABLE IX. MARRIAGES, 1880. Nativity of the Parties.

	BIRTH-PLACES OF BRIDES.							
BIRTH-PLACES OF GROOMS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands,	Other Countries.	Whole number of the
United States	750	48	25	2	20		4	M
Ireland	44	119	7		7			177
England, Scotland, and Wales	47	10	40		5			100
Germany	12	1	1	4			1	19
British America	27	3	4		20			M
Portugal and Western Islands	3					4		1
Other Countries	7	3			3	1	10	3
Whole number of Brides	890	184	77	6	55	5	13	1,22

Bride and Groom born in the same country. In 947 marriages, of 76.87 per cent. of the whole number, the bride and groom were natives of the same country; this is 116 more marriages than were found in this class in 1879, but the per cent. is 0.72 less. There were 750 marriages, or 60.88 per cent. of all, in which both parties were natives of the United States; this is 97 more marriages that there were of this class in the preceding year, and 0.08 per cent. less. Of marriages where both parties were natives of Ireland the number was 119, or 9.65 per cent. of all; this was 6 less marriages, and 2.02 per cent. less than in 1879. There were 40 marriages where both persons were English or Scotch; 13 more than in 1879; and 20 where both were British American, or 11 more than in the previous year.

Bride and Groom born in different countries. Every year this table shows some curious facts as to the intermarriage of the different nations represented in our population. There were 285 marriages or 23.13 per cent. of all, in which the persons were of different

nationalities; this is 45 more than in 1879, and 0.72 per cent. less. Of these 285 marriages, there were 99, or 34.73 per cent., in which the groom was born in the United States; and 140 others, or 49.12 per cent., in which the bride was born in the United States. The 99 grooms born in the United States, who married women of other countries, were 11.66 per cent. of the whole number of grooms (849) who were natives of this country; of the 177 grooms born in Ireland, 58. or 32.77 per cent., married women of other countries than their own; of the 102 born in England or Scotland, 62, or 60.79 per cent.; of those born in Germany, 78.95 per cent.; of those born in British America, 62.96 per cent. Of the 890 brides born in the United States, 140, or 15.73 per cent., married men of other countries; of the 184 born in Ireland, 65, or 35.33 per cent.; of those born in England or Scotland, 48.05 per cent.; of those born in Germany, 33.33 per cent; of those born in British America, 63.63 per cent.

The 24 grooms born in "other countries" than those specified in the table, were natives of the following countries: Italy, 9; Sweden, 8; France 3; Poland, 2; Brazil and British West Indies, 1 each.

The 15 brides born in "other countries" were natives of the following countries: Sweden, 6; Switzerland and Italy, 2 each; Austria, Poland, Cuba, South America and Sandwich Islands, 1 each.

The number and percentage of American, foreign and mixed marriages in each of the twenty-six years from 1855–1880, inclusive, are given in the following table. By mixed marriages are understood those where one of the parties was of American, and the other of foreign nativity:—

	Ame	rican.	For	elgn.	Mixed.
Year.	Number.	Per cent.	Number.	Per cent.	Number. Per o
1855	. 308	48.73	265	41.93	59
1856	. 852	54.74	919	32.97	791
1857	. 384	62.54	175	28.50	55
1858	. 283	50 54	214	.8 8.21	63I
1859	. 333	54.50	224	36.66	54
1860	836	54.19	225	86.29	59
1861	. 804	55 58	182	83 27	61
1862	. 288	52.65	179	83.73	801
1863	. 849	56.07	192	81 47	76
1864	. 427	57.78	211	28.55	101
1865	. 417	58.82	902	28.49	90
1966	. 471	58.00	244	80.05	97
1867	. 485	56.73	269	81.46	101
1868	. 530	60.78	215	24.66	127
1869	. 530	56.99	286	80.75	114
1870	. 552	57.56	262	27.82	145
1871	. 523	55.46	278	29.48	143
1872	. 591	54.52	805	28.14	188
1873	. 652	56.69	817	27.57	191
1874	. 648	55.10	346	29.42	
1875	. 586	51.31	820	29 66	
1876	. 547	57,64	241	25.39	
1877	. 545	57.19	233	24.45	
1878	. 601	59.15	245	94.13	
1879				20.26	
1880	. 750	60.88	243	19.72	239

The proportion of foreign marriages has steadily decreased for a number of years, and in 1880 was smaller than ever before; on the other hand, the proportion of mixed marriages has increased, and was larger in 1880 than in any previous year. The proportion of American marriages, though slightly less than in 1879, was larger than any other year of the record except 1857, when it was exceptionally large. The whole number of marriages in this table is 21,913; in Table VIII., the total number for the same period was given at 21,919. This discrepancy is owing to omissions in some of the early years of the records, and is explained in the report for 1879, page 22. There are almost every year reports of marriages brought in too late.

to be reckoned in making up these tables; they are afterwards added to the numbers for each year as given in Table XXI. at the end of the book, and the true number of marriages for the twenty-six years, 1855–1880, as given there, is 21,950.

Marriages of Colored Persons. There were 57 marriages, in 1880, among the colored people of the city; this was 14 more than in 1879, and is more than in any year since 1874. The aggregate and average age of the persons married in 1880 were as follows:—

57 malesAyerage age, 31.16 yearsAverage age, 31.16 y	years.
57 femalesAggregate age, 1,444 yearsAverage age, 25.33 y	ears.

In the twenty-five years, 1856-1880, inclusive, the number of marriages of colored persons in Providence was 928; the aggregate and average age of the persons as follows:—

928 males Aggregate age,	, 27,342 yearsAverage ag	e, 29.46 years.
928 femalesAggregate age	, 24,299 years Average ag	e, 26.18 years.

Age of persons married. Table X. gives the number and percentage of brides and grooms in each division of ages:—

TABLE X. MARRIAGES, 1880. Ages of the Parties.

	AGES OF THE BRIDES.							ber .	بدة
AGES OF THE GROOMS.	Under 20 years.	20 to 25.	25 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	Whole number of Grooms.	Percentage Grooms at each age.
Under 30 years	17	4				 .		21	1.70
29 to 25	106	250	87	4				897	82.2
95 to 30	44	218	98	94	•••••			879	80.76
20 to 40	14	82	110	80	11			297	24.30
40 to 50	2	8	14	48	12	1		85	6.90
50 to 60			8	16	14	1	8	87	8.05
60 to 70		•••••		4	5	1	2	12	0.97
70 to 80		1			2	1		4	0.82
Whole number of Brides	183	558	262	176	44	4	5	1,289	
Percentage of Brides	14.86	45.29	21.97	14.98	8.57	.82	.41		100.00

Of these 1,232 marriages, there were 460, or 37.34 per cent. of all, in which the bride and groom were in the same division of age. There were 418 grooms, or 33.93 per cent. of all, who were less than 25 years old; and 741 brides, or 60.15 per cent. of all, who were in the same division of ages. There were 4 grooms who were more than 70 years old, one of whom married a woman between 20 and 25 years of age.

The aggregate and average age of all persons married during the year were as follows:—

The following is a statement of the same facts for each of the twenty-five years, 1856-1880, inclusive, and for the whole period:-

	Males.	Females.
Years.	No. Aggregate age. Average age.	No. Aggregate age. Average age.
1856	639 18,104 years28.33 years	637 15,559 years24.42 years.
1857	607 16,753 years27.59 years	606 14,427 years23.84 years
1858	560 15,850 years28.29 years	560 13,702 years24.47 years
1859	611 16,989 years27.81 years	611 14,616 years23.92 years
1860	620 17.135 years27.60 years	620 14,667 years 23.66 years
1861	547 15,448 years28.24 years	547 13,316 years 22.33 years
1862	517 15,778 years28.83 years	547 13,565 years24.80 years
1863	610 17,208 years28.21 years	610 14,792 years24.24 years
1864	739 21,361 years 28.90 years	789 18,075 years 24.45 years
1865	709 20,539 years 28.96 years	709 17,605 years24.83 years
1886	812 22,801 years28.08 years	812 20,036 years24.67 years
1867	855 24,443 years28.58 years	855 21,078 years24.65 years
1868	872 24,546 years28.15 years	872 21.440 years24.59 years
1869	930 26,548 years28.55 years	930 23,135 years 24.88 years
1870	959 27,343 years28.51 years	959 23,773 years24.79 years
1871	943 26,743 years28.37 years	943 23,290 years24.70 years
1872	1,084 31,110 years28.70 years	1,084 26,835 years24.76 years.
1873	1,150 32,879 years28.50 years	1,150 28,701 years 24.96 years
1874	1,176 33,927 years28.85 years	1,176 29 510 years25.00 years
1875	1,079 30,848 years28.59 years	1,079 27,090 years25.11 years
1876	949 27,496 years28.97 years	949 23,640 years24.91 years
1877	953 27,904 years29.28 years	953 24,092 years 25.28 years
1878	1,016 29,460 years28.99 years	1,016 25,796 years25.40 years.
1879	1,071 31,016 years28.96 years	1,071 27,103 years25.32 years.
1880	1,232 36,127 years29.32 years	1,232 31,015 years25.18 years
25 years.2	11,270608,351 years28.60 years	21,267526,858 years24.77 years

The number of males and females does not agree in this table, because in 1856 the ages of 17 males and 19 females, and in 1857 the ages of 7 males and 8 females, were not given.

Number of the Marriage. Table XI. shows how many of the 1,232 marriages in Providence during the year 1880, were the first, second, third, etc., marriages of the parties:—

GROOMS.	BRIDE	Whole			
Number of the Marriage.	First.	Second.	Third.	Fourth.	number of Grooms.
Pirst	924	61	1	1	987
Second	187	86	4		237
Third	8	5	2	 	15
Fourth	3	1			8
Whole number of Brides	1 071	189	7		1 999

TABLE XI. MARRIAGES, 1880. Number of Times Married.

The 924 cases where it was the first marriage of both parties, were 75 per cent. of the whole number; in 86 cases, or 6.98 per cent. of all, it was the second marriage of both; in 137 cases it was the second marriage of the groom, and the first of the bride; this was 11.12 per cent. of all the marriages.

Denominational. The 1,232 marriages of 1880, were solemnized by the clergymen of different denominations in the city as follows:—

Roman Catholic407	Seaman's Mission 11
Calvinistic Baptist 182	Advent Christian 5
Free Baptist 60	Evangelical Advent 14
Methodist208	Latter Day Saints 10.
Episcopalian155	Hebrew4
Congregational	Bethel Pastor 2
Unitarian 89	Denomination not specified 4
Universalist 59	
Presbyterian 17	Total1,362
Christian 8	

Nativity and Residence. There were 101 marriages in 1880, or 8.20 per cent. of the whole number, in which both parties were natives of Providence; there were 128 other marriages, or 10.39 per cent. of all, in which the groom only was born in this city, and 217, or 17.61 per cent., where the bride only was born here; there were in all, therefore, 547 persons married here in 1880, who were natives of the city; this was 22.20 per cent. of all the persons married during the year.

There were 206 marriages, or 16.72 per cent. of all, where neither party was a resident of the city; of these 206 marriages there were 132 where both parties were not only not residents of the city, but came from outside of the State; and in 104 of these cases both were residents of Massachusetts; in 15 others one of the parties was from that State.

Marriage and Education. There were 2,464 persons married in Providence in 1880, of whom the following particulars are given, so to sex and nativity:—

Born in United States		Females,	890To	otal, 1,78
Born in foreign countries	. Males, 383	Females,	, 342 T	otal, %

Of these the number and proportion in each class who could be write their names, and signed the marriage certificate with a mark, were as follows:—

1880.	Whole Number Married.	Signed with a Mark.	In Each 100.
Males, born in United States	849	24	3.86
Females, born in United States	890	69	6 17
Total, born in United States	1,789	86	4.5
Males, born in foreign countries	383	64	16.71
Females, born in foreign countries	842	120	
Total, born in foreign countries	725	184	95.3
Totals	0.401	970	•

Of the 86 persons of American nativity who signed the marriage certificate with a mark, there were 22 (10 males and 12 females)

who were colored persons; and 55 (11 males and 44 females) who were children of foreign parents; there were, therefore, only 9 persons (3 males and 6 females) of white American parentage, who were unable to sign their names; this was 5 less than in 1879.

The whole number of persons who signed with a mark in 1880 was 30 more than in 1879; but the proportion of these to the whole number of persons married was 0.24 less; of those born in the United States the number was 6 more; the number of males in this class being 8 less, and of females 14 more. The proportion of Americans signing with the mark in 1880 was 0.36 less than in 1879. Of those born in foreign countries the number in 1880 who signed with a mark was 24 more than in the previous year; of males, 2 less; of females, 26 more. The proportion of all in this class in 1880 was 0.18 more than in the preceding year.

The statistics for the twenty-one years, 1859-1879, inclusive, are as follows:—

	Whole Number	Signed with	In Each
1859-1879.	Married.	a Mark.	100.
Males, born in United States	11,594	561	4.84
Pemales, born in United States	11,806	794	6.78
Total, born in United States	23,400	1,855	5.79
Males, born in foreign countries	6.638	2,159	32.52
Females, born in foreign countries	6,426	3,105	
Total, born in foreign countries	13.064	5,964	40.95
Totals	88 484	6 61Q	10 01

DEATHS.

The number of deaths reported in Providence for the year 1880 was 2,080. This was 54 more than in 1879; males, 9 less; females, 48 more; whites, 70 more; colored, 16 less; natives of the United States, 6 more: of foreign countries, 48 more; of American parage, 21 less; of foreign parentage, 75 more.

The whole number of deaths in the twenty-six years, 1855-1886, inclusive, was 35,570; the average annual death-rate for that period was one in 50.78 of the population. In 1880 the death rate was one in 50.41 of the population, or 19.83 in each 1,000. In the different classes of the population the rate was as follows:—

	Population, 1880.	Deaths, 1880.	To population one death in	Deaths to each 180 of population
Whole number	104,857			
Males				
Females				
White	101,211	1,984	51.01	
Colored	8,646	96		
Born in United States				
Born in foreign countries.	27,972	499		
American parentage				
Foreign parentage	55,575	1,160	47.91	

In the above statistics the population of each class is given from the official figures of the United States Census of 1880, except the population by parentage, which is obtained by assuming that the procentages of American and foreign, by parentage, are the same as the State Census of 1875; namely, 47 per cent. of American parentage, and 53 per cent. of foreign parentage.

The following are the general statistics relating to the 2,080 deaths of the year 1880:—

DEATHS: GENERAL FACTS.

TABLE XII. DEATHS, 1880.

Sex, condition, color, nativity, parentage, locality, season.

1880.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Whole Year.
Whole number of Deaths	201	176	183	191	189	153	208	174	143	153	146	163	2,080
Sex.—												- 1	
Males	84 117	97 79	97 86	91 100	89 100	67 86	90 118	74 100	74 69	84 69	74 72	74 89	995 1,085
Condition.—	61												
Married	55	54	57	54	59	40	44	45	45	45	41	60	599
Single	124	96	97	114	106	92	137	106	88	75	84	71	1,190
Widows	15	14	18	13	15	14	22	17	6	17	17	20	188
Widowers	6	11	11	10	9	6	5	6	4	16	3	12	99
Divorced	1	1				1					1		4
Color.—													
Whites	197	167	174	184	181	144	200	163	137	150	134	153	1,984
Colored	4	9	100	7	8	9	8	11	6	3	12	10	96
Sativity.—					11								
United States	147	133	140	143	147	114	173	132	109	112	116	115	1,581
Ireland	34	100	32	33	26	28	25	26	25	29	21	35	346
England and Scotland	12	100		12	6	7	3	10	7	8	5	10	97
Germany	2				3		2	1	2		2	2	15
British America	2			9	3	100	5	3		3	2	1	26
Portugal & West'n Islands.	1				1								2
Other countries	3	1		1	3	2		2		1			13
Parentage													
American	78	84	83	78	94	74	92	69	59	67	60	82	920
irish	81	1		84	58	55	88	71	67	61	62		827
English and Scotch	26		100	16	16		11	21	10		1000		189
German	5			3	4	1	5	3	2	1	100	2	39
British American	8	1	4	7	7	3	6	4	4	5	6	6	61
Portuguese	1	2	2	1	3	1	1	4	1				16
Other foreign	2	2	3	2	7	2	5	2		2	1		28
Locality, Wards													
1	22	21	16	18	23	21	22	20	22	23	15	30	253
Ш	1		1		100		2	6	1	7	100		65
III	17		15	18	17	12	20	14	9	10	16	17	183
IV	12	2 7	12	10	10	8	9	4	7	5	3	3	90
V	20			15	18	8	17	10	11	16	10	8	162
VI	15	5 7	8	11	7	3	5	14	7	5	8	8	98
VII	15	17	19	22	21	15	16	13	12	17	7	16	187
VIII	10	9	16	26	25	18	30	21	23	16	14	22	248
1X	19	9 25	26	23	20	21	33	19	19	23	27	24	279
X		2 37	41	39		1	100	45	25		1		435
Public Institutions	. 1	9 1	7	4	7	4	11	4	7	5	11	6	80
Totals	20	1 176	183	191	189	150	208	174	148	153	146	163	2,080

Season. The year 1880, like the two preceding, is an exception to the rule that has been found to prevail very generally in the distribution of mortality in the different seasons of the year. The largest number of deaths is generally in the third quarter of the year, and the smallest number in the second quarter. But in 1880 the largest number was in the first quarter; and the smallest in the fourth; the second quarter, also, had more deaths than the third; that the third quarter was third in rank, which has only happened twice before: namely, in 1858, and in 1879.

The number of deaths in each quarter of 1880, and of each of the five years preceding, is given below; also the quarterly aggregate and average numbers for the twenty-five years, 1855-1879, inclusive:—

						25 years, 1855 15%.
	1880.	1879.	1878.	1877.	1876.	1875. Aggregate. Aviga-
January-March	. 560	. 514	. 541	. 411	. 383	488 8,033 🗷
April-June	. 533	439	. 423	440	. 419	401 7,348 🗯
July-September	. 525	. 493	. 520	. 557	. 578	531 9,846 🖼
October-December.	462	580	. 505	530	. 490	495 8,263 和
Whole year	2,080	2,026	.1,989	1.938	.1,865	1,91533,4901,340

The number and percentage of deaths in each quarter of the year 1880, and for the period of twenty-five years, 1855-1879, were st follows:—

	1	880.	25 years, 1865-		
	Deaths.	Per cent.	Deaths.	Per cest.	
January-March	. 560	26.92	8,088	2.9	
April-June	. 528	25.63	7,348	21.84	
July-September	. 525	25.24	9,846	9.0	
October-December	. 462	22.21	8,963	si G	
Totals	.2,080	100.00	88,490	100.00	

The number and percentage are given below for each quarter of each of the twenty-six years, 1855-1879, inclusive:—

		y-March. Per cent.				-Sept. Per cent.		
555							•	
:6 6.	308	23.52	24	22.91	596	5.39	217	
	214		173	ĸ.a	9%		533	. Si.B
356	279	57.43	53		\$7		265	19 9
359	213	23.00	205	28	579		211	\$
360	343		200	39 88	257	9A	\$32	25 18
961	240	23.84	534	24	396		229	22.76
82	197	21.55	192	21.01	571	23.65	354	27.78
85	267	23.64	365	23.6	579	31.14	301	24.75
84	302	23.56	273	n.n	445	31.46	205	.93.00
65	299	91.69	310	19.81	373	\$4.72	300	24.77
966	238	22.97	218	21 04	338	21.02	313	. 23.36
367	258	26.88	302	21.04	286	22.79	214	.23.21
368	231	20.81	344	21.98	346	81.17	2:0	.96.04
989	331	96.25	250	19.90	253	98.91	312	-24.64
70	331	96.50	271	21.46	आ	29.37	290	. 22.97
71	811	94.80	263	20.98	385	30.70	295	.33.59
79	330	90.59	361	9.71	518	59.31	39 1	.34.39
3	427	24.84	407	23.68	504	29.22	381	.22.16
574	444	92.35	501	25.21	558	97.98	486	.94.46
75	488	25.48	401	20.94	531	27.73	495	.95.85
ns	383	20.54	419	23.47	573	30.79	490	.96.97
	411	21.21	440	92.70	557	28.74	530	.27.35
78	541	97.90	423	21.27	590	98.14	505	. 25.39
79	514	25.37	439	21.67	498	94.83	580	.28.63
20	560	26.92	533	25.63	525	25.24	463	.22.21

The number of deaths in the first quarter of 1880 is larger than in any previous year at that season; the percentage, though larger than in 1879, is less than in 1878, 1858 and 1856. In the second quarter of 1880 the number of deaths is larger than in any other year, and the percentage larger than in any year but 1858. In the third quarter the number and percentage are both larger than in 1879; but the percentage is smaller than any year in the record except 1879. In the fourth quarter the number is less than in any year since 1873; and the percentage less than in any year since 1874. The reason for so large a percentage in the first quarter of 1880 was the large number of deaths from scarlatina at that season. A severe epidemic of this disease prevailed throughout the year 1879, and especially in the latter part of the year, making the percentage of the fourth quarter of that year exceptionally large; the number of deaths from this

cause in 1879 was 252, of which 63.89 per cent. were in the last quarter of the year. The disease continued to prevail through the first five months of 1880, and 47.33 per cent. of the whole number of deaths caused by it, were in the first quarter of the year; after May it declined very rapidly and ceased to be epidemic. Nearly one-third of the deaths from Pneumonia were in the first quarter of the year.

Sex and Parentage. The following are the statistics of the 2,060 deaths, as to sex and parentage:—

American parentage	442 males	478 femalestotal,	23)
Foreign parentage	553 males	607 femalestotal, l	,300
		 -	_
Totals	995 males	l,085 femalestotal, 1	(0)

The number of females was 90 more than the number of males; of American parentage, 36 more; of foreign parentage, 54 more. The number of foreign parentage was 240 more than that of American parentage; of males, 111 more; of females, 129 more. During the twenty-six years, 1855–1880, inclusive, the decedents of both classes of parentage were divided as to sex, as follows:—

1855-1880......total, #579

Condition. The number of married persons who died in 1880 was 599; 4 more than in 1879; 285 of American, and 314 of foreign parentage. The number of single persons who died during the year was 1,190, or 24 more than the previous year; 472 were of American, and 718 of foreign parentage. The number of decedents who were widows was 188; 10 more than in 1879; 103 of American, and 85 of foreign parentage. The number of widowers who died was 99; 13 more than in 1879; 56 of American, and 43 of foreign parentage. There were 4 divorced persons among the decedents in 1880, which was 3 more than in 1879; they were all of American parentage.

Color. There were 1,984 white decedents in 1880; 70 more than in 1879; and 96 colored decedents, or 16 less than in the preceding year. There were 2,536 white children born in 1880; the proportion of births to deaths in this class was, therefore, 100 to 78.23.

The number of colored children born during the year was 91; the proportion of births to deaths, 100 to 105.5. In the twenty-six years, 1855-1880, inclusive, the number of colored children born was 1,938; the number of deaths among colored people in the same time was 2,049; a proportion of births to deaths of 100 to 105.7.

Nativity and Parentage. The figures of Table XII. show how important it is in giving the statistics of mortality to register the decedents by parentage, as well as by nativity. A large number of decedents who were born in the United States were children of foreign parents, and therefore under the same conditions as to locality and habits of living; they belong more properly in the foreign than in the American class. The number of decedents of American nativity was 1,581, or 76.01 per cent. of all, the number of American parentage was 920, or only 44.23 per cent. of all. The number of American nativity was 6 more than in 1879; the percentage 1.73 less: the number of American parentage was 21 less than in 1879; the percentage 2.22 less.

Of decedents of foreign birth the number in 1880 was 499, or 23.99 per cent. of all; this was 48 more than in 1879, and 1.73 per cent. more. Of those of foreign parentage the number was 1,160, or 55.77 per cent. of all; the number was 75 more and the percentage 2.22 per cent. more than in the year before.

The nativity of the 13 decedents born in "other countries" than those specified in the table was as follows: 2 each were born in Norway, Sweden and France, and 1 each in Italy, Spain, Switzerland, Holland, Austria, Poland and Liberia.

Of the 28 of "other foreign" parentage there were 7 of Italian, 6 of French, and 5 of Swedish parentage; 2 each Norwegian and Polish; 3 Swiss; and one each Dutch, Austrian and African.

Locality. Table XII. gives the number of deaths in each ward of the city and in Public Institutions for each month and for the whole year. The number of deaths in the first ward in 1880 was 26 less than in 1879; in the second ward, 3 less; in the third, 11 less; in the fourth, 26 less; in the fifth, 47 less; in the sixth, 8 less; in the seventh, 7 more; in the eighth, 50 more; in the ninth, 11 less; in the tenth, 130 more; in Public Institutions, 1 less. The seventh, eighth and tenth wards, collectively, showed an increase of 187 over the previous year; the other wards a loss of 133. It will be seen

that nearly all the increase was in the eighth and tenth wards. This was owing to the large number of deaths from scarlatina in those wards; in the tenth ward there were 94 deaths from this cause, which was 38.68 per cent. of the whole number of deaths from scarlatina; in the eighth ward 38 deaths, or 15.64 per cent. of all. This fact will be more fully commented upon in the consideration of scarlatina among the causes of death.

The number of decedents in each ward, according to parentage, was as follows:—

Wards.	American.	Foreign.	Wards.	A merican.	Porting.
I	96	157	VII	96	¶
II		12	VIШ		I
III	68	115	1X	155	
IV	44	46	x	96	
v	65	97	Public Ins	titutions 39	
VI	77	21			
Who	le citv			920	

The number of deaths in the different Public Institutions, is given below, for 1880; and for each of the two years preceding:—

	1880.		1879.	160
Rhode Island Hospital	40		84	1
Butler Hospital	8		12	1
Dexter Asylum	13	• • • • • • • • • • • • • • • • • • • •	16	f
Home for Aged Men	1		1	•••••
Home for Aged Women	7	•••••	. 5	
Children's Home	0	•••••	4	
Children's Hospital and Nursery	6		. 7	
R. C. Orphan Asylum	8	•••••	. 0	•••••
R. C. Convent	1	•••••	. 0	
.Reform School	0		. 0	*******
Central Police Station	1	•••••	. 0	•••••
Total	80	- ••••••	. 79	•••••

Aggregate and Average Age. Table XIII. gives the number of deaths and the aggregate and average age of the decedents, according to parentage, for each month, and for the whole year 1880:—

TABLE XIII. DEATHS, 1880.

Aggregate and Average Age and Parentage.

	Amer	ican Par	ratage.	Fore	ign Pare	ntage.	Total American and Foreign.			
MONTHS.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregato Age.	Average Age.	
January	. 78	2,298	39.46	193	8,091	25.13	901	5,389	26.81	
February	84	2,571	39.61	22	2,513	27.30	176	5,063	28.88	
March	83	3,964	39.23	100	2,425	24.25	183	5,686	31.04	
April	78	3,011	3 R. 6 1	113	2,995	26.50	191	6 006	31.44	
May	94	3,963	34.89	95	2,245	23.62	189	5,598	99.9	
June	74	2,618	35.3 8	79	3.298	29.00	153	1,916	32.13	
July	92	2,536	27.46	116	3,236	19.27	208	4,773	23.9	
August	69	2,914	33.00	105	2,478	23.60	174	4,693	96.97	
September	59	1,451	34.59	84	2,054	24.45	143	3,505	94.51	
Oetober	67	2,953	44.01	86	2,632	30.49	153	5,575	36.44	
November		2,309	38.48	86	1,911	22.92	146	4,220	28.90	
December	82	3,967	89 .84	81	2,863	35.33	163	6,129	87.60	
Whole Year	920	31,775	34.54	1,160	29,729	25.63	2,080	61,504	29.57	

The average age of all decedents in 1880 was 29.57 years; 0.77 years more than in 1879. Of American decedents the average age in 1880 was 34.54 years; 2.77 years more than in 1879; of foreign decedents the average age was 25.63 years, which was 0.59 years less than in the preceding year.

Comparing the year 1880 with the year before, there are the following differences in the average age of the decedents in the different months. Among the decedents of American parentage the average age in 1880 is greater than in 1879 in the months of March, April, June, August, October, November, and December; and less in 1880 in the months of January, February, May, July and September. Among the decedents of forcign parentage the average age in 1880 was greater than in 1879, in the months of March, October, November and December; and less in the other eight months of the year.

Table XIV. is a recapitulation of the same statistics for the period of twenty-five years, 1856-1880, inclusive:—

Table XIV. Deaths, Twenty-Five Years, 1856-1880.

Recapitulation. Aggregate and Average Age by Parentage.

TWENTY-FIVE	Ameri	can Par	entage.	Fore	ign Pare	ntage.		America Foreiga	
YEARS, 1856-1880.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average
January	1,525	52,017	84 .11	1,382	34,904	25.26	2,907	86.921	2
February	1.384	46,696	33.74	1,305	29,798	22.83	2,689	76,494	Ì
March	1,448	50,909	85.16	1,336	83,016	24.71	2,784	83,925	3
April	1,404	47,919	84.18	1.894	86.871	26.45	2,798	84.790	3
May	1,333	48,553	86.42	1,254	32,090	25.56	2,567	80,633	3
June	1,107	88,055	84 88	1,181	29,441	24.93	2,288	67,496	2
July	1,539	89,926	25.94	1,837	84,074	18.55	3,376	74,000	1
August	1,647	45,530	27.64	1,931	86,105	18.70	3,578	81.635	1
September	1,457	42,540	29. 2 0	1,621	81,911	19. 6 8	3,078	74,451	1
October	1,827	44,139	83.26	1,481	84,171	23 07	2,808	78,310	2
November	1,357	45,115	3 3.25	1,422	85,251	24.79	2,779	80,366	1
December	1,489	52,969	85.57	1,418	36,895	26.02	2,907	89,864	1
Total	17,017	554,368	32.58	17.562	404,517	23.03	34,579	938,885	!

These two tables show very decided differences between the average of the two classes of decedents. In the table for the year 1 (Table XIII.) the highest average age of decedents of American parentage is 44.01 years, in October; the lowest, 24.59, in Sept ber. Of those of foreign parentage, the highest average age is 3! years, in December; the lowest, 19.27 years, in July. The great difference between the two classes was in November, when the age age of American decedents was 16.26 years more than that of foreign class; the least difference was in September, when the exin favor of the American decedents was only 0.14 of one year.

In the statistics for a period of twenty-five years, as given in Ti XIV., we find differences of the same nature. The highest aver age of American decedents was 36.42 years, in May; of for decedents, 26.45 years, in April; the greatest average age of Americans is nearly 10 years above that of the foreign class.

lowest average age of American decedents was 25.94 years, in July; of foreign decedents, 18.55 years, in the same month; the lowest average age of the foreign is 7.39 years less than that of the American decedents. The average age of Americans was less in 1880 than in the long period of time, in the months of January, February, May and September; and more in the other eight months of the year. Of foreign decedents, the average age was less in 1880 than in the longer period, in the months of January, March, May and November; and greater in the remaining eight months. For the year the average age of American decedents in 1880 was 1.96 years above the average of the period of years; that of foreign decedents was 2.60 years less.

The following table shows the difference between the two classes in each month, both for the single year 1880 and for the twenty-five years, 1856-1880, inclusive. In every instance the average age of American decedents was greater than that of the foreign class:—

Difference in years between the Average Age of American and Foreign Decedents.

		1880.		25 years, 1856–1880.				
MONTHS.	Averag	e Age.		Averag	ge Age.			
	American.	Foreign.	Difference.	American.	Foreign.	Difference		
Janu ary	29.46	25.13	4.83	84.11	25.26	8.85		
February	30.61	27.30	8.81	83.74	22.83	10.91		
March	39.32	24.25	15.07	85.16	24.71	10.45		
April	88.61	26.50	12.11	84.13	26.45	7.68		
Мау	84.89	23 .63	11.26	86.42	25.58	10.84		
June	\$5.88	29.09	6.29	84.88	24.93	9.45		
Joly	27 48	19.27	8.19	25.94	18.55	7.39		
August	32.09	28.60	8.49	27.64	18.70	8.94		
September	24.59	24.45	0.14	29.20	19.68	9.52		
October	44.01	30.49	13.52	88.26	23.07	10.19		
November	88.48	22.22	16.98	83.25	24.79	8.46		
December	89.84	35.33	4.51	35.57	26.02	9.55		
Whole year	34.54	25.63	8.91	82.58	23.03	9.55		

It has generally appeared from this table that the difference between the two classes is greater in the long period of years than in a single year. But in this instance there are six months where there is more difference between them in 1880 than in the twenty-five years, namely: March, April, May, July, October and November; and six months where it is the reverse of this.

The largest number of deaths in one month in 1880 (Table XIII.) among those of American parentage was 94 in May; the least number, 59 in September. Of the foreign decedents the largest monthly total was 123 in January; the smallest, 79, in June. In the twenty-five years, 1856–1880, (Table XIV.), the largest monthly aggregate was in August for both classes; of American decedents, 1,647; of foreign, 1,931; the smallest monthly aggregate for both classes was in June, American, 1,107; foreign, 1,181. The percentages of all the deaths in the different quarters of the year, according to parentage, for the period of twenty-five years, 1856–1880, were as follows:—

	1st Quarter.	2d Quarter.	3d Quarter. 4th Quarter.
American parentage	25.60	22.59	37.29
Foreign parentage	22.91	21.80	90.6991.690

Table XV. gives the number of decedents and the aggregate and average age, according to parentage, in each of the twenty-five years, 1856-1880, inclusive, and for the whole period:—

DEATHS: AGGREGATE AND AVERAGE AGE.

TABLE XV. DEATHS, 1856-1880.

Recapitulation. Aggregate and Average Age by Parentage.

	Amer	ican Par	entage.	Fore	ign Pare	ntage.	Total	Americ Foreign	
YEARS.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregate Years.	Average Years.
356	511	14,554	26.90	524	8,258	15.75	1,065	22,812	21.4
\$57	523	16,314	81.19	402	8,007	19.91	925	24,321	26.9
36,	51%	15,910	81.07	505	8,478	16.78	1,017	24.388	23.9
350	479	14.993	81.30	420	8,640	20.57	899	23.633	26.9
860	512	16,660	32.51	480	9,751	19.94	1,001	26,411	26.8
961	538	16,327	20.34	513	9,605	18.72	1,051	25,932	24.0
863	460	14,589	81.71	454	10,994	24.21	914	25,583	28.0
963	597	19,396	32.49	617	13,757	22.29	1,214	33 .155	27.
864	619	21,782	83.48	632	11,865	17.98	1,281	83,097	25.
865	631	19.628	81.10	580	580 11,991 20.67 523 12,762 24.40		1,211	81,619	26.
860	518	18,906	3 8.85	523	12,762	24.40	1,036	31,668	80.
967	528	19.258	3 6.47	432	- I I II		960	81,407	82.
866	596	21,779	36.54	514	12.539	24.89	1.110	84,318	80.
949	627	20,490	32.6 8	629	14,776	23.49	1,256	85,266	28.
870	642	21,078	32.83	621	14,393	23.18	1,203	85,471	28.
871	643	22,725	35.34	611	17.005	27.83	1,254	89. 730	81
873	832	27,572	83.14	771	19,160	24.86	1,608	46,782	29.
573	862	28,675	83.27	857	19,061	22.24	1,719	47,736	27.
87L	885	27,588	81.17	1,102			1,987	52,390	26.
875	914	80,419	83.28	1.001	24.802	24.28	1,913	54,721	28.
876	875	29,025	83.17	990	23,959	24.20	1,865	52,984	29.
877	908	28,853	81.22	1,030	25,410	24.67		53,763	27.
578	889	26,724	80.06	1,100	25,178	22 89	1,989	51,992	26.
1879	941	29,896	81.77	1,085	28,446	26.22	2,026	58,342	2 8.
890	990	81,775	84.51	1,160	29,729	25. 63	2,080	61,504	29.
b years' totals	17,017	551,368	82.58	17,562	404,517	23.03	34,579	958,885	27.

The greatest difference between the average age of the two of decedents was 15.50 years in 1864; the least difference years in 1879. The difference between them in the whole per 9.55 years.

Number and percentage of decedents at different ages. Tabigives the number and percentage, according to parentage, dents at different periods of life in 1880; and the same fact decedents in the twenty-five years, 1856-1880, inclusive:—

Table XVI. Deaths, 1880.

Percentages, at Different Ages, according to Parentages.

	Amer Paren 188	tage,	Fore Paren 188	tage,	Total. can an eign,	d For-
AGES.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Under l year	175	19.02	206	17.76	381	18.32
1 and under 2	62	6.74	85	7.33	147	7.07
2 and under 5	74	8.04	129	11.12	203	9.76
Total under 5	311	33.80	420	36.21	731	35.15
5 and under 10	48	5.22	88	7.59	136	6.54
10 and under 15	14	1.52	27	2.33	41	1.97
15 and under 20	9	.98	43	8.71	59	2,50
20 and under 30	65	7.07	142	12.24	207	9.95
30 and under 40	82	8.91	87	7.50	169	8.12
40 and under 50	57	6.20	99	8.53	156	7.50
50 and under 60	60	6.52	88	7.50	148	7.12
60 and under 70	84	9.13	80	6.89	164	7.88
70 and under 80	119	12.93	55	4.74	174	8.37
80 and under 90	60	6.52	21	1.81	81	3.89
90 and over	11	1,20	10	.86	21	1.01
Total	920	100.00	1,160	100.00	2.080	100.00

From Table XIII. we found that the average age of all A decedents in 1880 was a little more than in 1879; from Tab

years of age in 1880 was slightly more than in the preceding year, yet the percentages of deaths among the aged were also very much larger than in 1879; thus of decedents between 70 and 80 years old, the number in 1879 was 83, and the percentage 8.82; in 1880 the number was 119, and the percentage 12.93; also of persons over 90, the number and percentage in 1880 were more than double those of 1879; this, of course, raised the average age above that of the previous year. Of decedents of foreign parentage, the number under 5 years of age in 1880 was larger than in 1879, but the percentage 1.03 less; but among the aged decedents of this class the number and percentage were considerably less in 1880 than in 1879; of those living beyond the age of 60 years the percentage in 1880 was 1.55 less than in the preceding year; so that the average age of all was reduced, as was shown by Table XIII.

The following statement gives the percentage, according to parentage, of the decedents under 5 years old, in each of the twenty-five years, 1856-1880, inclusive:—

	American.		otal American & Foreign.
1000	Under five years.		Under five years.
1806	37.15 per cent	54.95 per cent .	45.91 per cent.
1857	32.31 per cent	51.74 per cent	40.75 per cent.
3 86 8	36.35 per cent	55.41 per cent .	45 33 per cent.
1859	. 28 60 per cent	48.34 per cent	37.82 per cent.
3000	32.81 per cent	48 87 ner cent.	
1861	35 87 ner cent	59 68 per cent	44.05 per cent.
1663	21.00 per centiti	49.07 man cant	36.54 per cent.
1963		42.07 per cent	per cont.
1964	28.11 per cent	46.81 per cent	
	28.81 per cent	51.11 per cent.	39.82 per cent.
1000	29.95 per cent	46.37 per cent	37.82 per cent.
1506	23.78 per cent	40.53 per cent.	32.24 per cent.
1867	24 05 per cent	36.35 per cent	29.59 per cent.
1366	27.35 per cent	43.19 per cent.	34.69 per cent.
1369	30.78 per cent	43.24 per cent.	87.02 per cent
1670	31.00 per cent	42.67 per cent	
1871	28.69 per cent	86 01 per cent	32.22 per cent.
1879	30.90 per cent	40.00 per cent	
1673.		40.00 per cent	рег оещь
187L	au.as per cent	43.05 per cent.	36.71 per cent.
1675.	34.02 per cent	43.55 per cent.	
1878	82.06 per cent	40.96 per cent.	
1877	84.29 per cent	48.13 per cent.	38 98 per cent.
	32.93 per cent	41.26 per cent.	
	28 78 nor cont	49 AL ner cent	40 03 ner cent
**********	\$2.94 per cont	87.94 per cent.	
	33 .80 per cent	36.21 per cent.	85.15 per cent

During the forty-one years, 1840-1860, inclusive, there were 47,724 decedents in Providence, of whom 18.742 were less than five years old; the number and percentage in each division of age under 5 years were as follows:—

Whole number of decedents						_
Under one year				per e	cent. o	€ all.
One, and under two years	4.820	or	10 10	per e	cei.t o	f all.
Two, and under five years	4.704	or	9 86	per e	cent o	f all.
Total, under five years	15.743	- Ot	30.27	per c	cent. o	f all.

Table XVII. shows the number and percentage of decedents, in the different divisions of age, according to parentage, in the aggregate for the twenty-five years, 1856-1880, inclusive:—

Table XVII. Deaths, 1856-1880.

Number and Percentage at Different Ages, according to Parentage.

	A merican 25 years,	Parentage 1856–1860.	Foreign 25 years,	Parentage, 1856-1880.	American Parentage 1856	& Foreign c. 25 years, -lee0.
AGES.	Number of Deaths.	Percentage in each division of age.	Number of Deaths.	Percentage in each division of age.	Number of Deaths.	Percentage in each division of age.
Under 1 year	2,953	17.85	3,756	21.39	6,709	19.40
1 and under 2	1,118	6.57	1,919	10.93	8,037	8.78
2 and under 5	1,815	7.78	1,946	11.08	8,261	9.43
Total under 5	5,886	31 65	7,621	43.40	13,007	87.61
5 and under 10	887	5.21	925	5.27	1,812	5.24
10 and under 15	877	2.21	886	1.91	713	2.06
15 and under 20	584	8.48	555	8.16	1,139	8.29
20 and under 30	1,565	9.20	1,914	10.90	8,479	10.06
30 and under 40	1,474	8.66	1,645	9.87	8,119	9.03
40 and under 50	1,202	7.59	1.884	7.59	2,626	7.60
50 and under 60	1,871	8.06	1,110	6.32	2.481	7.18
80 and under 70	1,468	8.63	996	5.61	2,454	7.10
70 and under 80	1,560	9.17	757	4.81	2,817	6.70
80 and under 90	893	5.25	298	1.67	1,186	8.43
90 and over	160	.94	86	.49	246	.n
Totals	17,017	100.00	17,563	100.00	84,579	100.00

These Tables, XVI. and XVII., show that the percentage of children under five years of age is much greater among the decedents of foreign than of American parentage. In the year 1880 (see Table XVI.) the percentage of this class among foreign decedents was 2.41 more than among American decedents; in the long period of time the difference was much greater; the percentage of foreign was 11.75 more than that of American decedents. This corresponds with the tact proved from Tables XIII. and XIV., that the average age of foreign decedents is much less than that of American decedents. In the 25 years, 1856-1880, the average age of decedents of American parentage was 32.58 years, while that of the other class was only 23.03 years. The number of American decedents during this period of years, who lived beyond fifty years of age, was 5,452, which was 32.05 per cent. of all American decedents in that time; the number of foreign decedents in the same period, who lived to be more than fifty years old, was 3,232, or 18.40 per cent. of all the foreign decedents.

In comparing Tables XVI. and XVII. we see that the percentage of children of American parentage under five years old was more in 1880 than in the long period of time; but of decedents over sixty years of age the percentage was 5.79 more in the single year than in the term of years, which caused the average age of American decedents in 1880 to be greater than during the long number of years; as was shown to be the case in Tables XIII. and XIV. From those tables also we saw that the average age of decedents of foreign parentage was higher in 1880 than for the long period of years; this agrees with the facts found from Tables XVI. and XVII., that the percentage of children in this class under five years old in 1880 was much less than the average for the twenty-five years, and the percentage of decedents over sixty years old much larger.

Deaths by age and parentage in each month and quarter of the year. Table XVIII., on page 47, shows the comparative mortality of the two classes of decedents in different seasons, and at the various ages, by giving the number of deaths at each age, and of all ages, according to parentage, in each month and quarter of the year. The whole number of deaths in the year was 2,080; of American parentage, 920; of foreign parentage, 1,160; divided as follows in the different seasons:—

	lst Quarter.	2d Quarter.	3d Quarter. 4th Quarter-
American Parentage .	26.63	26.74	23.9112.77
Foreign Parentage	27.16	21 74	26.29 21.8

Of the decedents of American parentage the largest percentage was in the second quarter of the year,—a very unusual occurrence; the first quarter is next in order; and the third quarter is third in order. Of the deaths in this class 53.37 per cent. were in the first half of the year. Of decedents of foreign parentage the largest percentage was in the first quarter of the year; the next in the third quarter; 51.90 per cent. of the deaths were in the first half of the year.

Of children under one year old, the deaths in 1880 were as follows: American parentage 175, or 35 more than in 1879; 77 of these deaths, or 44 per cent., were in the third quarter of the year; the whole number of deaths, of all ages, in the third quarter was 220, of which 77, or 35 per cent., were children less than a year old.

Of foreign parentage the decedents of this class numbered 206, which was 30 more than in 1879; 85 of these children, or 41.26 per cent., died in the third quarter of the year; these 85 deaths were 27.87 per cent. of the whole number of deaths in that quarter.

Causes of death; sex, age, parentage and percentage from boom causes. Table XIX. gives these particulars for each of the 2,080 deaths during the year.

Season and Disease. Table XX. gives the number of deaths from each cause in each month and quarter of the year, showing the comparative prevalence and mortality of different diseases in the different seasons.

Table XVIII. Deaths by Age and Parentage in each Month and Quarter, 1880.

=							<u>, -</u>									
					Div	ISIO	NAL	PE	RIOD	8 01	, l.i	FB.				
	DIVISIONS OF THE YEAR.	Under 1.	1 to 2.	2 to 5.	5 to 10.	10 to 15.	15 to 20.	20 to 80.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.	80 to 90.	90 to 100.	ALL AGES.
	January February March	11 12 7	4 2 6	15 8 10	7 6 5	i	1 1 2	7 9 4	6 10 7	5 4 8	4 7 4	3 7 7	12 7 14	3 3 8	i	78 84 88
	lst quarter	30	18	83	18	1	4	20	23	17	15	17	33	14	2	245
AGE.	April	11 18 16		5 6 6	7 5 4	1 3 1	1 i	5 9 5	6 8 5	5 6 6	4 6 8	10 10 8	13 11 8	5 5 4	1 2 4	78 94 74
ARENT	2d quarter	45	13	17	16	5	2	19	19	16	13	28	32	14	7	246
Q.	August	85 18 24	8	5 4 5	3 8 	 2 1	 i	4 2 3	4 5 4	2 4 2	7 2 5	7 10 5	12 6 8	6		92 69 59
MERICAN	3d quarter	77	18	14	6	3	1	9	13	8	14	23	21	14	••••	220
₹	October November December	6 7 10	5 2 6	2 3 5	4 1 3	2 8	; j	1 6 10	8 12 7	5 5 6	7 4 7	6 6 5	12 10 11	6 1 11	9	67 60 82
	4th quarter	23	13	10	8	5	2	17	27	16	18	17	33	13	2	209
	Whole year	175	62	74	48	14	9	65	82	57	60	84	119	60	11	990
	JanuaryFebruary	14 10 17	11 7 6	16 9 15	11	2	8 5 8	9 12 13	9 4 10	9 13 6	9 7 14	11 5 6	4 3 4	 3	1 1 1	128 92 100
	ist quarter	41	24	40	3 6	6	11	81	23	28	30	22	11	6	3	815
ARENTAGR.	April	17 17 15	_5	16 12 9	11	ĺi	6 4 1	9 12	7	7 8 6	8 5 8	10 5 4	7 4 8	1 3 1	1 1 2	118 95 79
HEN	2d quarter	49	15	87	24	7	11	39	16	21	21	19	19	5	4	287
FOREIGN P.	July	44 28 13	15 12 6	5 9 7	5 4 2	8 2 4	2 3 6	11 10 14	8 12	6 5 8	1 10 7	7 8 2	4 2 2	3 4 1	2	116 105 84
FOR	3d quarter	85	33	21	11	9	11	85	28	19	18	17	8	8	2	305
	October November December	9 14 8	3	16 7	9 7 1	1 8 1	8 5 2	10 11 13	9 5 6	10 8 13	9 5 5	7 6 9	4 3 10	i	1 	86 86 81
	4th quarter Whole year	31 206	13 85	31 129	17 88	5 27	10 48	34 142	20 87	89 31	19 68	22 80	17 55	2 21	1 10	258 1,160
T	otal Amer. & For	381	147	203	136	41	53	207	169	15%	148	164	174	81	21	2,080

Ratio of desiries from each cause to whole nom-ber from known therefore 8 Disease, sex, age, parentage, and percentage from known causes. Jant Foreign. DUR OF as values letter the ության Dan: 07 19bau DIVISIONAL PERRORS OF LIFE. tinn 18 Dan 02 | gg ameun]; [i I-IIH OF ரு கூடிய bur oc արվեւ 30 Dan 02 ban c Og reban nuder 15 Ol Tobut pun g S wad under 5. year. I and Under 2 Lasha's .fatoT SEX. Решавса. **DEATHS**, 1880. Mules. Accidents (various).... Aneurlam Bones, lifsenses of...... Biain, Congestion of...... Asphyxia..... Drowning..... Anæmia...... Apoplexy...... Atelectasis Pulmonum...... Railroad Polsoning..... Falls..... A becesses...... Bladder, Diseases of...... CAUSES OF DEATH. Burns and Scalds TABLE XIX. Asthma..... Diseases of

TABLE XIX.—CONTINUED. D	EATE	DEATHS , 1880.	80	Dis	808	, 86	8,	de.	pa	e a c	ge.	ž	a	rce	nta	20	rom	knor	on ca	Disease, sex, age, parentage, and percentage from known causes.
7						Ā	DIVISIONAL PERIODS OF LIFE.	OMAL	M	1012	0	3	4				PAN	PARENTAGE.	i i	nown Cause Ca Ca Ca Ca Ca Ca Ca Ca Ca Ca Ca Ca Ca
CAUSES OF DEATH.	sə[s]	emsjes.	otal.	nder I year. Ina	2 Tebau bas	bas	Of 19bau bna (Al 19bau	Sand noder 99.	6 and 100 and 100.	ьпя 0 Оф төрии	бая 0 .00 төраи	рия 0 ппдет 60.	6 жл б 107 төрпи	рая 0 .08 тэбаи	bas 0 under 90.	band over.	asohom.	.agle10i	latol.	tatto of d rom each o whole sent from k
Brain Indammation of	PE 33	_ 1	13		<u> </u>	<u> </u>	ж	8	8	1.	17	\neg :	1-	,	1:	6 :	7 3	4 ×	12	8 8
Bronohitis	2	64	\$	5	•	4		:	-	:		65	4	=	8	_	82	ž	3	2.71
a	2	18	4	:	<u>:</u>	- :			4	64	•	0	a	9	-04	:	8	9	4	8.8
of Breast	:		ä	<u>:</u>	<u>:</u>	÷	÷		:	:	:	_	÷	i	÷	:	Н	_	64	.10
" of Stomach	-	-	66	$\frac{\cdot}{\cdot}$	<u>:</u>	÷	$\stackrel{:}{\leftarrow}$:	64	01	_	-	-01-	÷	:	4	4	6 0	8.
" of Uterus	:	9	8	÷	:	<u>:</u>	<u>.</u>		7	00	60		64	÷	÷	:	40	60	16	2.
Catarrb		:	_	=	:	\div	<u>.</u>		•	:	i	÷	÷	÷	÷	:	_	i	_	8
Childbirth	:	9	2	÷	<u>:</u>	÷	:	:	-8	**	=	i	÷	÷	÷	:	4	•	2	84.
" Puerperal Convul	:	*	**	$\frac{\cdot}{\cdot}$:	\div	<u>:</u>		•	-	i	i	Ť	i	÷	:	•	=	•	.15
" Puerperal Fever		=	Ħ	$\frac{\cdot}{\cdot}$:	\div	\div	:	-	4	Ħ	i	÷	÷	i	:	4	-	=	3.
" Abortion		64	ei	$\frac{\cdot}{\cdot}$	<u>:</u>	÷	÷		-	_		i	÷	÷	÷	:	~	-	C4	.10
Cholera Infantum	\$	23	100	12	84	<u>.</u>	÷	:		i	:	÷	:	÷	÷	:	7	28	8	4.84
" Morbus	_	**	Ť	$\frac{\cdot}{\cdot}$	÷	÷	_;.		_	_	Ì	÷	:	_	-	:	*	-04	*	.19
Colto	64	:	A	<u>:</u>	<u>:</u>	:	<u>:</u>				:	=		÷	÷	:	i	- 64	64	.10
Consumption	18	188	88	4	4	4	_	81	117	8	\$	9	8	10	=	:	Ħ	211	328	15.60
Convulsions	2	a	9	ձ	60	-	÷	:	•	_	Ť	Ì	:	:	÷	;	8	8	9	2.57
Cronn	-	31	8	65	4	7				_	-	- <u>:</u>	-	_	_	-	F	81	23	1.41

TABLE XIX.—CONTINUED. D	DEATHS,	8, 18	1880.	3	eas	Lisease, sex, age, parenlage, and	, ag	e.	are	gal	, a		§	arta L	8	percentage from	Knoton	200 2	causes.	
		SEX.				Ę	DIVISIONAL PERIODS OF	[TE	PERIO	8	Į į	LIFE.				PAR	PARENTAGE.		enna -mun	UMOU
CAUSES OF DEATH.	.105.	.solam	.faic	der l year. bud	ander 2.	nnder 5. nnder 10.	ander 15.	ora nador 20. baa	nder 80.	ander 40.	nnder 50. and onder 60.	.07 19brin	рия .08 төрии	ander 90.	Rnd OVer,	.naoi1em	.nglero	.iate of d	M hole	er itom ki
	אי	24		7	3 8	3 0	οι	۱ ۱	08	07	<u> </u>	<u>09</u>	01	1		Y),d		oi	90 90
Debility	1	छ-	22	2	=	<u>:</u> -	Ė	<u>:</u> -	÷	÷	·	≟	_	-	:	2	60	8		Ŕ
" Premature Birth	•	6 0	14	=	÷	<u>:</u>	i	-	<u>:</u>	÷	<u>:</u>	<u>:</u>	:	:	i	10	a	7		ŝ
Diabetes	-	10	- 6	÷	-:	_ <u>:</u> -	Ė	÷	:	<u> </u>	··	_		_	ī	10	A	6		×
Diarrhoa	51	9	88	8	65	<u>.</u>	i	÷	:	÷	<u>.</u> ;	_	7	81	:	a	9	8		¥.
" Chronic	_	F	Ø4	÷	$\frac{\cdot}{\cdot}$	<u>:</u>	i	÷	\div	<u>:</u>	∔		7		<u> </u>	#	i	N		ř
Diphtheria	22	8	5	**	8	<u> </u>	64		- 84	-	-			:	:	8	8	9	•	×
Dropsy	ю		•	:	÷	-	i	:	-;	<u> </u>			_	_	-	60	**	6		ş.
Dysentery	4	- 66	13	•	•		_	-	-:		-	_	34	Ī	- :	6	•	育		ž
Embolism		•	Ä	:	<u>:</u>	<u>:</u>	i	:	<u>:</u>	÷	<u>.</u>			:	:		:	F		Ŗ
Enteritis	0	2	19	4	64	<u>.</u>	:	÷	-	-01	65	*		:	-	-	월	9		8
Epilopsy	_	- 65	*	÷	\div	<u>:</u>	i	:	84	÷	<u>:</u>		_		:	60		-		Ŧ.
Erysipelas	10	4	۵	÷	\div	<u>:</u>	i	:	-	:	<i>(</i> 4	_	_	=	:	•	83	_a_		ź
Exposure and Neglect	-	85	1-	60		<u>:</u>	1	:	:	-	-	<u>.</u>		:	<u>:</u>	60	4	7		ž
Fever, Malarial		:	Ħ	÷	÷	<u>:</u>	Ì	:	:	:	<u>:</u>	<u>:</u>	:	:	÷	:				ş
" Typhoid	7	88	잂	÷	÷		•	a	8	1-	01	-		:	:	3	88	25		8
Gall Bladder, Disease of	_	A	84	<u>:</u>	÷	:	Ė	÷	÷	:	÷	•		•	•	- i-	Ī	on		Ĕ.
Gangrene, Senile		77	Tá	⋮	-			-	-	_	_:	_	_	-	7	a.	-	•		÷

		SEX.				^	IVIE	IOMA	DIVISIONAL PERIODS OF LIFE.	N N	8	Į,	,				PA	PARENTAGE.	Η.	edine essa: mun nwon
CAUSES OF DEATH.	Males.	Pennales.	Total.	Jabri year.	ander 2. Sand Bas 6	ander 5.	of rabous bas 0	ander 15.	og 19pau Dang (i Og 19pau	50. 200 and 60. 40. 40. 40.	bna 0. Jog 19bnu	рал 0 .00 тэраи	osnd Or 19bau	.08 төрпи .08 төрпи	o and under 90.	Dand Over.	merican.	.nglato	'otal.	tatio of de rom each o rom bole eer from kr
Hematocele	1:	1	17	T	<u> </u>	1:	나.	۲Ļ:	<u>; :</u>	1	<u>. [</u>	<u>. [</u>	1:	1:	· :	B :	7		17	- I -
Heart, Diseases of	- F	\$	8	i	-:	:	-:	_		=	2	22	18	8	5	~	23	17	88	4.65
" Hypertrophy of	•	*	-60	-	÷	<u>:</u>	$\frac{\cdot}{\cdot}$	-:		_				64	69	:	4	4	80	×.
Heat		i	-	÷	:	:	\div	÷		≟	_	:		:	:	i		H	-	8.
Hemorrhage	4	60	7	8		÷	$\frac{\cdot}{\cdot}$	_ <u>:</u> -		<u>:</u>		:	_:	:	:	:	4	95	-	क
" from Lungs	••	•	Ħ	÷	•	÷	÷	<u>:</u>	<u>:</u>			-6	64	:	:	:	•	15	=	8
Hernia	•	₩	Ю	7	=	÷	÷	<u>:</u>	:			:	64	i	i	:	64	65	=0	ž
Hip Joint, Disease of	:	-	ਜਂ	$\frac{\cdot}{\cdot}$:	:		:	<u>:</u>	<u>:</u> -	<u>:</u>	:	<u>:</u>	-	i	:	H		-	8.
Hooping-Cough		10	=	æ	:	*	÷	<u>:</u>	:		<u>.</u> .		:	:	:	•	4	-	11	8
Hydrocephalus	7	2	8	=	=	-6	04	<u>:</u>			<u>:</u>	:	_:	:	:	:	8	=	8	1.41
Hydrothorax	*		64	÷	<u>:</u>	<u>:</u>	÷	$\frac{\cdot}{\cdot}$	<u>:</u>			:	:	_	:	:	61	•	01	.10
Insanity.	04	•	80	÷	$\frac{\cdot}{\cdot}$	\div	$\frac{\cdot}{\cdot}$:	:		_	04	60	F	:	·	10	8	80	8.
Intemperance	•	16	œ ·	÷	÷	<u>:</u>	$\frac{\cdot}{\cdot}$	<u>:</u>	<u>:</u>	- -		61	:	i	İ	:	64	-	0	₹
Intrasusception		1	F	÷	÷		-	÷	<u>:</u>	∔	:	:	<u>:</u>	-	i	i	-	7	-	8.
Kidneys, Diseases of	_	80	8	÷	÷	:	=	<u></u>	\vdots	-		89	80	01	_	Ī	•	8	12	85.
"Addison's Disease of		_	+	÷	$\dot{}$	÷	÷	÷	:	÷	<u>:</u>			:	:	÷	:		-	8
" Bright's Disease of	<u>=</u>	9	8	-	-	=	-	<u> </u>	-	-æ: -	_	a	6 5		7	=	18	7	8	1.59

TABLE XIX.—CONTINUED. D	DEATHS , 1880.	8, 18	80.	Ä	eas	3	a, a	ge, 1	par	nta	g,	and	8	ا رو د	tage	Disease, sex, age, parenlage, and percenlage from known causes.	knon	m ca	28	
		SEX.				5	DIVISIONAL PERIODS OF LIFE.	MAL	P	Į O	0	LI	, i			PA PA	PARENTAGE		SAURS	DOMD
CAUSES OF DEATH.	Males.	Решалев.	Total.	Under I year. baa I	bunder 8.	d 19bnu bra d 01 19bnu	ol rehan has ol di rehan	5 49 pur 30 02 49 50 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19 02 19	So and of so.	50-a-08 bna-08 	.05 Tehnn has 05	.08 18 brar Dan 08	Date 70.	OS YSDAU Dan OS	90 and 80.	очет, Атаетісап.	Foreign.	Total.	Ratio of districtions of distr	Der from k
Knee Joint, Disease of	ai		100	:		 	<u> </u>		İΞ	İĖ	 	╁	 :	H	 		-	-	İ	15
Laryngitis		6	63	÷	-	<u>:</u>	:	:	Ī	=	:	÷	÷	÷	<u>:</u>		**	67		55.
Leucocythæmia	÷	:		$\frac{\cdot}{\cdot}$:	<u>:</u>	<u>:</u>	:	Ì	•	=	÷	÷	<u>:</u>	<u>:</u>			7		8
Liver, Diseases of	2	2	8	÷	:	<u>:</u>	<u>:</u>	:	:	_	6	*	10	<u>;</u>	÷	2	2	8		8
" Inflammation of	- 61	i	ak	÷	-	÷	<u>.</u>	:	÷	:	$\frac{\cdot}{\cdot}$	-:	_	-	÷		~	61		2.
Locomotor Ataxia	**		63	÷	÷	÷		:	Ť	:	*	:	÷	÷	÷		:	65		25
Lungs, Diseases of	-	•	-	\div	- :	÷	:	:	÷	:	:	:	÷	÷	<u>:</u>	:	=	=		8
Malformations	•	0	=	=	:	\div	<u>:</u>		i	÷	:	\div	÷	÷	<u>:</u>	•	90	=		23
Marasmus	2	9	23	22	84	÷	<u>:</u>	:	÷	;	7	-	_	-	÷	. 18	17	3		8.
Measles	:	45	65	÷	.	=	<u>:</u>		÷	÷	:	÷	÷	÷	÷	i	65	65		.13
Meningitis, Cerebro-Spinal	3 1	84	4	æ	=	=	<u>.</u>	:	÷	:	<u>:</u>	÷	÷	÷	<u>.</u> -		60	4		2
Metritis	:	_	Ä	÷	÷	÷	<u>:</u>	:	-	÷	<u> </u>	÷	:	÷	<u>:</u>		:	_		8
Murder	:	_	Ä	÷	÷	\div	<u>:</u>	:	:	_	$\frac{\cdot}{\cdot}$	÷	÷	÷	<u>:</u>	<u>:</u>				S
Old Age	8	4	Ę	\div	<u>:</u>	÷	<u>:</u>	:	÷	÷	÷	-	_	<u></u>	=	28	5	Į.		83.58
Paralysis	1	*	8	$\frac{\cdot}{\cdot}$	÷	÷	÷	$\overline{\vdots}$	÷	;	80	60	<u>.</u>	<u>±</u>	<i>(</i> 4)	18	۵	8		1.46
Peritonitis	6	•	7	÷	÷	÷	-	_	64	**	_	- 76	-	-	<u>.</u> .		65	7		S.
Plearing	- ; 	_ :	- 10		-	-	≟]	-	=	=	-	- <u>:</u>	-		 -	-	7		2.

fo whole ber from ki seenso	25.	1.11	8	.15	s	11.71	8.	ş	8	.15	8	97.	ૹ઼	.10	٩.	7 8.	91.
Ratio of do																	
Total.	170	83	-	60	11	243	1	н	1	60	-	04	90	61	Q3	9	0
Foreign.	16	10	:	64	1-	155	-	-	7	1	7	04	49	61	1	03+	1
American.	4	13	F	1	4	88	H	1	1	01	1	:	60		Н	94	-
90 and Over.	-	1	:	1	:	1	:	:	÷	1	Ť	÷	i	1	:	:	
80 and under 90.	00	-	î	i	H	i	i	:	;	1	:	:	÷	i	i		
70 and under 80.	18	2	H	i	H	i	i	1	÷	ŕ	1	i	01	i	i	i	
60 and under 70.	83	60	:	:	-	1	İ	i	i	:	i	-	-	:	•	i	P
50 and under 60.	98	1	i	1	÷	i	i	i	i	İ	Ť	i	4	F	•	i	
40 and under 50.	8	9)	Ť	i	G)	:	Ť	÷	:	i	i	F	-	:	F	ri	B
30 and under 40	=	00	i	i	4	:	i	;	i		ë	1	:	-	-		1
20 and under 30.	120	:	İ	-	:	-	i	i	i	İ	:	i	Ť	:	:	H	
15 and under 20.	10	i	i	;	F	60	÷	i	÷	ri	i	:	İ	i	i	;	
ander 15.	24	i	i	i	:	15	Ť	÷	÷	:	÷	÷	i	Ť	Ť	Ť	-
under 10.	100	i	i	÷	i	13	÷	÷	1	÷	÷	1	i	:	Ť	Ť	1
ander 5.	100	÷	i	i	-	98	-	i	Ť	-	÷	i	i	:	i	-	
l and under 2. bas 2	=	-	1	1	:	53	:	i	:	1	÷	i	i	÷	:	1	
year.	18	£-	i	01	1	13	i	:	-	i	•	i	i	:	i	01	•
Total.	170	22	-	80	n.	243	1.	1	1	60	-	03	80	61	01	10	•
Females.	68	30	1	1	*	121	-	1	П	Т	-	69	9	91	1	7	-
Males.	81	13	-	OV	1-	199		1	:	61	:	1	05	-	1	1	-
1	1:	:	1		-			÷	÷	1	÷	÷	-	÷	-	:	
CAUSES OF DEATH.	Pneumonía	" Congestion of Lungs	Prostate, Disease of	Purpura, Bemorrhagica	Rheumatism	Soarlating	Scrofula	Septicemia	Skin, Diseases of	Spine, Diseases of	Spleen, Disease of	Stomach, Diseases of	" Inflammation of	Buiolde	Surgical Operation	Syphilis	Tahes Mesenteries

Table XIX.—Continued. Deaths, 1880. Disease, sex, age, parentage, and percentage from known causes.

		SEX.					DIV	OISI	NAL	DIVISIONAL PERIODS OF LIFE.	TOD	8 01	LI	FE.			T	PAI	PARENTAGE,	GE,	mnt
CAUSES OF DEATH.	Males,	Females.	Total.	Under I year.	f and t	g and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	70 and under 80.	80 and under 90.	90 and Over,	Апретевп	Foreign.	Total.	пот евсь с
reething	-	6	16	00	00	1	1				1	1	:		1	1	1	4	15	16	.78
Throat, Diseases of	1	1	64	H	-	:	1		-		:		:	1	:	:	:	21	-	-01	.10
[uperculosis	89	9	6	-	1	1	1	-	:	00	65	-		:	1		:	7	01	6	.44
ramors	1	10	9	;	1			:	1	:	1	1	61		01	-		9	-	9	.29
Ulcers	-	64	c,	:	1	-	1		1		1	1	:	-	-	:	1	-		01	-10
Uramia	1	1	-	İ	1	H	1	1	:	•	1	-		:	:	1	:	1	1	-	.05
Unknown	1-	10	17	13		-		1	1	-	0,	1	-	1	-	:	1	6	00	11	
Totals	892	1,085	2,080	381	1	203	136	4	52	202	169	156	148	164	174	8	22	950	1,160	2,080	100.00
								1													

CAUSES OF DEATH.	.Tibuast.	Гергаагу.	Матер.	First Quarter.	April.	May.	Jane.	Second Quarter.	JuJA.	Jangua	September.	Third Quarter.	October.	November.	December.	Fourth Quarter.	Whole Year.
Aocidents (all kinds)	8	8		181		100	1	2	00	1	8	22	 -	1	10	188	8
Apoplexy	4	a	61	60	•	-1	04	12	4			9	60	4	-	19	3
Brain, Diseases of	11	4	•	-61	4	21	4	8	9	•	•	81	60	4	-	19	85
Bronchitis	99	8	20	18	OS.	*	4	19	_	*	4	•	*		O.	18	28
Cancer (all kinds)	-	64	*	13	40	7	a	8	gh	18	64	71	8	- 25	•	11	8
Childbirth	4	•	æ	11	1	64	60	•	=	:	61	**	:	64	*	9	8
Cholera Infantum	:	i	:	:	:	Ø1	40	-	3	- 83	=	88	65	H	H	10	100
Cholera Morbus		i	:		:		i		•	-	:	÷	:	i	:	:	
Consumption	88	54	8	88	22	19	18	\$	88	8	8	8	2	8	8	88	8
Convulsions	*	∞	8	6	-	24	0 0	11	∞	+	66	9	8	4	•	13	
Croup	10	:	- 8 5	80	_	i	:	÷	<u>:</u>	:	60	60	64	=	7	17	28
Diarrhœa		:	69	69	-	i	_	61	7	ю	40	*	oi	Ī	:	69	8
Diphtheria	۲	•	4	11	6	*	80	138	•	ĸ	10	91	10	10	0	15	5
Dysentery	i	<u> </u>	•	:	61	_	61	¥Ģ.	OH	_	-	4	*	:	:	*	51
Enteritis	4	63	:	+	:	04	_	65	64	- 6	:	-	:	01	i	61	19
Erysipelas		34	H	4	N	i	_	95	÷	:	:	-	i	:	_	-	
Fever, Typhoid	61		*	9	4	G	•	17	_	-61	00	=	=	8	69	8	23
Heart, Diseases of	Ħ	=	11	2	•	2	91	8	ю	15	8	16	-	77	20	8	30
												_					

TABLE XX.—CONTINUED. DEATHS, 1880. Season and Disease.

Whole Year.	66	80	6	94	55	85	80	*	75	30	11	193	243	01	16	11	188	St. Dest
Fourth Quarter.	9	1	01	19	19	60	1	-	19	9	90	43	30	-	28	et	387	Ton a
ресешьет.	1	1	1	10	80	1	1	-	6	62	60	16	0	1	-		14	1000
November.	80	-	-	9		-	-	1		-		H	Je.	-	-	-	3.6	New.
October.	G.	-	1	00	20	64	-	-	00	04	1	16	90		-	*		1000
Third Quarter.	10	10	-	00	10	17	CH	01	13	10	G#	14	55	-	10	•	47	-
September	2	01	ON	10	*	4	-	1	-	64	i	01	-	i	61	21	a a	-
.tsuZuA	01	-	1	C4	-	10	H	1	9	62	F	00	13	ì	+		17	1
Jaja.	00	01	-	,	B	100	7	ci.	-	10	1	*	13	-	-	61	11	1
Second Quarter.	9	0.0		90	1-	9	-	1	24	00	10	7	76	1	100	•	107	Ì
.enut.	C.		-	-	-	**	1	1	10	64	69	119	15	i	64	1	3.0	1
May	20	05	Ì	*	01	01	H	1	00	69	1	ă	68	T	1	60	18	1
April.	-	1	F	00	9	1	1	İ	0	4	63	15	60	1	-	-	118	-
First Quarter.	-	1	64	п	*	6	İ	i	19	9	4	62	115	1	1	9	800	1
March.	04	1	1	01	00	60	Ì	1	10	54	O1	26	8	1	1	-	3.6	1
February.	20	1	-	*	-	10	1	1	5	-	G4	13	98	-		*	310	1
January.	Ca	1		10	-	H	-	1	*	*	1	17	90	-	-	1	30	-
CAUSES OF DEATH.	Hydrocephalus	Insanity		Kidneys, Diseases of	Liver, Diseases of	Marasmus	Measles	Meningitis, Cerebro-Spinal	Old Age and Senile Gangrene	Paralysis	Peritonitis	Pneumonia and Congestion of Lungs	Scarlatina	Saleide	Teething	Unknown	Other Causes, as given	

NOTES ON THE CAUSES OF DEATH,

DURING THE YEAR 1880.

In the following notes information is given about the various causes of death, which could not be presented in the tables on the preceding pages, and comparisons are made with the statistics of previous years.

The percentage of deaths from the different causes is always reckoned on the number of deaths from known causes.

The number of deaths in Providence in 1880 was 2,080; in 17 cases, or 0.82 per cent. of all, the cause of death was not stated; this was 8 more deaths, and 0.38 per cent. more than in 1879.

The number of deaths from known causes, therefore, in 1880, was 2,063; and on this number all the percentages are reckoned. During the twenty-five years, 1856–1880, the total number of deaths was 34,579; of which 550, or 1.59 per cent., were from unknown causes; 34,029 from known causes. In the forty-one years, 1840–1880, inclusive, the number of deaths recorded in Providence was 47,724; 46,172 from known causes; and 1,552, or 3.25 per cent., the causes of which were not specified.

Table XX. shows to what degree different diseases prevail in different seasons of the year. It is expected to find an excess of diarrhoeal diseases in the third quarter of the year, and the figures show that of the 165 deaths during the year from this class of diseases 127, or 77 per cent., occurred during the three months of July, August, and September. On the other hand diseases of the respiratory system prevail less at this season and much more extensively in the colder months; thus of 249 deaths from pneumonia and bronchitis, only 22, or about .09 per cent., were in the third quarter.

The number of deaths from diphtheria was much less than for some years past; but it prevailed to some extent throughout the year, the number of deaths being very evenly distributed in the different seasons. Scarlatina prevailed most severely in the early part of the year; 47.33 per cent. of the deaths from this cause being in the first quarter.

Abscesses. The number of deaths caused by abscesses in 1880 was three, which was 0.15 per cent. of all deaths from known cause; the number was two less, and the per cent. 0.10 less than in 1873. One was a "lumbar abscess," and one of the parotid gland; one was not specified. Other abscesses are reckoned under the head of disease of the organ affected. Of these three decedents, 1 was a male, and 2 were females; 1 of American, and 2 of foreign parentage. The statistics for the 25 years, 1856–1880, inclusive, are as follows:—

Number of deaths from all known causes, 34,029.
From abscesses: males, 49; females, 45; total, 94; or, 0.28 per cent of all American parentage, 42; foreign parentage, 52.

Accidents. The number of deaths resulting from accidents, in 1880, was larger than in any previous year, though there have been a few years when the percentage was greater. The number was 65, which was 9 more than in 1879, and the percentage was 0.37 more; the percentage was larger than in any year since 1873. In Table XIX., 2 of these deaths are recorded as caused by asphyxia; 11, by burns or scalds; 15, by drowning; 9, by falls; 1, by poisoning; 4, by railroad accidents, and 18 are included under the heading 6 "various." Of the latter, one was "run over by a low gear;" one was "hit in the face by a sky-rocket;" one had the "simple fractured by falling brick;" two were "thrown from a wagon;" three had a "leg broken;" two had an "arm broken;" one bad "ribs broken;" one "spine dislocated;" and three died from "injury at birth."

The numbers of each sex and parentage were as follows:-

Males, 47; females, 18; total 65, or 3.15 per cent. of all. American parentage, 19; foreign parentage, 46.

The numbers of each sex and parentage for the twenty-five years, 1856-1880, are:—

Males, 739; females, 247; total, 986, or 2.90 per cent. of all.

American parentage, 382; foreign parentage, 604.

Aneurism. There were two deaths from this cause in 1880, or 0.10 per cent. of all from known causes. The decedents were 1 male, and 1 female; 1 of American, and 1 of foreign parentage. The statistics of this cause of death for the twenty-five years, 1856–1880, are:—

Males, 22; females, 4; total, 26, or 0.08 per cent. of all. American parentage, 9; foreign parentage, 17.

Apoplexy. The number of deaths caused by apoplexy in 1880, was 48, which was 2.32 per cent. of all. The number was 8 less, and the per cent. 0.46 less than in the previous year.

Males, 21; females, 27; total, 48, or 2.32 per cent. of all.

American parentage, 32; foreign parentage, 16.

The statistics for twenty-five years, 1856-1880, are:—

Males, 354; females, 370; total, 724, or 2.13 per cent. of all.

American parentage, 506; foreign parentage, 218.

Further particulars on this subject are given under the head of "Paralysis."

Brain, Diseases of. The number of deaths, reported under this head in 1880, was 80; of these, 9 were specified as "congestion," and 49 as "inflammation." The number of deaths was 4 more than in 1879, and the per cent. 0.10 more. The statistics of sex and parentage are:—

Males, 42; females, 38; total, 80, or 3.87 per cent. of all. American parentage, 38; foreign parentage, 42.

For the twenty-five years, 1856-1880, the statistics are as follows:—

Males, 715; females, 605; total, 1,320, or 3 88 per cent. of all. American parentage, 692; foreign parentage, 628.

It will be seen that there is very little difference between the number of decedents of American and of foreign parentage, both in the single year 1880, and in the period of twenty-five years, for which the statistics are given. But from apoplexy, a disease which also affects chiefly the brain, the number of American decedents is considerably more than twice that of the foreign class.

Bronchitis. This was reported as the cause of 56 deaths in 1880, or 2.71 per cent. of all. It is a disease which affects chiefly the very young and the very old; 21 of the decedents, or 37.5 per cent. of all were less than one year old; and 15, or 26.8 per cent., were more than 70 years old. The number of deaths was 22 more than in 1879, and the per cent. 1.03 more. The statistics as to sex and parentage are as follows:—

Males, 29; females, 27; total, 56, or 2.71 per cent. of all. American parentage, 22; foreign parentage, 34.

For the twenty-five years, 1856-1880, the statistics are:-

Males, 201; females, 258; total, 459, or 1 35 per cent. of all. American parentage, 204; foreign parentage, 255.

Cancers. The number of deaths from cancers was larger than ever before in a single year, and the per cent. more than in any year except 1877. The number in 1880, was 68, or 3.30 per cent. of all from known causes; it was 2 more, and 0.03 per cent. more than in 1879. In Table XIX. 2 are specified as cancer of the breast; 8, se cancer of the stomach; and 16, as cancer of the uterus; the 42 which were included under the term "various," were located as follows: liver, 14; rectum, 3; ovaries, 2; mesentery, 2; kidneys, 2; interpretable, perinæum, omentum, bowels and bladder, one each; tongot, 2; neck, pharynx, lip and throat, lung and sub-maxillary gland. one each; one epithelial, one abdominal, and one bone cancer; of four the location was not specified.

The following statistics of sex and parentage are given:

Males, 25; females, 43; total, 68, or 3.30 per cent. of all. American parentage, 39; foreign parentage, 29.

For the twenty-five years, 1856-1880, the statistics are:

Males, 225; females, 604; total, 829, or 2.44 per cent. of all. American parentage, 525; foreign parentage, 304.

Of the 829 decedents, 72.9 per cent. were females; and 63.3 per cent. were of American parentage.

Child-birth. There were 26 deaths from child-birth in 1880, including 3 from puerperal convulsions; 11, from puerperal fever; and 2 from abortion; the 26 deaths were 1.26 per cent. of all deaths from known causes, and just 1.00 per cent. of the 2,600 mothers bearing children in 1880. The number of deaths was 6 more than in 1879, and the per cent., 0.26 more. The decedents were 11 of American, and 15 of foreign parentage. In the twenty-five years, 1856–1880, the deaths from this cause have been 405; 172 of American, and 233 of foreign parentage. This was 1.19 per cent. of all deaths during that period, and 0.84 per cent. of the 48,442 mothers bearing children in the same time.

Cholera Infantum. There were 100 deaths in 1880 from this cause, a much larger number than for the two years previous. The percentage of all from known causes was 4.84, which was 1.92 more than in 1879. The statistics are as follows:—

Males, 42; females, 58; total 100, or 4.84 per cent. of all. American parentage, 41; foreign parentage, 59.

The following are the statistics of sex and parentage for each of the twenty-five years, 1856-1880, and for the whole period:—

	SK	x.	PARENT	AGE.	Whole
	Males.	Females.	American.	Foreign.	Number. Percentage
856	85	22	16	41	575.5
857	27	23	26	24	505.5
838	27	22	18	86	49
859	14	19	13	90	333.8
860	35	33	26	42	687.0
861	29	82	25	86	616.0
863	25	19	17	27	444.9
863	36	80	19	47	665.5
864	29	29	18	40	584.6
865	20	80	13	87	504.9
866	80	17	15	32	474.6
867	24	25	23	26	495.1
868	38	82	22	48	70
b 69	85	80	 2 7	38	655.9
870	48	45	87	56	93
671	30	82	26	86	625.0
1873	72	79	58	98	151 9.5
873	42	50	39	58	925.4
1874	79	62	44	90	1846.8
875	62	74	50	86	136
1876	66	56	48	79	1226.8
			43		1226.5
1878	35	85	28	44	70
1879	80	29	28	81	592.9
1000			41	***	1004.8

Cholera Morbus. The number of deaths reported from this cause each year is very small; in 1880, it was 4; 1 male, and 3 females; 2 of American, and 2 of foreign parentage; total 4, or 0.19 per cent. of all from known causes. In the twenty-five years, 1856-1880, the number of deaths from this cause was 116, or 0.34 per cent. of all.

Consumption. There were 322 deaths from consumption during the year 1880, which was 15.60 per cent. of the whole number of deaths from known causes; in 1879, the number of deaths from this cause was 293, and the per cent. 14.52. The particulars of sex and parentage in 1880, were as follows:—

American parentagemales, Foreign parentagemales,	•
Totalmales,	
The statistics for the twenty-five	e years, 1856-1880, are:-

Whole number of deaths from known causes, 84.029; deaths from consumption

Males, 45.54 per cent......females, 54.46 per cent. American parentage, 47.01 per cent....foreign parentage, 53.99 per cent.

The population of Providence by the United States census of June, 1880, was 104,857; the death-rate from consumption for the year was, therefore, one in 326 of the inhabitants, or 3.07 in each thousand persons. The average annual rate for the fifteen years, 1840–1854, was one in 209.9; and for the twenty-six years, 1855–1880, it was one in 306.1.

By periods of five years from 1840 to 1879 inclusive, and for the whole period of forty years, the whole number of deaths from known causes in Providence, and the number and percentage from consumption, were as follows:—

	Number of deaths	Number from	Percentage of deaths
	from known causes.	Consumption.	from Consumption
1840-1844	2,824	745	26.38
845-1849		761	19 55
850-1854	4,465	958	21.46
855-1859	4,753	930	19.57
1860-1864	5,298	997	18.82
1865-1869	5,474		18,43
1870-1874	7,784	1,174	
1875-1879	9,688		15.26
		<u> </u>	
1840-1879, 40 vear	rs44,109		18.25

The number and percentage of deaths in each of the last ten years have been:—

	Number.	Per cent.
1871	194	15.66
1872		15.26
1873		13.57
1874		18.76
1875		15.79
1876		
1877		15.22
1878		15.46
1879	293	14.59
1880		

The number and percentage of the decedents from consumption in each division of age, were as follows for the year 1880, and for the period of 25 years, 1856 to 1880 inclusive:—

Periods.	Under 1	Carrie	under 2.	2 and	under 5.	6 and under 10.	10 and under 15.	15 and		20 and under 30.	30 and	under 40.	40 and under 50.	50 and	60 and	under 70.	nnder 80.	80 and under 90.	Total.
1890. Number		4	4	-	4	1		1 -	12	117	1	65	-	1		29	5	-	822
Per cent	1.9	4	1.24	1.5	24	.31	2.49	8.8	33	36.84	20	. 19	14.25	4.9	79.0	01(1	1.55	.31	100.00
1836-1880. Number		18	67		61	58	117	51	12	1,769	1,5	900	79	51	7 8	75	139	13	5,690
Per cent	1.1	19	1.18	1.0	07	1.09	2.00	9.0	00	81.09	21	.09	13.9	9.0	9 6.	59	2.44	.28	100.60

The largest proportion of the decedents are between twenty and fifty years old; 70.82 per cent. of all, in 1880, being within those ages; 50.32 per cent. lived beyond the age of thirty. The average age of all decedents in 1880, was 29.57 years. Of the 5,690 decedents from consumption in the twenty-five years, 1856–1880, 66.13 per cent. were between twenty and fifty years of age; 53.39 per cent. lived beyond the age of thirty. The average age of all decedents during this period was 27.73 years.

Convulsions. There were 49 deaths from convulsions in 1880, which was 10 less than in the previous year; 34 of the decedents were less than a year old.

Males, 24; females, 25; total, 49, or 2.37 per cent. of all. American parentage, 23; foreign parentage, 26.

For the twenty-five years, 1856-1880, the statistics are:-

Males, 495; females, 402; total, 897, or 2.64 per cent. of all American parentage, 371; foreign parentage, 526.

Croup. There were 29 deaths from croup in 1880, which was it less than in 1879, and less than in any year since 1874, when the number was the same as in 1880. Of the 29 deaths, 17, or 58.82 per cent., were in the fourth quarter of the year; 3 of the decedent were under one year old; 4 were between 1 and 2; 14, between 2 and 5 years; 7, between 5 and 10; and one between 10 and 15. The statistics of sex and parentage were as follows:

Males, 8; females, 21; total, 29, or 1.41 per cent. of all. American parentage, 7; foreign parentage, 22.

For the twenty-five years, 1856–1880, the numbers are:

Whole number of deaths, 768; annual average, 31.

Males, 384; females, 384: total, 768, or 2 26 per cent. of all.

American parentage, 314; foreign parentage, 454.

The following table shows the number of deaths from croup for each of the twenty-six years, 1855–1880 inclusive; also the percentage each year, of the whole number from known causes; and the particulars as to sex, age, parentage, season, locality and color:—

797		327	5888427145511	F125828855
29	ac 22	1-81	© :00 : : : : ∞01 E.4.	94 -4-04 -HICKS
2.13	161	212	O100048191 - 141-91	00 -01 -00 -00 -00 -00
38	828	18	****************	D-10-100-4-1
2.07	823	16	E-854- 01600	F-614 : 81 : 81 6 62 4
3.48	25 25	258	WRONE IN CEUE	80-30-480-880-880
2.95	55	35	Oc. 8 14 1 10 8 54	0 :0 :4 : :446
65.	100	12	м-шей : ::- менена	4
22	15.6	156	******	******
121	22	, e	**************************************	4
33.82	178	128	0000H 0-H4F	∞-∞-∞≈≈+r-
2.16	118	22	*****	онао : овио о
1.53	00	92	- m- m-m-	0
1.01	66	+1+		01 14 00 00 00 11 00 10 10 10 10 10 10 10 10
19	60	100	00 00 00 00 100 100 mmmon	*******
2.27	22	F 21	10 H H H H H H 10 H 10 H 10 H 10 H 10 H	8 :0 :0 H
2.44	18	200		@01.00 may = 111 m
3.59	88	13	**************************************	ಅಚಕಾರ್ವವಿ
2.78	988	28	01:00:00:00:01 : :01:4:00:10	æ .→∞≈∞∞ · · ·
3.51	22	52	10 00 01 01 10 11 11 11 11 11 11 11 11 11 11 11 11 1	8 48888
20.00	22	66		********
25.80	P =	44	01-01-4-1-1-1-1-101	H010001H04
24	20	2122	F : 00 00 - 00 00 00	64 : 55 55 55 12-
617	212	212	*********	9999F99
82.13	93	10	2044224422000	00 +00 00 to to
96	90	900	4001-001-00 :0004	01-00-00-00
3.12	27	138	010000000000000000000000000000000000000	0-5:000 0-000
Whole number		American	February January March April April Andry June July Suptember October December	Ward II.

Debility. The number of deaths reported from this cause in 1880 was 27, including 14 from premature birth; in 1879, the number was 33, including 10 from premature birth. The particulars for the year 1880 are as follows:—

Males 13; females, 14; total, 27, or 1.31 per cent. of all. American parentage, 10; foreign parentage, 17.

For the twenty-five years, 1856-1880:-

Whole number of deaths, 600; annual average, 24.

Males, 314; females, 286; total, 600, or 1.76 per cent. of all.

American parentage, 248; foreign parentage, 352.

Diarrhæa and Dysentery. There were 42 deaths from these causes in 1880, or 2 more than in 1879, and 2.03 per cent. of all deaths from known causes. The statistics for each disease are as follows:-

	Males.	Females.	American.	Foreign.	Total. Percentage
Acute Diarrhœa	12	16	9	19	:28
Chronic Diarrhœa	1	1	2	-	2
Dysentery	4	8	6	6	12 #
Total					
The deaths from given in connection					
	Males.	Females.	American.	Foreign.	Total. Percentage.
Cholera Infantum	42	58	41	59	100
Cholera Morbus	1	8	2	2	4
Enteritis					

The total number of deaths from these five diseases, for the jest 1880, was as follows:

Males, 69; females, 96; total, 165, or 7.98 per cent. of all.

American parentage, 67; foreign parentage, 98.

The number of deaths is 45 more than in 1879, and the per cent., 1.99 more; the number of males, 8 more; of females, 37 more; of American parentage, 15 more; of foreign parentage, 30 more.

The statistics of the five diseases, for the twenty-five years, 1856–1880, are:—

Number of deaths, 3,780; annual average, 151.

Males, 1880; females, 1,900; total, 3,780, or 11.11 per cent. of all.

American parentage, 1,535; foreign parentage, 2,245.

Of decedents from these diseases in 1880, the proportions between the two classes of parentage were as follows: 146.3 decedents of foreign parentage to each 100 of American parentage, or 59.39 of foreign, and 40.61 of American parentage in each 100 decedents. Of decedents from all causes in 1880, the proportions were 126.1 of foreign to each 100 of American parentage, or 55.77 foreign, and 44.23 American in each 100.

Though the excess of foreign decedents from diarrheal diseases is much greater than when all causes are compared, yet the number of deaths is not so disproportionate, when we consider the number of children living of each class, and that most of the decedents from these causes are very young children. Thus in 1880, 78.79 per cent. of the decedents from diarrheal diseases were children less than two years old; of the 2,627 children born during the same year, 52.33 per cent. were of foreign, and 47.67 per cent. of American parentage; 109.8 children of foreign to each 100 of American parentage. In 1879, the proportions of children born were 51.31 per cent. of foreign, and 48.69 of American parentage, or 105 foreign to 100 American.

The following table shows the number of deaths from each of these causes, with the number from the whole class of causes, and their percentage of all from known causes, reported in Providence during the period from 1840 to 1854 inclusive; during each year, from 1855 to 1880 inclusive, and in the aggregate during the whole period of 41 years, 1840 to 1880 inclusive:—

		311					Per cen
	Cholera		laba D	To		All	of all fro
			arrhœa. Dy	•			
			849		102		
l	61	4	87				
· · · · ·	57	5	29	85	6	132	11
)	50	2	3 8	29	5	194	13
)	49	2	27	13	8	98	9
3	83	1	28	13	4	79	8
3	68	2	21	18	9	118	11
l .	61	5	22	32	6	126	11
ł	44	1	81	22	9	107	11
3	66	5	85	77	9	192	13
3	58	4	45	40	18	160	н
)	50	7	34	62	4	157	19
7	47	17	28	44	9	145	1
9	49	2	15	43	10		
)	70	8	24	15	8	120	10
5	65	6	24	23	6		
	93		22		15		1
	62		24		18		16
		10	52		14		10
	92		87		25		
	134		50	•	25		1
	186		36		81		1
	122		40		18		
	122		84		21		1
	70						
	•		28		18		
		5	31		16		••••••
v	100	. 4	30	13	19	165	••••••

Deaths from Asiatic cholera are not included in these statistics; they have been as follows: in 1849, 158; in 1854, 159; in 1866. 14; total, 331.

Diphtheria. The epidemic of this disease, which had prevailed to such an alarming extent in Providence during the four years, 1876-1879 inclusive, diminished very decidedly in 1880, though some deaths occurred in each month, yet the whole number during the year was only 61, which was 46 less than in the previous year, and the percentage was 2.35 less. The deaths were distributed through the year with remarkable equality, as follows:

1st quarter17	3d quarter16
2d quarter13	4th quarter 15

As usual, the decedents were mostly young children; 55, or 90.16 per cent., were less than ten years old.

The statistics of sex and parentage were as follows:—

Males, 32; females, 29; total, 61, or 2.95 per cent. of all.

American parentage, 35; foreign parentage, 26.

During the four years' epidemic of diphtheria, the decedents of foreign parentage were greatly in excess of those of American parentage, the proportion being 56.39 of foreign, and 43.61 of American parentage in each 100; previous to the epidemic the reverse of this was true, the figures being 58.68 of American, and 41.32 of foreign parentage in each 100 decedents. In 1880, the American class is again in excess; 57.38 per cent. of American, and 42.62 of foreign parentage in each 100 decedents.

The first cases of diphtheria in Providence were reported in 1858; for the twenty-three years, 1858–1880, the statistics are as follows:—

Deaths from known causes in 23 years, 32,101.

From diphtheria, 1,154; annual average, 50.

Males, 575; females, 579; total, 1,154, or 3.59 per cent. of all.

American parentage, 562; foreign parentage, 592.

In the following table are given the statistics as to sex, parentage, season and locality, for each of the twenty-three years, 1858–1880, and for the whole period:—

DEATHS FROM DIPHTHERIA IN PROVIDENCE, TWENTY-THREE YEARS, 1858-1880.

DIPHTHERIA.	1858.	1859.	1860	1861.	1863.	1863.	1864.	1865.1	866.1	867.	808	1869.	1870.	1871.	1872.	1873.	874.	1875.	.978	1877.	1878.	1879.	.0880	1838, 1838, 1803, 1803, 1803, 1803, 1804, 1805, 1805, 1805, 1805, 1805, 1871, 1872, 1873, 1874, 1875, 1877, 1878, 1879, 1880, Total.
Whole number	0.50	1.96	2.60	2,17	1.92	3.54	46	1.85	1.08	0.53	97.0	1.05	0.80	1.69	12.1	1.35	1.02	0,745	111	296	246	107	28.9	1,154
SEX: Males	-4	so co.	71	101	48	18	88	21.4	100	02.01	:10	00 KG	00	51.0	118	14	100	2000	25 25	146	136	46	50 61	575
PARENTAGE: American	44	11	16	9 33	11 6	38	183	10	910		.0	91-	40	70	81-	52.8	91	10	5.80	129	198	75	10.00	503
SEASONS: Fabruary March April May June June Joly Soutember I Soctober Docomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber I Socomber	0444	9 1 1 400			0 : 1 : 1 : 1 : 1 : 0 0 0	F-100-00 0400	001-00-00000000000000000000000000000000	- 01	m .mm .m .0100	01-1 1 1 1 1-1	00				Q41201-02 - 001 4	**************************************	010H : 010H : F00H		C .40000000000	42882222538528	122215.55558	Örene a örer a siste	F-04040500000	81-88-45-48-48-48-48-48-48-48-48-48-48-48-48-48-
Vard II 8 9 1 1 1 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1				2000 - 2004 · ·	нначакн	200-20-50 ::	r-r-Bauara	PENGONA HERRORA E 100000H	227 4 4		n ::::::::::::::::::::::::::::::::::::	muuun i4	85 IIII	4 4		n-0	- I I I I I I	H0101 :01- :00	B-1000+0000	8.2822819	84455525	Hadian-ded	600 HOLES	52 25 25 25 25 25 25 25 25 25 25 25 25 2

The connection of season with the mortality from diphtheria is a subject which has received considerable attention in these reports, since the disease has prevailed so extensively in our city; and facts of importance and interest are shown by the study, particularly when compared with similar statistics in regard to scarlatina and croup.

The following table shows the number and the percentage of deaths in Providence, from diphtheria, scarlatina, and croup, in each month and quarter of the year. The figures for scarlatina and croup include the aggregate mortality from these diseases during the twenty-six years, from 1855 to 1880 inclusive. The figures for diphtheria are for twenty-three years, from 1858 to 1880 inclusive.

		HERIA. -1880.		ATINA. 5-1880.		OUP. -18 80.
Months.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
January	108	9.36	. 257	12.57	106	13.82
February	27	6.67	220	10.76	89	11.18
March	90	7.80	188	9.20	89	11.18
lst quarter	275	23.83	665	32.53	284	85.68
April	78	6.76	183	8.95	65	8.16
Мау	72	6.24	196	9.10	42	5.28
June	70	6.06	160	7.83	27	8.89
2d quarter	220	19.06	529	25.88	134	16.83
Jaly	47	4.07	109	5.83	27	8.89
August	49	4.95	87	4.26	11	1.88
September	98	8.49	95	4.65	43	5.41
3d quarter	194	16.81	291	14.24	81	10.18
October	175	15.16	136	6.65	75	9.42
November	163	14.18	180	8.81	110	13.82
December	127	11.01	213	11.89	119	14.07
4th quarter	465	40.30	559	27.35	297	87.81
Totals	1,154	100.00	2,044	100 00	796	100.00

Dividing the year into periods of four months, the numbers and proportions are as follows:—

	DIPHT	HERIA.	SCARL	ATINA.	Сво	CP.
Seasons.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.
1st 4 months	853	80.59	848	41.48	349	43.84
2d 4 months	238	20.63	542	26.53	107	13.44
8d 4 months	563	48.79	654	82.60	840	41.77

Dividing the year into two periods, the numbers and proportions are:—

	DIPRT	HERIA.	SCARL	AKITA.	Cas	OUP.
Seasons.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.
1st 6 months		42.89 57.11	1,194 850	58.41 41.59	418 378	3.0 47.0

These figures show that, while the percentage of deaths from each disease is smallest in the second four months of the year, there are decided differences as to the prevalence of the three in the first third and last third of the year.

In Diphtheria, the percentage of the last four months is much larger than that of the first four; while in Croup, which is frequently compared with Diphtheria, notwithstanding the marked difference between the two, the percentages of the first and the last third of the year are nearly equal; that of the first third being slightly greater. When the year is divided into two equal perioder the difference in the prevalence of these two diseases is still more marked; the percentage of Diphtheria in the second six months being much greater than in the first half of the year; and the percentages of Croup being quite the reverse.

With regard to Scarlatina, the percentage of the first third of the year is considerably larger than that of the last third; and the per-

centage of the first six months much greater than that of the second

In the whole period the smallest percentage of mortality in any month was as follows for each disease: diphtheria, 4.07 in July; scarlatina, 4.26, and croup, 1.38, both in August. The largest monthly percentage of each was: diphtheria, 15.16 in October; scarlatina, 12.57 in January; and croup, 14.07 in December.

Dropsy. This was assigned as the cause of 6 deaths in 1880; 5 males, and 1 female; 3 of American, and 3 of foreign parentage; total, 6, or 0.29 per cent. of all.

Erysipelas. The number of deaths reported from this cause in 1880 was 9, which was one less than in 1879.

Males, 5; females, 4; total, 9; or 0.44 per cent. of all. American parentage, 6; foreign parentage, 3.

For the twenty-five years, 1856 to 1880, the statistics are:—

Males, 87; females, 92; total, 179; or 0.53 per cent. of all.

American parentage, 105: foreign parentage, 74.

Exposure and Neglect. The number of deaths ascribed to exposure or neglect in 1880, was 7, which was as many as in the two years previous combined. Three of the decedents were less than a year old; two were between 1 and 2 years of age; one, between 40 and 50; and one between 70 and 80.

Males, 4; females, 8; total, 7; or 0.34 per cent. of all. American parentage, 3; foreign parentage, 4.

Fever, Typhoid. There were 52 deaths from this cause in 1880, which was 12 more than in 1879, and 0.54 per cent. more. There was also one death reported from malarial fever, which is included in the following figures:—

Males, 25; females, 28; total, 53; or 2.57 per cent. of all. American parentage, 24; foreign parentage, 29.

In the following statistics for the twenty-five years, 1856-1880, besides the deaths from typhoid, are a few that have been reported from "fever," and "bilious fever," and about 30 from typhus fever:—

Whole number, 1,075; annual average, 48.

Males, 552; females, 523; total, 1,075, or 3.16 per cent. of all.

American parentage, 594; foreign parentage, 481.

Heart, Diseases of. The number of deaths from diseases of the heart in 1880, was 104, including 8 from "hypertrophy." Among other affections of the heart, the following were specified on the returns: "pericarditis," "endocarditis," "angina pectoris," "atrophy," "paralysis," "dilatation," and "fatty degeneration" of the heart. The number of deaths was 7 less than in 1879, and the per cent. 0.47 less.

Males, 55; females, 49; total, 104, or 5.04 per cent. of all.

American parentage, 59; foreign parentage, 45.

In the twenty-five years, 1856-1880, the number of deaths from heart disease was 1,582, or 4.65 per cent. of all; annual average, 63.

The particulars for each year are given in the following table:—

	81	x.	PAREN	TAGE.	Whole	
	Males.	Females.	American.	Foreign.	Number.	Percent
856	8	18	16	5		
857	18	18	22	9	31	
858	18	23	98	8	36	
859	13	11	17	7	34	
960	15	16	21	10	81	
361	28	18	29	19	41	
969	96	23	83	16	49	
63	23	19	80	19	49	
364	26	82	44	14	58	
85	18	29	83	14	47	
366	19	21	27	18		
87	26	23	85	14		
368	25	27	84	18	59	
369	83	29	85	27	68	
370						
371		• • • • • • • • • • • • • • • • • • •				
379					•••••	
78					100	
374	200000				104	
75					80	•••••
376					99.	•••••
377					02	
378					es	•••••
379						
340	,					
30V	50	19	99	±0	104	
years' total.						_

Hemorrhage. There were 18 deaths reported from this cause in 1880, or 0.87 per cent. of all. This was 8 more than in 1879, and 0.37 per cent. more.

Males, 9; females, 9; total, 18, or 0.87 per cent. of all. American parentage, 10; foreign parentage, 8.

In the table, 11 are specified as hemorrhage from the lungs; the others were, 2 from the bowels, 1 uterine, and 4 umbilical hemorrhage.

Hernia was reported as the cause of 5 deaths in 1880, the same number as in each of the two years previous.

Males, 3; females, 2; total, 5, or 0.24 per cent. of all.

American parentage, 2; foreign parentage, 3.

Hooping Cough. There were 11 deaths from this cause during the year; 4 less than in 1879. Eight of the decedents were less than a year old; and 3, between 2 and 5 years.

Males, 6; females, 5; total, 11, or 0.53 per cent. of all.

American parentage, 4; foreign parentage, 7.

For the twenty-five years, 1856-1880, the statistics are:—

Whole number, 397; annual average, 16.

Males, 170; females, 227; total, 897, or 1.17 per cent. of all.

American parentage, 165; foreign parentage, 232.

Hydrocephalus. This was stated as the cause of 29 deaths in 1880, or 6 less than in 1879; all the decedents were less than ten years old, and 22 of them were less than 2 years old. The particulars are as follows:—

Males, 14; females, 15; total, 29, or 1.41 per cent. of all.

American parentage, 12; foreign parentage, 17.

For the twenty-five years, 1856-1880:-

Whole number, 762; annual average, 80.

Males, 414: females, 348; total, 762, or 2.24 per cent. of all.

American parentage, 382; foreign parentage, 380.

Insanity. There were eight deaths from insanity during the year 1880, which was 4 less than in 1879.

Males, 2; females, 6; total, 8, or 0.39 per cent. of all. American parentage, 5; foreign parentage, 3.

The statistics for the twenty-five years, 1856-1880, are:-

Whole number, 256; annual average, 10.

Males, 122; females, 184; total, 256, or 0.75 per cent. of all.

American parentage, 182; foreign parentage, 74.

Of the 256 decedents, 71.09 per cent. were of American parentage Most of these deaths occur at Butler Hospital for the insane, which is within the city limits, but where the patients are from many different places. Of the 8 deaths in 1880, only 1 occurred outside of the Hospital.

Intemperance was the cause of 9 deaths in 1880, or 2 more than it the previous year.

Males, 4; females, 5; total, 9, or 0.44 per cent. of all. American parentage, 2; foreign parentage, 7.

For the twenty-four years, 1857-1880, the statistics are:—

Whole number from known causes, 33,001.

From intemperance, 238; annual average, 10.

Males, 172; females, 66; total, 238, or 0.72 per cent. of all.

American parentage, 68; foreign parentage, 170.

Of the 238 decedents, 72.27 per cent. were males, and 71.43 p cent. of foreign parentage.

Kidneys, Diseases of. There were 46 deaths from diseases of t kidneys, in 1880, including 33 from "Bright's Disease," and 1 fr "Addison's Disease." The whole number was 3 more than 1879. The particulars are as follows:—

Males, 21; females, 25; total, 46, or 2.22 per cent. of all.

American parentage, 22; foreign parentage, 24.

For the twenty-five years, 1856-1880:—

Whole number, 526; annual average, 21.

Males, 307; females, 219; total, 526, or 1 55 per cent of all.

American parentage, 296; foreign parentage, 230.

Liver, Diseases of. The number of deaths in 1880, from diseases of the liver, was 22; 1 more than in 1879, and 1.06 per cent. of all from known causes.

Males, 12; females, 10; total, 22, or 1.06 per cent. of all.

American parentage, 11; foreign parentage, 11.

For the twenty-five years, 1856-1880, the numbers are:-

Whole number, 380; annual average, 15.

Males, 193; females, 187; total, 380, or 1.12 per cent. of all.

American parentage, 186; foreign parentage, 194.

Malformations. Eleven deaths from malformation were reported in 1880, the same number as in 1879. Eight cases were malformation of the heart; 2, of the spine, and 1 was not specified.

Males, 6; females, 5; total, 11, or 0.53 per cent. of all.

American parentage, 8; foreign parentage, 3.

For the twenty-five years, 1856-1880, the numbers are:—

Males, 125; females, 71; total, 196, or 0.58 per cent. of all.

American parentage, 126; foreign parentage, 70.

Marasmus. The number of deaths ascribed to this cause in 1880, was 35, which was 7 more than in 1879; 31 of the decedents were less than 2 years old.

Males, 19; females, 16; total, 35, or 1.69 per cent. of all.

American parentage, 18; foreign parentage, 17.

The statistics for the twenty-five years, 1856-1880, are as follows:—

Whole number, 653; annual average, 26.

Males, 348; females, 305; total, 653, or 1.92 per cent. of all.

American parentage, 329; foreign parentage, 324.

Measles. There were 3 deaths from measles in 1880; in 1879, there were none. This is a disease which has appeared in Provi-

dence, in a very irregular manner; in several of the years in which these records have been kept, there have been no deaths from this cause; in other years, a very few; while in only five of the twenty-five years has it prevailed to any extent. The following table shows the number of deaths in each year since 1856:—

1877 1	187016	1868 16	1856 1
1878	1871 1	1864 7	1857 2
1879 0	1872 7	1865 7	185840
1880	187898	1866 9	1859 0
_	1874 4	1867 0	1860 0
Total, 25 yrs.29	1875 0	1868 4	1861 6
	1876 0	1869 6	1862 1

The three deaths in 1880, were 0.15 per cent. of all from known causes; all were females of foreign parentage; one decedent was less than a year old; one, between 2 and 5 years; and one between 5 and 10 years. The following statistics are for the twenty-five years, 1856–1880:—

Males, 100; females, 110; total, 210, or 0.62 per cent. of all.

American parentage, 60; foreign parentage, 150.

Of the 210 decedents, 71.43 per cent. were of foreign parentage.

Meningitis, Cerebro-Spinal. The number of deaths from this cause in 1880 was 4, or 3 less than in 1879.

Males, 2; females, 2; total, 4, or 0.19 per cent. of all. American parentage, 1; foreign parentage, 3.

The first cases of this disease in Providence were reported in 1864; the number in each year since then is as follows:—

1864 2	1870 0`	1876 4
1865 2	1871	1877 7
1866 1	187210	1878 7
1867 0	187339	1879 7
1868 1	187410	1880 4
1869 0	1875 6	
		Total, 17 years112

Whole number from known causes, in 17 years, 26,192.

From cerebro-spinal meningitis, 112.

Males, 63; females, 49; total, 112,.or 0.43 per cent, of all.

American parentage, 53; foreign parentage, 59.

The number and percentage of deaths in each division of age and according to sex and parentage, were as follows for the seventeen years, 1864-1880 inclusive:—

AGES.	Under 1 Year.	1 and under 3.	2 and	5 and under 10.	10 and under 15.	16 and under 90.	90 and under 30.	80 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	Males.	Females.	American.	Foreign.	Total.
No.	80	•	15	28	81	7	13	10	2	-	1	8	49	S	. 69	118
PER CENT.	7.14	5.36	18.89	22.83	10.73	18.50	11.61	8.58	6.36	88.	88.	56.25	43.75	47.82	52.68	100.00

Of the 112 decedents, 66, or 58.93 per cent., were less than 15 years old.

Murder. Only one death from murder was reported in 1880, that of a foreign female between 30 and 40 years of age.

Old Age. The number of deaths ascribed to this cause in 1880, was 73, which was 9 more than in the previous year. Fourteen of the decedents were more than 90 years old, or 9 more than in 1879.

Males, 31; females, 42; total, 73, or 3.58 per cent. of all. American parentage, 36; foreign parentage, 37.

For the twenty-five years, 1856-1880, the statistics are:-

Whole number, 1,327; annual average, 53.

Males, 447; females, 880; total, 1,327, or 8.90 per cent. of all.

American parentage, 811; foreign parentage, 516.

Of the 1,327 decedents, 66.31 per cent. were females, and 61.12 per cent. of American parentage.

But a small proportion of those who die at an advanced age are reported under this head, as is shown by the following numbers. In 1880, there were 276 decedents who were more than 70 years old; 190 of American, and 86 of foreign parentage; this was 13.38 per cent. of all the decedents from known causes during the year.

In the twenty-five years, 1856-1880, the number of decedents of more than 70 years of age, was 3,749; 2,613 of American, and 1,136, of foreign parentage; 11.02 per cent. of all from known causes.

Paralysis. Though more properly a symptom of disease than a distinct disease, yet many deaths are reported each year under this head. In 1880, it was assigned as the cause of 30 deaths, the same number as in 1879.

Males, 17; females, 13; total, 30, or 1.45 per cent. of all.
American parentage, 21; foreign parentage, 9.

For the twenty-five years, 1856-1880, the numbers are:-

Whole number, 438; annual average, 18.

Males, 230; females, 208; total, 438, or 1.29 per cent. of all.

American parentage, 327; foreign parentage, 111.

The statistics for apoplexy and paralysis are generally given together in this connection; the following are the figures for each of the 25 years, 1856–1880:—

Years.	Apoplexy.	l'aralysis.	Total.	Years.	Apoplexy.	Paralysis.	Total
1856	21	2	23	1870	28	24	¥
1857		9	27	1871	83	28	a
1858	15	8	18	1872	81	20	51
1859	16	6	22	1873	86	21	57
1860	17	11	28	1874	36	\$3	39
1861	14	11	25	1875	41	35	%
1862	19	5	94	1876	44	24	6
1863	20	8	28	1877	50	24	74
1864		16	29	1878	43	24	#
1865	20	18	38	1879	56	30	85
1866		8	29	1880	48	20	78
1867		18	49				
1868	22	24	46	Totals, 25 y	ears724	438	1,161

The statistics of both causes, as to sex and parentage, for the whole period, are as follows:—

Whole number, 1,162; annual average, 46.

Males, 584; females, 578; total, 1,162, or 3.41 per cent of all.

American parentage, 838; foreign parentage, 829.

The decedents were 71.69 per cent. of American parentage, and 28.31 per cent. of foreign parentage.

Peritonitis. There were 14 deaths from peritonitis in 1880; the same number as in 1879.

Males, 5; females, 9; total, 14, or 0.68 per cent. of all. American parentage, 6; foreign parentage, 8.

The statistics for the twenty-five years, 1856-1880, are:-

Whole number, 229; annual average, 9.

Males, 101; females, 128; total, 229, or 0.67 per cent. of all.

American parentage, 128; foreign parentage, 101.

Pneumonia. There were 170 deaths from pneumonia in 1880, and 23 from congestion of the lungs; total 193, or 34 more than in 1879, and 53 more than in 1878. The deaths occurred as follows in the different seasons of the year:—

lst quarter62	3d quarter14
2d quarter	4th quarter43

Of the 193 decedents, 64, or 33.16 per cent. were under 10 years of age; and 80, or 41.45 per cent., were over 50.

Males, 94; females, 99; total, 193, or 9.35 per cent. of all. American parentage, 86; foreign parentage, 107.

The general statistics for the twenty-five years, 1856–1880, are as follows:—

Whole number, 2,509; annual average, 100.

Males, 1,286; females, 1,223; total, 2.509, or 7.37 per cent. of all.

American parentage, 1,253; foreign parentage, 1,256.

The following table gives the particulars as to the deaths from pneumonia for each of the twenty-five years, 1856-1880; giving the sex, parentage, season, age, whole number, and percentage of all deaths from known causes:—

Scarlatina. The epidemic of scarlatina, which was so severe in 1879, continued to prevail in 1880, through the first half of the year; after that time, though there were some deaths each month, the number diminished very rapidly. The whole number of deaths in 1880 was 243, or 11.77 per cent. of all; in 1879, there were 252 deaths. or 12.49 per cent. of all. The decedents in 1880, were divided as follows as to age: under 1 year, 21; between 1 and 2 years, 29; between 2 and 5 years, 95, or 39.09 per cent. of all from this cause; between 5 and 10 years, 73; between 10 and 15, 15; between 15 and 20, 3; between 20 and 30, 7.

.The deaths in the different seasons of the year were as follows:-

1st quarter115	8d quarter82
2d quarter 76	4th quarter20

The 115 deaths in the first quarter of the year were 47.33 per cent. of the whole number; the largest number in any one month was 50, in January; the largest number in any one ward was 94, in the tenth ward. The decedents were very evenly divided as to sex, there being only one more male than female; but as to parentage, 63.79 per cent. were of foreign, and 36.21 per cent. of American parentage.

The statistics for the year are as follows:-

Males, 122; females, 121; total, 243, or 11.77 per cent. of all.

American parentage, 88; foreign parentage, 155.

For the twenty-five years, 1856-1880:-

Whole number, 2,002; annual average, 80.

Males, 978; females, 1,024; total, 2,002, or 5.88 per cent. of all.

American parentage, 822; foreign parentage, 1,180.

The following table gives the particulars of sex, parentage, season, and locality, for each of the twenty-six years, 1855-1880:—

SCARLATINA IN PROVIDENCE, TWENTY-SIX YEARS, 1855-1880.

10.11.01.2 0.11.01.2	101	
3 % 8 %		
8.8 8.8		14 8 16 50 14 11 17 91
	342800004 -8-	342800004 -8-
1111 1110 1110 1110 1110 1110 1110 111	8 8 8 5 8 8 5 8 8 8 5 8 8 8 8 8 8 8 8 8	- 20 - 20 - 20 - 20 - 20 - 20 - 20 - 20

A comparison of the statistics of scarlatina, with those of diphtheria and croup, is found under the head of diphtheria.

Scrofula. There was but one death from scrofula in 1880; the decedent was a female of American parentage, between 2 and 5 years old. In 1879 there were 5 deaths from this cause.

Small-pox. There have been no deaths from small-pox in Providence since 1875. For the twenty-five years, 1856–1880, the statistics are as follows:—

Males, 58; females, 21; total 79, or 0.23 per cent. of all. American parentage, 48; foreign parentage, 31.

The particulars for each year are as follows:-

SMALL-Pox.

	Sı	EX.	PAREN	TAGE.	Whole		
	Males.	Females.	American.	Foreign.	Number. Percent		
356	7	2	5	4	9		
357	•••••			• • • • • • • • • • • • • • • • • • • •	•••••		
358	• • • • • • • • • • • • • • • • • • • •	•••••					
359	2	8	5		5		
960	4	1	8	2	5		
961		• • • • • • • • • • • • • • • • • • • •			•••••		
882	2	2	1	8	4		
363		•••••	• • • • • • • • • • • • • • • • • • • •	•••••	•••••		
364	7				7		
965	9	2		8	11		
966					••••••		
967	1		1		1		
968	1		1	•••••	1		
869	8		8	•••••	8		
870		2		9			
871		8	1	2	8		
872	9	4	9	4			
878	12	2	8		14		
					•••••		
875	1		•••••	1	1		
877		• • • • • • • • • • • • • • • • • • • •	•••••				
878		•••••					
879	•••••	•••••	•••••	••••	•••••		
880		•••••			***************************************		

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Skin, Disease of, was reported as the cause of one death in 1880; the decedent was a female of foreign parentage, under 1 year old.

Spine, Diseases of. Three deaths were reported from diseases of the spine during the year, 1880; this was 3 less than in 1879, and 0.15 per cent. of all from known causes.

Males, 2; females, 1; American parentage, 2; foreign parentage, 1.

Stomach, Diseases of. There were 10 deaths reported under this head in 1880, of which 8 were called "inflammation"; the number was 5 less than in 1879. The statistics of sex and parentage are:—

Males, 3; females, 7; total, 10, or 0.49 per cent. of all. American parentage, 3; foreign parentage, 7.

Suicide. There were but 2 cases of suicide in 1880, 4 less than in 1879; there have been only three years in the twenty-five years during which these records have been kept, when the number was so small; in 1856 there was but one case of suicide reported; in 1857, 2; in 1864, 1; in some years the number has been quite large, in 1877 amounting to 12 cases. In 1880, both cases were by cutting the throat; both decedents were females, of foreign parentage. The statistics for the twenty-five years, 1856-1880, are:—

Males, 103; females, 44; total, 147, or 0.43 per cent. of all.

American parentage, 81; foreign parentage, 66.

Tecthing. The number of deaths from this cause in 1880 was 16, or 3 more than in the previous year. The statistics of sex and parentage are:—

Males, 7; females, 9; total, 16, or 0.78 per cent. of all. American parentage, 4; foreign parentage, 12.

Tuberculosis. The number of deaths reported under this head in 1880, was 9, which was 11 less than in 1879.

Males, 3; females, 6; total, 9, or 0.44 per cent. of all. American parentage, 7; foreign parentage, 2. There are many tubercular diseases which might very properly be included under this general head; and the percentage of such a class would exceed that of any other. This was not specified as a separate head in these tables until the year 1867, since which time the deaths have been as follows:

Whole number from known causes in 14 years, 22,737.

From tuberculosis, 181; annual average, 13.

Males, 79; females, 102; total, 181, or 0.80 per cent. of all.

American parentage, 120; foreign parentage, 61.

Tumors. There were 6 deaths from tumors in 1880; 1 less than in 1879.

Male, 1; females, 5; total, 6, or 0.29 per cent. of all. American parentage, 5; foreign parentage, 1.

Those which were specified were 1 of the spleen; 1, ovarian; 3, uterine.

Unknown. The number of deaths in 1880, of which the cause was not stated, was 17, or 8 more than in 1879.

Males, 7; females, 10; total, 17, or 0.82 per cent. of all deaths.

American parentage, 9; foreign parentage, 8.

Twelve of the decedents were less than a year old.

The statistics of deaths from unknown causes, for the twenty-five years, 1856-1880, are as follows:—

Whole number, 550; annual average, 22.

Males, 284; females, 266; total, 550, or 1.59 per cent. of all.

American parentage, 185; foreign parentage, 365.

Of the 550 decedents, 66.36 per cent. were of foreign parentage.

STILL-BORN.

The following statistics are recorded concerning the still-born children in 1880:—

Number. The number of still-born children in 1880 was 121; 1 less than in 1879, and 4.61 per cent. of the 2,627 children born during the year, or 1 in 21.71.

Sex. The number of males was 76, or 13 more than in 1879; the number of females, 45, or 14 less than in the previous year.

Color. White, 111; colored, 10.

Locality.	Ward I14	Ward VII 11
	II 2	VIII 5
	III17	IX 17
	IV 7	X 22
	V12	
	VI14	Whole city121

Season. January, 9; February and March, 12 each; April, 9; May, 5; June, 14; July, 9; August, 5; September, 9; October, 11; November, 17; December, 9.

The number and percentage in each quarter of the year were as follows:—

Number.	Per cent.	Number.	Per cent.
lst quarter33	27.27	8d quarter28	19.01
3d quarter28	23.14	4th quarter87	30.58

Parentage. In the following table the first letter shows the nativity of the father; the second, that of the mother of the 121 still-born children in 1880.

A. and A 59	I. and A 1	B. A. and J 1
I. and I 23	E. and A 7	8. and I 1
E. and E 5	G. and A 2	A. and E 3
G. and G 1	B. A. and A 3	8. and E 1
B. A. and B. A 5	8. and A 1	A. and B. A 1
P. and P 2	A. and I 8 E. and I 8	E. and 8 1
8w. and 8w 1	E. and I 8	G. and Swiss 1
Total.		

A.—American; I.—Irish; E.—English; S.—Scotch; G.—German; B. A.—British American; P.—Portuguese; Sw.—Swedish.

The whole number of children born in Providence during the year 1880, was 2,627; of which number, the 121 who were still-born were 4.61 per cent. Of the 2,627 children born, 1,332, or 50.70 per cent., had American mothers; 804, or 30.61 per cent., had Irish mothers; and 491, or 18.69 per cent., had mothers who were of other foreign countries. Of the 121 still-born children, 65, or 53.72 per cent., were children of American mothers; 36, or 29.75 per cent., were children of Irish mothers; and 20, or 16.53 per cent., had mothers, who were natives of other countries.

The proportion of still-born children to the whole number born of mothers of each class, was as follows: of the 1,332 children of American mothers, 65, or 4.88 per cent., were still-born; of the 804 children of Irish mothers, 36, or 4.48 per cent., were still-born; of the 491 children of mothers of other foreign countries, 20, or 4.07 per cent.; to American mothers one still-born child in 20.49; to Irish mothers, one in 22.33; to other mothers, one in 24.55. The proportion of still-born children to the whole number of children borne by mothers of each class, has generally been largest among the Irish; but in 1879 and 1880, the proportion was largest among the American.

The number of children borne by American mothers in 1880, was 44 more than in 1879; but the percentage of these to the whole number of children born was 0.37 per cent. less; of still-born children, the number and proportion borne by American mothers in 1880 were both less than in the previous year. Of the children of Irish mothers in 1880, the number was larger than in 1879; the proportion to all the children born, slightly less; of still-born children in this class,

the number and proportion were both larger than in the previous year. Of children whose mothers were of various other nations, the number and proportion in 1880 were larger than in 1879; of still-born children, the number and proportion were also greater.

For the twenty-five years, 1856-1880, the statistics on this subject are as follows:—

Whole number of children born, 49,049.

Number of still-born, 2,848, or 5.81 per cent. of all; or one in 17.22. Of these 2,848 still-born children, 2,668 were white; 180, colored; 1,641, or 57.62 per cent., were males; and 1,207, or 42.38 per cent., females.

Reckoning the parentage according to the father's birth-place, 1,220 were American; 1,241 were Irish; and 380 were children of other foreign fathers; of 7 the parentage was not stated.

The following statement shows the whole number of children borne by mothers of different nativities, during the twenty-five years, 1856–1880 inclusive; also the number of still-born children to mothers of each class; and the percentage of the latter to the whole number of children:—

American mothers	Whole number of children.		Percentage of still-born. 5.31
Irish mothers	•	·•	
English, Scotch or Welsh mothers.	8,554	170	4.78
German mothers	998	29	2.92
British American mothers	1,628	69	4.24
Other foreign mothers	795	48	6.04
-	49,049	2,848	5.81

There are considerable differences in the numbers of still-born children, among the mothers of different nativities. The proportion of still-born is much the greatest among the children of mothers born in Ireland. Of all the children born of American mothers in Providence, during twenty-five years, 1856 to 1880 inclusive, one in 18.8 was still-born, while of the children of Irish mothers, one in 14.5 was still-born, and of children of English, Scotch, and Welsh mothers. one in 20.9; and of children of German mothers, one in 34.2 was still-born.

GENERAL SUMMARY.

The following recapitulatory table will give the general results of registration in Providence during the 26 years, 1855 to 1880 inclusive:—

TABLE XXI. BIRTHS, MARRIAGES, AND DEATHS.

YBAR.	Population.	Births.	To population, one birth in	Marriages.	Of population, one person married in	Deaths.	Of population, one death in
1855	47,785	1,600	99.86	686	87.56	991	48.23
1856	48,797	1,675	29.09	656	87.14	1,065	45.75
1857	49,457	1,666	29.30	614	40.27	925	58.46
1858	49,457	1,794	28.68	560	44.15	1,017	48.63
1850	50,100	1,598	81.45	616	40.86	899	55.72
1860	50,666	1,648	80.74	638	40.02	1,001	50.51
1861	50,686	1,795	29.87	547	46.31	1,051	48.21
1863	52,288	1,529	84.16	547	47.75	914	57.15
1863	58,024	1,307	40.56	610	48.46	1,214	48.67
1864	53,810	1,844	40.04	789	86.40	1,201	42.01
1965	54,595	1,298	42.06	709	88.50	1,211	45.08
1866	55,709	1,682	84.18	812	34.80	1,036	53.77
1867	56,824	1,624	84.99	856	88.28	960	59 19
1868	64,138	1,866	84.87	872	86.77	1,110	57.78
1869	66,522	1,819	86.57	980	85.76	1,256	59.96
1870	68,904	1,889	86.48	969	85.98	1,268	54.55
1871	78,011	1,960	86.74	948	88.18	1,254	57.42
1879	75,118	2,206	84.05	1,084	84.63	1,608	46.86
1878	78,225	2,128	36.76	1,150	84.01	1,719	45.51
1874	96,453	2,780	84.69	1,176	41.01	1,987	48.54
1875	100,675	2,864	87.79	1,079	46.65	1,915	59.57
1876	101,511	2,681	88.58	949	58.48	1,865	54.48
1877	101,000	2,548	89.73	958	52.99	1,938	59.13
1878	102,286	2,627	88.94	1,016	50.84	1,989	51.43
1879	108,579	2,522	41.07	1,071	48.85	2,096	51.12
1880	104,857	2,627	89.92	1.234	12.49	2,080	50.41
Average	69,551	1,948	85.70	844	41.90	1,366	50.84

BIRTHS AND DEATES BY FALENTAGE.

The following recognizating table shows the number of notice and deaths, by parentage, in Providence, it much if the twenty-my peace from 1856–1859 inclusive. In the times year, 1865, there was an excess of feaths over the further in the population of American parentage:—

TABLE XXX.

YEAR.	ARREST PARTIES.			FOLDES TARRESTANCE			
	Penths.	Brits.	Example of Bertin.	Teams	Rr:18	Farmer at his n	
1956	345	230	535	38.	:36	X	
1857	-	=	23	402	367	363	
1858	202	: = 1	339	125	56	atř	
1850	6	•	236	431	Sin	554	
1980	323	333	± .	€ ₹	SC	6	
1961	336	789	**	153	985	C	
1963	-	65	**	434	:00	423	
1963	36	355	_ _	Œ.	::39	362	
1864	60	£3	_ 	622	765	133	
1865	61	345	-85	544	732	228	
1966	523	736	223	123	946	2/3	
1867	206	725	197	433	200	867	
1868	596	862	286	514	364	479	
1869	627	936	330	(2)	208	354	
1870	68		968	ह्य	350	1 37%	
1871	643	200	337	611	200	369	
1879	833 :	1.455	. 388	771	เส		
1873	863	1,005	140	857	1.135	268	
1874	865	1,217	332	1.103	1,563	461	
1875	914	1,180	206	1,001	1.484	483	
1876	875	1,169	294	990	1.463	479	
1877	908	1,238	230	1.030	1,305	275	
1979	889	1,215	336	1.100	1.413	312	
1879	941	1,398	287	1,085	1,394	209	
1980	990	1,959	339	1,160	1,375	215	
25 years	17,017	22,343	5,396	17,569	96,706	9,144	

CENSUS OF PROVIDENCE.

Statement showing the nativity, sex, color, and total population. of the city of Providence by wards, according to the United States census of 1880.

W	Total	NATIVE	WHITE.	FOREIG	WHITE.	Colored.			
WARDS.	Population.	Males.	Females.	Males.	Females.	Males.	Females.		
<u> </u>	18,713	4,501	4,989	1,786	2,051	228	258		
11	5,002	1,596	1,812	248	806	280	308		
111	8,790	2,097	8,087	1,089	1,290	115	142		
ıv	6,201	2,202	2,353	649	851	70	76		
v	8,748	2.912	8,078	1,251	1,423	49	35		
v1	6,190	2,256	2,581	849	586	219	806		
vII	9,406	8,169	8,507	1,015	1,240	186	199		
vIII	12,816	4,615	5,112	1,199	1,563	185	192		
ıx	15,256	5,508	5,684	1,458	1,728	420	453		
x	18,810	5,717	5.576	8,666	8,831	7	18		
City	104,857	85,475	87.764	12,658	15.819	1,659	1,987		

CITY ORDINANCES.—CHAPTER XLVII.

REGISTRATION OF DEATHS.

SECTION 1. There shall be appointed by the City Council a sufficient number of persons to act as undertakers, removable at the pleasure of the City Council.

- SEC. 2. Whenever any person shall die in the city, the physician attending in his or her last sickness shall furnish to the undertaker attending the funeral, or to the city registrar, a certificate, giving the name of the person, date of death, and the disease or cause of his or her death.
- SEC. 3. No person shall bury, or place in a tomb, or remove from the city, or otherwise dispose of the body of any human being who shall die in the city, without first reporting the death to the city registrar, and obtaining a permit from him.
- SEC. 4. No permit shall be given as provided in section three, until the city registrar is furnished with the information in relation to the deceased person. required by the laws of the State, for record, so far as the same can be ascertained, together with the physician's certificate of the cause of death, whenever a physician has been in attendance, or a coroner's certificate, whenever a coroner's inquest has been held.

Whenever a permit for burial is applied for, in a case of death without the attendance of a physician, or if it is impossible to obtain the physician's certificate, the city registrar shall investigate the case so far as may be necessary, and when he has obtained satisfactory evidence in relation to the cause and circumstances of the death, he shall sign the certificate and give the required permit. If not satisfied in relation to the cause and circumstances of the death, or if, in his opinion, the public good requires it, he shall report the case to a coroner for investigation.

- SEC. 5. Whenever the body of a human being who has died out of the city shall be brought here for burial, the undertaker, or other person attending the funeral, shall furnish the report required in sections three and four, with the exception of the physician's certificate.
- SEC. 6. All funerals shall take place between sunrise and sunset, unless otherwise permitted or directed by the board of health. The top of every coffin deposited in the ground shall be at least three feet below the usual surface thereof.
- SEC. 7. The city registrar shall cause an abstract of the returns of deaths made to him to be published in all the newspapers which contract to do the city printing, monthly, and oftener if required by the board of health. He shall also, on or before the first day of April, annually, prepare and present to the city council a statement of the number of births, marriages and deaths which occurred in the city during the year ending with the thirty-first day of December next preceding, with such other information and suggestions in relation thereto as he may deem useful for the promotion of the public health, and other interests of the city.
- SEC. 8. No undertaker or other person shall bury or cause to be buried the body of any deceased person in the city, except in such grounds as are or may be designated as burying grounds, and authorized to be used as such.
- SEC. 9. Every person violating any provision of this chapter shall pay a fine of not less than five and not more than twenty dollars for each offence.

		·	
•			

TWENTY-SEVENTH ANNUAL REPORT

TPON THE

BIRTHS, MARRIAGES, AND DEATHS

IN THE

CITY OF PROVIDENCE:

FOR THE YEAR

1881,

By EDWIN M. SNOW, M. D.,

SUPERINTENDENT OF HEALTH AND CITY REGISTRAR.



PROVIDENCE;

PROVIDENCE PRESS COMPANY, PRINTERS TO THE CITY. > 1882.

• •

TWENTY-SEVENTH ANNUAL REPORT

UPON THE

BIRTHS, MARRIAGES, AND DEATHS

IN THE

CITY OF PROVIDENCE:

FOR THE YEAR

1881,

By EDWIN M. SNOW, M. D.,

SUPERINTENDENT OF HEALTH AND CITY REGISTRAR.



PROVIDENCE:

PROVIDENCE PRESS COMPANY, PRINTERS TO THE CITY.

THE NEW YORK
PUBLIC LIBRARY
118673

ASTOR, LENOX AND
TILDEN FOUNDATIONS.
1899.

THE CITY OF PROVIDENCE.

JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved June 80, 1882.]

RESOLVED, That the twenty-seventh annual report of the City Registrar, for the year ending December 81, 1881, be received, and that six hundred copies thereof, and of the communication accompanying the same, be printed in pamphlet form for the use of the City Council, under direction of the joint committee on printing.

A true copy: witness,

HENRY V. A. JOSLIN, CITY CLERK.

INTRODUCTION.

CITY REGISTRAR'S OFFICE, PROVIDENCE, May 18, 1882.

To the Honorable the City Council:

The report herewith presented upon the births, marriages, and deaths in Providence, during the year ending December 31, 1881, is my twenty-seventh annual report upon this subject.

During the year 1881, there were 2,803 births, 2,145 deaths, and 1,202 marriages, reported in Providence. The births were 176 more, the marriages 30 less, and the deaths 65 more than in 1880.

Estimating the population of the city, in the middle of the year 1881, at 107,500, we have the following proportions of births, marriages, and deaths to population: Births 2,803, or one in 38.35, or 26.07 in each 1,000; Persons married 2,404, or one in 44.72, or 22.36 in each 1,000; Deaths 2,145, or one in 50.11, or 19.95 in each 1,000.

Among the white population of the city, there was, in 1881, one birth in 38.65; one person married in 45.46; and one death in 50.91.

Among the colored population, there was, in 1881, one birth in 30.83; one person married in 30.63; and one death in 34.93.

The report herewith presented contains the usual tables of past reports, with some additional tables relating to important diseases in the city, and also tables relating to the census of the population.

EDWIN M. SNOW, M. D.,

Superintendent of Health and City Registrar.

CONTENTS.

The state of the s	PAGE
BIRTHS	1-1
Wards, months, sex and color, (Table I.)	
Births in connection with season, sex and locality, (Table II.)	1
Proportion of the sexes born at different seasons	
Color:—Number and proportion of the sexes, 20 years. Color:—Births, marriages and deaths in 26 years. Parentage of children born in each ward (Table III)	3
Parentage of children born in each ward, (Table III.)	4.16
Number and percentage of American and of foreign children, 25 years.	>11
Number of 1st 2d etc. children in 1881, and 1855-1880, (Table V.)	72
Average number of children to each mother, 27 years, (Table VI.)	11
Number of the child, with age of the mother, (Table VII.)	14-16
Plurality-births, statistics for ISSI	17.19
Color:—Births, marriages and deaths in 26 years. Parentage of children born in each ward, (Table III.). Number and percentage of American and of foreign children, 25 years. Parentage:—Mixture of different nations, (Table IV.). Number of 1st, 2d, etc. children in 1881, and 1855-1880, (Table V.). Average number of children to each mother, 27 years, (Table VI.). Number of the child, with age of the mother, (Table VII.). Plurality-births, statistics for 1881. Plurality-births, 27 years' summary.	11-28
MARRIAGES. Marriages at different seasons, 27 years, (Table VIII.) Birthplaces of parties married in ISSI, (Table IX.) Bride and groom born in same country, etc. Proportions of American, foreign, and mixed marriages, 27 years Marriages of colored persons	16-05
Marriages at different seasons, 27 years, (Table VIII.)	W-81
Birthplaces of parties married in 1881, (Table IX.)	13 m
Bride and groom born in same country, etc	96
Marriages of colored persons	5
Age of persons married in 1881, (Table X.)	3
Average age of persons married, 26 years. (Table)	3
Marriages of colored persons. Age of persons married in 1881, (Table X.) Average age of persons married, 26 years. (Table). Number of times married, (Table XI.) Denominational.	20
Nativity and residence	0.30
DEATHS Deaths to population in 1991, proportion to each clear	31-97
Deaths by sex, condition, color, nativity, parentage, and locality, (Table XII.). Deaths by season; numbers and proportions; recapitulation.	見
Dooths by sesson's numbers and proportions: reconitulation	23-31
Sex, condition, color, nativity, parentage, and locality of decedents,	
1881	25-11
Aggregate and average age by parentage, 1881, (Table XIII.)	91
Aggregate and average age by parentage, 26 years, (Tables Alv. and	18-11
Sex, condition, color, nativity, parentage, and locality of decedents. 1881. Aggregate and average age by parentage, 1881, (Table XIII.) Aggregate and average age by parentage, 26 years, (Tables XIV.and XV.). Difference between average age of American and foreign decedents, (Table) Percentages at different ages by parentage, 1881, (Table XVI.)	
(Table)	10
Percentages at different ages by parentage, 1881. (Table XVI.)	47
Percentages under five years of age by parentage, 26 years (Table)	44-45
Deaths by age and parentage in each month and quarter 1881. (Table	
XVIII.).	45-47
Causes of death, sex, age, parentage and percentages. (Table XIX)	45-34
(Table). Percentages at different ages by parentage, 1881. (Table XVI.). Percentages under five years of age by parentage, 26 years (Table). Percentages at different ages by parentage, 26 years, (Table XVII.). Deaths by age and parentage in each month and quarter, 1881, (Table XVIII.). Causes of death, sex, age, parentage and percentages, (Table XIX) Causes of death in connection with season, (Table XX.)	0070
Notes on the Causes of Death in 1881. Cholera Infantum; decedents by sex and parentage, 26 years. (Table). Consumption; sex and parentage. Consumption; change in rate of mortality. Consumption in Providence, 26 years, (Table). Crom: statistics for 1881.	57-8
Cholera Infantum; decedents by sex and parentage, 26 years, (Table)	- 61
Consumption; sex and parentage	61.5
Consumption; change in rate of mortality	61-0
Crown; statistics for 1821	W.
Croup: deaths in 26 years. (Table)	
Di rrhœn and dysentery; recapitulation for 42 years	70-1
Diphtheria; statistics for 1881	- 5
Diphtheria in Providence, 24 years (Table)	75-7
Heart, diseases of decedents by sex and parentage 26 years, (Table).	8
Meningitis, Cerebro spinal; deaths at each age, 18 years	- 5
Pneumonia; statistics for 1881 and 26 years, (Table)	87-5
Scarlatina; statistics for 1881 and 27 years, (Table)	0
Unknown causes	. 9
Still-born children in 1881	16-0
Consumption in Providence, 26 years, (Table). Croup; statistics for 1881. Croup; deaths in 26 years, (Table). Di rrhen and dysentery; recapitulation for 42 years. Diphtheria; statistics for 1881. Diphtheria in Providence, 24 years (Table). Fever, Typhoid, 1881, and 26 years, (Table). Heart, disease of, decedents by sex and parentage, 26 years, (Table). Meningitis, Cerebro spinal; deaths at each age, 18 years. Pneumonia; statistics for 1881 and 26 years, (Table). Scarlatina; statistics for 1881 and 27 years, (Table). Small pox; recapitulatory table, 26 years. Unknown causes. Still born children in 1881.	Y
Oneman in Deservations	
GENERAL SUMMARY:- Births, Marriages and Deaths; numbers and proportions, 27 years,	
(Table XXI.)	98
(Table XXI.)	10
XXII.)	644
Census of Providence, 1880, sex, nativity, parentage, and color	-

CITY REGISTRAR'S REPORT.

1881.

BIRTHS.

The number of children born in Providence, during the year 1881, was 2,803.

Table I. gives the number of births in each ward of the city, and in the whole city, in each month and quarter of the year; also, for the city, the number of each sex, and the number of colored children in each month and quarter:—

TABLE I. BIRTHS, 1881. Wards, Months, Sex and Color.

NEW A		WARDS.									Whole City.			Colored children in- cluded in the preceding.		
MONTHS.	r.	п.	m.	ıv.	v.	vi.	VII.	viii	ıx.	x.	Males.	Females.	Total.	M.	¥.	Total.
January February March	28 32 37	8 6 12		8 7 14	17 12 17	7 9 8	20 17 12	22 23 16	24 35 28	52 45 63	106 112 120	100 96 100	206 208 220	3 6 2	10 2	16
Ist Quarter	97	26	55	29	46	24	49	61	87	160	338	296	634	11	16	27
April May June	34 34 32	4 7 10	31 22 22	6 14 20	21 23 16	7 12 12	21 24 22	26 24 31	38 34 49	44 47 43	125 111 133	107 130 124	232 241 257	3 5 11	5 6 4	11 15
d Quarter	100	21	75	40	60	31	67	81	121	184	369	361	730	19	15	34
July August September	24 22 19	9	22 24 28	5 11 10	16 22 16	11 9 13	25 24 20	26 26 43	33 37 41	61 49 54	115 117 132	110 116 116	225 283 248	6 6 12	6	12 12 16
2d Quarter	65	15	.74	26	54	33	69	95	111	164	364	342	706	24	16	40
October November December	33 30 29	7.85	21	10 13 10	17 29 19	16 6 12	16 21 21	88 25 22	48 29 28	58 62 53	140 139 112	129 105 108		2	811	18
th Quarter	92	20	68	33	65	34	58	85	105	173	391	342	733	10	10	20
Whole Year	354	82	272	128	225	122	243	399	424	631	1,462	1,341	2.808	64	57	12

The number of births in 1881 was 176 more than in 1880, and much larger than in any previous year. In the fifth, seventh and ninth wards, the number of children born in 1881 was 29 less than in 1880; in the remaining seven wards, the number was 205 more, the greatest increase in any one ward being 65 in the tenth ward. The population of the city in the middle of the year 1881 was estimated at 107,500; the ratio of births, therefore, was one in 38.35 d the population, or 26.07 of births to each thousand inhabitants.

Season. There were four months in 1881 when the number of births was less than in the corresponding months of 1880, namely: January, March, July and December; in each of the remaining eight months the number in 1881 was greater. In each quarter of 1881 except the first, the number was larger than in the previous year. The largest monthly total in 1881 was 269 in October, he smallest, 206 in January. The largest quarterly total was 783 in the fourth quarter, the smallest, 634, in the first. In the first amonths of the year, the number of births was 1,364, or 89 months in the first half of the year 1880; in the last six months, the number was 1,439, or 87 more than in the previous year. The difference between the number of births in the first half of the year and that in the last half was 75; in 1880 it was 77.

The following statement shows the number of births in each quiter of the year 1881; the aggregate number in the corresponding seasons of the twenty-five years, 1856–1880, and the number for the twenty-six years, 1856–1881, inclusive:—

In the twenty-seven years, 1855-1881 inclusive, the aggregate monthly totals, and the daily average for each month, have been at follows:—

24,1354,422 6 5.10 5.28	
6 5.10 5.28	5.83
. Oct. Nov.	Dec.
04,6414,523	4,746
	7 5.54 5.58 verage, 5.42.

The largest aggregate number of births in any month was in December; the smallest, in February; the highest daily average was in June; the lowest, in January. There were six months in which the daily average was above the general average of the whole period, namely: March, June, September, October, November and December; in the other months the daily average was less than the general average.

Sex and Locality. The number of children of each sex, born in each ward of the city, in each quarter of the year, and in the whole year, is shown by the following table:—

TABLE II. BIRTHS, 1881. Wards, Sex, and Season.

	1st Quarter.			2d Quarter.			3d Quarter.			4th Quarter.			Whole Year.		
WARDS.	M.	F.	Total.	M.	F.	Total.	М.	F.	Total.	м.	F.	Total.	м.	F.	Total
I	51	46	97	45	55	100	28	37	65	46	46	92	170	184	354
II	11	15	26	7	14	21	9	6	15	12	8	20	39	43	82
11	29	26	55	40	35	75	42	32	74	35	33	68	146	126	279
ıvv	20	9	29	24	16	40	15	11	26	19	14	33	78	50	128
v	20	26	46	35	25	60	25	29	54	29	36	65	109	116	225
VI	13	11	24	16	15	31	18	15	33	16	18	34	63	59	122
VII	21	28	49	34	33	67	33	. 36	69	31	27	58	119	124	248
νш	34	27	61	42	39	81	54	41	95	48	37	85	178	144	322
IX	49	38	87	57	64	121	58	53	111	54	51	105	218	206	424
X	90	70	160	69	65	134	82	82	164	101	72	173	342	289	631
Whole City	238	296	634	369	361	730	364	342	706	391	342	733	1,462	1,341	2,803

Proportions of the Sexes. There were 121 more male than female children born in Providence in 1881. In the month of May alone, the number of female children was the larger; in the first, second, fifth and seventh wards, there were more female children born; in the other wards, more males. The whole number of children born was 2,803; of males, 1,462; of females, 1,341; a proportion of 109.0 males to 100 females; or 52.16 boys and 47.84 girls in each 100 children born.

The number and proportion of each sex, in each quarter of the year, were as follows:—

Males. Females. Males. Females. Males. Females.

Jan.-March.....388 and 296, or 114.2 to 100, or 53.31 to 46.69 in each 100 born-April-June.....369 and 361, or 102.2 to 100, or 50.55 to 49.45 in each 100 born-July-Sept......304 and 342, or 106.4 to 100, or 51.56 to 48.44 in each 100 born-

Oct.-Dec......391 and 342, or 114.3 to 100, or 53.34 to 46.66 in each 100 both.

Whole year....1,462 and 1,341 or 109.0 to 100, or 52.16 to 47.84 in each 100 both.

The proportions in the first and fourth quarters of the year are almost the same, and the excess of males much greater than in the second and third quarters. The aggregate number of children, and the proportion of each sex in each quarter of the year, for the twenty-six years 1856–1881 inclusive, are given below. The whole number of children born during this period was 51,852; males, 26,675; females, 25,177:—

Males. Females. Males. Females. Males. Females.

Jan.-March...6,522 and 6,062; or 107.6 to 100; or 51.83 and 48.17 in each 100 born.

April-June...6,649 and 6,269; or 106.1 to 100; or 51.47 and 48.53 in each 100 born.

July-Sept....6,555 and 6,350; or 103.2 to 100; or 50.79 and 49.21 in each 100 born.

Oct.-Dec....6.949 and 6,496; or 107.0 to 100; or 51.68 and 48.32 in each 106 born.

Whole period 26.675 and 25,177; or 105.9 to 100; or 51.44 and 48.56 in each 100 born.

In the twenty-eight years, 1854–1881 inclusive, the number of children born in the city was 54,821; males, 28,193; females, 26,628. The number and proportion in each year, and in the whole period, were as follows:—

```
Males.
            Females. Males. Females. Males.
                                                Females.
- 711 and
              658; or 108.0 to 100; or 51.91 and 48.09 in each 100 born.
   807 and
              703; or 101.7 to 100; or 50.41 and 49.56 in each 100 born.
   891 and
              784; or 113.6 to 100; or 53.19 and 46.81 in each 100 born.
   872 and
              816; or 106.8 to 100; or 51.66 and 48.34 in each 100 born.
   890 and
              834; or 106.7 to 100; or 51.62 and 48.38 in each 100 born.
   825 and
              768; or 107,4 to 100; or 51.79 and 48.21 in each 100 born.
    809 and
              839; or 96.4 to 100; or 49.09 and 50.91 in each 100 born.
   936 and
              789; or 118.6 to 100; or 54.26 and 45.74 in each 100 born.
   791 and
              738; or 107.2 to 100; or 51.73 and 48.27 in each 100 born.
   689 and
              618; or 111.4 to 100; or 52.71 and 47,29 in each 100 born.
- . 663 and
              681; or 97.3 to 100; or 49.33 and 50.67 in each 100 born.
. 691 and
              607; or 113.8 to 100; or 53.24 and 46.76 in each 100 born.
. 849 and
              783; or 108.4 to 100; or 52.02 and 47.98 in each 100 born.
. 830 and
              794; or 104.5 to 100; or 51.11 and 48.89 in each 100 born.
-, 944 and 922; or 102.4 to 100; or 50.59 and 49.41 in each 100 born.
. 945 and 874; or 108.1 to 100; or 51.95 and 48.05 in each 100 born.
-. 967 and 922; or 104.8 to 100; or 51.19 and 48.81 in each 100 born.
-. 956 and 1,004; or 95.6 to 100; or 48.78 and 51.22 in each 100 born.
-. 1,079 and 1,127; or 95.7 to 100; or 48.91 and 51.09 in each 100 born.
.. 1,109 and 1,019; or 108.8 to 100; or 52.11 and 47.89 in each 100 born.
-. 1,408 and 1,372; or 102.6 to 100; or 50.65 and 49.35 in each 100 born.
.. 1,390 and 1,274; or 109.1 to 100; or 52.18 and 47.82 in each 100 born.
-- 1,359 and 1,272; or 106.8 to 100; or 51.65 and 48.35 in each 100 born.
-. 1,302 and 1,241; or 104.9 to 100; or 51.20 and 48.80 in each 100 born.
-. 1.358 and 1,269; or 107 0 to 100; or 51.69 and 48.31 in each 100 born.
.. 1,297 and 1,225; or 105.9 to 100; or 51.43 and 48.57 in each 100 born.
-. 1,363 and 1,264; or 107.8 to 100; or 51.88 and 48.12 in each 100 born.
-.. 1,462 and 1,341; or 109.0 to 100; or 52.16 and 47.84 in each 100 born.
rs.28,193 and 26,628; or 105.9 to 100; or 51.43 and 48.57 in each 100 born.
```

tor. The number of colored children born in Providence in 1881 121; this was 30 more than in 1880, and a much larger number in any previous year. The colored population of the city in middle of the year was estimated to be 3,737; this gives the of births in this class as one in 30.88 of the population, or 3 births in each thousand of the colored inhabitants. The ratio c whole number of births to the total population of the city has been given as one in 38.35, or 26.07 births in each thousand itants. The white population of the city was 103,763, and the er of white children born 2,682; the ratio, one birth to 38.65

persons or 25.85 births to each thousand of the white inhabitants. The number of colored children born in each ward of the city was as follows: First ward, 15; Second, 10; Third, 13; Fourth, 4; Fifth, 2; Sixth, 13; Seventh, 12; Eighth, 17; Ninth, 35; Teuth, 0. The number in each month: January, 7; February, 16; March, 4; April, 8; May, 11; June, 15; July, 12; August, 12; September, 16; October, 13; November, 3; December, 4; Total, 121. The number of males was 64; of females, 57; a proportion of 112.3 males to 100 females, or 52.89 males and 47.11 females in each 100 children born.

In the twenty-six years, 1856-1881 inclusive, the number of ordered children born in Providence was 2,015; 1,074 males and 31 females; the number and proportion of each sex, in each year, and for the whole period were as follows:—

```
Year. Total. Males. Females. Males. Females. Males. Females.
1856..... 50..... 26 and 24; or 108 3 to 100; or 52.00 and 48.00 in each 100 bors
1857 ..... 31 ..... 17 and 14; or 121.4 to 100; or 54.84 and 45.16 in each 100 bots
1858 . ... 51 ..... 25 and 26; or 96.2 to 100; or 49.02 and 50.98 in each 100 bors
1859..... 36..... 20 and 16; or 125.0 to 100; or 55.56 and 44.44 in each 100 hors-
1860..... 54..... 31 and 23; or 134.8 to 100; or 57.41 and 42.59 in each 100 bors.
1861..... 51..... 28 and 23; or 121 7 to 100; or 54.90 and 45 10 in each lothers
1862..... 61..... 26 and 35; or 74.3 to 100; or 42 62 and 57.38 in each 100 both
1863..... 32..... 16 and 16; or 100.0 to 100; or 50.00 and 50.00 in each 100 born-
1864..... 28..... 25 and 13; or 192.3 to 100; or 65.79 and 34.21 in each 100 borns
1865..... 46..... 27 and 19; or 142.1 to 100; or 58.70 and 41.30 in each 100 bots
1866..... 67..... 39 and 28; or 139 3 to 100; or 58.21 and 41.79 in each 100 bottom
1867..... 75..... 38 and 37; or 102.7 to 100; or 50.67 and 49.33 in each 100 bern-
1868.... 90..... 49 and 41; or 119.5 to 100; or 54.44 and 45.56 in each 100 bons-
1869..... 85..... 51 and 34; or 150.0 to 100; or 60.00 and 40.00 in each 100 bors-
1870..... 104..... 56 and 48; or 116.7 to 100; or 53.85 and 46.15 in each 100 lens
1871 ..... 91..... 50 and 41; or 121.9 to 100; or 54.95 and 45.05 in each 100 borns
1872..... 99..... 51 and 48; or 106.2 to 100; or 51.52 and 48.48 in each 100 both
1873 ..... 112...... 62 and 50; or 124.0 to 100; or 55.36 and 44.64 in each 100 bors
1874..... 109..... 55 and 54; or 101.9 to 100; or 50.46 and 49.54 in each 100 bors
1875..... 101..... 49 and 52; or 94.2 to 100; or 48.51 and 51.49 in each 100 bers.
1876..... 108..... 48 and 60; or 80 0 to 100; or 44.44 and 55.56 in each 160 bots
1877..... 104..... 60 and 44; or 136.4 to 100; or 57.69 and 42.31 in each 100 bors.
1878..... 104..... 52 and 52; or 100 0 to 100; or 50.00 and 50.00 in each 100 bers
1879..... 104..... 58 and 46; or 126.1 to 100; or 55.77 and 44.23 in each 100 bots
1880..... 91 ..... 51 and 40; or 127.5 to 100; or 56.04 and 43.96 in each 100 both
1881..... 121..... 64 and 57; or 112.3 to 100; or 52.89 and 47.11 in each 100 both
26 years.2,015.....1,074 and 941; or 114.1 to 100; or 53.30 and 46.70 in each 100 born
```

The following figures show the number of births, marriages and deaths among the colored population of Providence in 1881, in the twenty-five years previous, and in the whole period of twenty-six years, 1856—1881 inclusive:—

Twenty-five years, 1856-1980	1,894 births.	928 marri	ages1.995 deaths.
1881	121 births	61 marri	ages 107 deaths.
Twenty six years, 1856-1861	 2,015 births.	989 marri	 ages2,103 deaths.

In the year 1881, the proportion of deaths to births in this class was 88.4 to 100; for the whole period of twenty-six years, the proportions were 104.4 deaths to 100 births.

It is quite unusual to find so large an excess of births over deaths in this class of the population, as in 1881. Though there have been eleven years, in the course of the twenty-six, in which there were more births than deaths, the excess, with one or two exceptions, has been quite small; and taking the whole period together, there have been 87 more deaths than births. The number of births, marriages and deaths among the colored population of Providence, in each year since 1856, has been as follows:—

		 989 marriages	
881	121 births	61 marriages	107 death
880	91 births	57 marriages	96 death
		43 marriages	
		54 marriages	
		46 marriages	
		38 marriages	
		48 marriages	
		60 marriages	
		49 marriages	
		54 marriages	
		46 marriages	
		52 marriages	
		60 marriages	
		58 marriages	
		35 marriages	
		36 marriages	
		23 marriages	
		12 marriages	
		55 marriages	
		13 marriages	
		15 marriages	
		15 marriages	
		9 marriages	
		37 marriages	
		14 marriages 90 marriages	

PARENTAGE.

The parentage of the children born in Providence in 1881 is shown by the following table, which gives the number of children, according to parentage, born in each ward; also the percentage of each class, in the whole city. Where the parents were of different foreign nations, the parentage is given according to the birth-place of the father.

TABLE III. BIRTHS, 1881. Wards and Parentage.

				,	₩ A E	DS.					CH7.	h 100 here
PARENTAGE.	ī.	n.	III.	ıv.	v.	VI.	vn.	VIII	ıx.	x.	Whole City	In each
American	148	55	96	49	65	81	112	185	209	143	1,141	40.7
Irish	80	7	73	83	55	16	58	45	88	195	650	23.1
English, Scotch and Welsh	22	5	16	7	28	2	18	27	84	108	202	9.3
German	4	2			12	1	2	4	11	21	57	2.0
British American	18	4	7	2	16	8	5	9	10	15	84	3.0
Portuguese			24				1	••••	8	2	30	1.0
Other Foreign	14	1	5	15	5	1	10	••••	10	7	68	2.6
Total Foreign	183	19	125	57	111	28	94	85	156	848	1,151	41.0
American father and For- eign mother	28	8	24	10	27	8	17	25	80	50	224	7.9
Foreign father and American mother	45	8	27	12	32	10	20	27	29	92	987	10.24
Ward totals	854	82	279	128	225	122	243	323	494	631	2,808	100.00

The "other foreign" mentioned in the table were as follows: in the First ward, 12 of Russian parentage, 1 Austrian, 1 Italian; in the Second, 1 Russian; in the Third, 2 Russian, 1 Swedish, 1 Spanish, and 1 of St. Helena Island; in the Fourth, 12 Italian, 2 Swedish, 1 French; in the Fifth, 2 French, 1 each Swiss, Norwegian and Swedish; in the Sixth, 1 Italian; in the Seventh, 10 Italian, in the Eighth, none; in the Ninth, 7 Swedish, 1 each French, Danish, and of St. Helena Island; in the Tenth, 4 Italian, 2 French, 1

Bohemian. Thus there were in all the city 28 of Italian parentage; 15 of Russian; 11 Swedish, 6 French; 2 of St. Helena; 1 each of Norwegian, Austrian, Bohemian, Danish, Swiss, and Spanish.

In the Third, Fourth, Fifth and Tenth Wards, the number of children of purely foreign parentage was larger than the number of American parentage; in the whole city the number of foreign was 10 more than the number of American parentage.

The following table gives the number and percentage of children of American, foreign and mixed parentage, in each of the twenty-six years, 1856-1881; and the number and percentage for the whole time:—

		•	Amer. father.	For. father.
	American.	Foreign.	For. mother.	Amer. mother
Year.	Number. Per cent. N	lumber. Per cent. N	umber. Per cent	. Number. Per c
356	. 76542.09	85751.17	553.22	588.
857	65038.51	98755.51	513.02	502.9
1856	. 71241.30	88651.39	583.36	683.
1659	. 59937.60	88755.69	598.70	498.0
1880	. 66140.11	85752.00	694.19	618.
1861	. 65487.91	91653.10	864.99	694.0
1862	. 58938.52	79952.26	744.84	674.3
1863	. 46735.73	71154.40	624.74	675.
1864	. 52038.69	69952.00	6)4.47	634.
1865	. 47836.67	68452.70	70 5.89	68 5.
1806	. 65640.20	82150.30	70 4.29	855.
1867	. 64539.73	. 79148.70	804.93	1086.
1988	. 77941.75	. 86846.51	1035.52	1166.
1889	. 82245.19	. 77642.16	1045.72	117
1870	. 800 42.85	. 86245.64	904.76	1877.
1871	. 87944.85	82842.21	1015.15	1527
1872	. 90741.12	.1,00245.42	1285.80	1697
1873	. 86940.84	. 94144. 22	1346.29	1848
1674	.1.06888.42	1,32647.70	1495.80	2378
1875	. 99987.50	.1,25647.15	1816.79	2288
		.1,23046.75		
1877	.1.054 41.45	.1,06741.96	1847.23	2389
1878	.1.03589.40	.1,17444.69	1806.85	2389
		.1,08541.04		
		.1.09141.52		
	•	.1,15141.06		

Of children of American parentage, the percentage is a little larger than in 1880, and the number larger than in any previous year; of children of foreign parentage, the percentage is slightly less than in the preceding year, although the number is larger than it has been since 1878; of children of American fathers and foreign mothers, the number and proportion are greater than in 1880; of children of foreign fathers and American mothers, the number is greater and the percentage less than in 1880.

The following table gives the percentage of children of each nationality, born in Providence, in each of the twenty-six years, 1856-1881:—

			English.		British.	Othe
	American.	Irish.	8. & W.	German.	American. Port	uguese. for'n
	Per cent.	Per cent.			Per cent. Per	cent. Per ct.
1856	42.09	41.79	4.84	2.27		2.27
1857	38.51	44.32	5.92	8.73		1.84
1858	41.30	42.98	4.82	2.26		1.8
1859	87.60	45.56	5.34	8.26		1.5
1880	40.11	40.96	6.53	2.85		1.64
1861	87.91	42.61	6.03			1.97
1802	88.52	41.58	6.02	8.07		1.64
1863	35.73	48.08	6.35	2.98		1.96
864	88.69	41.86	6.55	2.75		1.86
1865	36.67	42.99	5.93	1.85		1.90
863	40.20	89.53	6.25	9.21		9.89
867	39.72	37.81	5.66	8.02		2,21
868	41.75	86.01	6.48	1.88	1.07	1.07
869	45.19	84.14	4.01	1.81	1.82	1.86
870	42.35	84.66	6.19	1.96	1.91	90
871	44.85	82.95	4.51	1 63	2.09	1.78
873	41.12	82.46	5.67	9.77	9.31	2.31
873	40.84	80.50	7.19	1.88	2.49	2.16
874	. 8 8.42	82.63	8.13	2.84	2.191	.22 1.19
875	87.50	81.58	7.69	2.85	2.67	.981.45
876	38.27	81.05	7.75	2.59	2.89 1	.23 1.25
877	41.45	28.08	6.96	1.96		.10 1.30
878	89.40	29.65	6.74	2.55	2.861	181.71
•••					2.54	
					8.271.	
••••					3.00 1.	******

The British American were included in the "other foreign" until

1868, and the Portuguese until 1874. The highest percentage of American parentage in any year was 45.19 in 1869; the lowest, 35.73, in 1863; of Irish parentage, the highest percentage was 45.58 in 1859, and the lowest, 23.19, in 1881: while of English or Scotch parentage, the largest percentage was in 1881, the least, 4.01, in 1869.

The parentage of the children born in 1881 is shown more definitely in Table IV., which gives the birth-places of both parents:—

	P	RTII-	-PLA	CES O	F MO	THE	RS.	pirth
BIRTH-PLACES OF FATHERS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number secon to the father's birth place.
United States	1,141	135	56	3	25		5	1,365
Ireland	132	615	24		11			782
England, Scotland and Wales	78	35	207	1	17		2	340
Germany	21	3	2	43	2		7	78
British America	39	11	3		70			123
Portugal and Western Islands	3	2	1			27		33
Other Countries	14	8	- 4	3	1	1	51	82
Whole number according to the mother's birth-place	1,428	809	297	50	126	28	65	2,603

TABLE IV. BIRTHS, 1881. Parentage.

The 1,141 children whose parents were both American, were 40.71 per cent. of all the children born during the year; this was 93 more children, and 0.81 per cent. more than in 1880. There were also 224 children of American fathers and foreign mothers, which made 1,365 children with American fathers, or 48.70 per cent. of all; this was 113 more children, and 1.03 per cent. more, than in 1880. The children of foreign fathers and American mothers were 287, which gives a total of 1,428 children of American mothers, or 50.95 per cent. of all the children born; this was 96 more children, and 0.24

FIRM

Bixth

Reventh

Biahth

Ninth

Tenth

Eleventh "

per cent. more, than in the preceding year. Of the two classes of children of mixed parentage, the number was 511, or 18.23 per cent. There were 1,151 children of purely foreign parentage; this was 41.06 per cent. of the whole number; 60 more children, but 0.46 per cent. less than in 1880. Of these 1,151 children, 1,013 had parents who were of the same foreign country; this was 88.01 per cent. of the children of foreign parentage, and 36.14 per cent. of all the children born during the year. The 138 children whose parents were natives of different foreign countries were 11 99 per cent. of all the children of foreign parentage; and 4.92 per cent. of the whole number of children born during the year.

NUMBER OF THE CHILD.

Table V. shows how many of the 2,803 children born in 1881 were the first, second, third, etc., children of their mothers; also the same facts in regard to the 50,649 children born during the twenty-six years preceding, 1855-1880, inclusive:-

NUMI	ER	OF (CHILD.	1881.	26 years, 1855–1880.	NUMBE	R C	o r ci	IILD.	1881.	29 years, 1835–1880.
First ch	ild c	f the	mother	624	12,436	Twelfth ch	1d (ofthe	moth'r	15	200
Second	66	"	"	558	10,362	Thirteenth	44	44	44	5	83
Third	16	**	"	414	8,113	Fourteenth	**	46	66	5	40
Fourth	4.	44	•	846	6.119	Fifteenth	**	46	••	7	16

8,324

1.447

940

611

833

4,440 Sixteenth

Eighteenth "

Nineteenth "

Whole number of chil-

802

173

146

58

86

14

**

"

46

46

6

7

1

50,649

2,803

TABLE V. BIRTHS, 1881. Number of the Child.

The 624 children, who were the first children of their mothers, were 22.26 per cent. of the 2,803 children born during the year; this was 1.04 per cent. less than the same class of children in 1880. Of the 50,649 children born in the 26 years, 1855-1880, 12,436, or 24.55 per cent., were the first children of their mothers.

Average number of children to each mother. There were 23 twin births in 1881; the number of mothers, therefore, was 2,780. From Table VII. we see that these mothers had borne in all 10,111 children, which would average 3.64 children to each mother.

The next table shows the average number of children to each mother, in each of the twenty-seven years, 1855 to 1881, inclusive. This table shows the number of children the mothers included in the figures had borne, previous to the end of the year 1881; but it does not show the average number of children the same mothers may have in their life-time:—

TABLE VI. BIRTHS, 1855 TO 1881.

Average number of children to each mother.

		No. of	Aggregate.
earr.	Mothers.	Children. No	o. of Children. Average.
855	1,588	1.600	5,2143.28 children to each mothe
854	1,658	1,675	5,562 3.35 children to each mothe
837	1,664	1,688,	5,7363.44 children to each mothe
858	1.706	1,724	5,685 3.33 children to each mothe
859	1,570	1,598	5,6183.58 children to each mothe
860	1.627	1,648	5.8413.59 children to each mothe
961	1,694	1.725	6,0173.55 children to each mothe
963	1,512	1,529	5,3223.52 children to each mothe
863	1,396	1,307	4.717 3.64 children to each mothe
3 61	1,327	1.844	4,8873.68 children to each mothe
98 3 .	1,279	1,298	4.6253.60 children to each mothe
266	1,614	1,633	5,7443.56 children to each mothe
867	1,595	1.634	5,497 3.50 children to each mothe
358	1,841	1,866	6,338 3.44 children to each mothe
969	1,797	1,819	5,954 8.30 children to each mothe
870	1,850	1,889	6,145 3.31 children to each mothe
871	1,937	1,960	5,420 3.31 children to each mothe
879	2,174	2,206	7,3463.38 children to each mothe
878	2,104	2,128	6,7313.20 children to each mothe
874	2,738	3,780	8,9603.27 children to each mothe
875	2,634	2,664	8,9773.41 children to each motion
876	2 604	2.631	8,9143.42 children to each mothe
877	2,516	2,543	8,7053.46 children to each mothe
			9,417 3.63 children to each mothe
			8,9423.58 children to each mothe
880	2,600	2.627	9,2863.61 children to each mothe
			10.1113.64 children to each mothe

Table VII. gives the number of the child in connection with the age of the mother; shows how many first, second, third, etc., children were borne in 1881 by mothers at the ages given; the number of mothers of each age, and the aggregate and average number of children they had borne:—

Table VII. Births, 1881. Number of child with age of mother.

				NUM	BER	OF	СНІ	LD		_		_	_	_		_	each	1881.	Total children they had borne.	Average number of children to mothers at each use.
Age of the																	mothers at	Children they bore in 1	ey had	verage number of child to mothers at each use.
															•		oth	Ž	Ē	1
MOTHERS													П					hey	E E	E
IN YEARS.																	Number of Age.	E	ğ	1 2 5
										اند	غ	÷	اغا	ے	نے	ė	amb	Į.	7	2 5
	1st.	3 d.	3d.	ĘÞ.	Stb.	etb.	7t b.	Stb.	91 D.	10Ch	11th	lth.	13th	14th.	15th.	l Ifth.	Na	కే	Ē	*
6	5				<u></u>				:	-	-		$\overline{\cdot \cdot}$:	-:	-	5		5	1.00
7	18	1						•-	••				••	••	••	••	14	14	l .	1
8	18	8		••••		••••	•••	••	••		••	• •	••	••	••	•••	16	16	l	
9	82	10	2	•••		••••	•••	••	••	••	••	••	••	•••	••	••	44	44	56	
0	40	18	5	••••		•••	••••		••	••	•••	•••	•••	••	••	• •	56	58	ध	1.40
1	45	. 29	7	. 2	••••	••••	••••	٠٠,	•••	••	••	••		• •	••	••	88	88		1.59
3	68	41	8	8		••••	••••		•••	•••	••	•••		••	••	•••	120 149	1 2 0		1.77
3	72 •50	•48 •45	24 29	14	5	1	••••	1	•••	•••	٠.	••		••	••	••	152	154	328	L
4 5	•78	61	83	12	5	1	••••		•		•	•••		•••			184	185	372	
A	48	58	*41	23	ı	_				ı							180	181	476	2.64
7	+36	38	87	19	ı	4	8										155	156	1 .	2.13
8	181	89	84	*48		13			1								186	190	806	8.27
9	21	87	31	84	+27	11	7	1	1								169	170	586	8.47
0	28	87	38	29	184	21	6	8	8				١				192	194	706	3.68
1	8	17	21	27	+28	7	6	3	8			••	.	••	٠.		113	114	450	3.98
2	11	28	22	•24	+2K	17	15	4	1	1	••	• •		••		••	147	149	1	4.13
B	5	14	121	24	19	14	16		1	2	2		•••		••		121	122	I I	4 65
4	4	7	15	10	_	7			4	7	••	1	••	••	••		90	90		5.29
5	12	18	14	19	ı	18		19	7	6	•••	1	••	••	••	•••	146	_		5.26
6. 	1	8	7	•17	i .	15			7	2		8	••	••	••	•••	101	108		6.00 6.15
7	1	4	8	18	ı	•8			9	4	8	2	١.	•••	•:		82	8 8	1 1	6.46
8	2	4	5	12 7	12 8	18 +7	18 6		7 6	2	1 8	8	1	••	1	•••	87 57	58	≀ I	5.74
9 O	8	1	8	+1	6	6	6	1 -	5	6	1	••	1	,	1	••	50	51	1	7.36
1	ı	1	8	5	1	1	2		1	٠	•	1		. *	i		20	20	1	5.95
3	ĺ		l	1	i		1		i	8	1	•	2	2	ì		27	27		8.59
8	l <u>.</u>		1	1	8	1	8		ī			1		1	9		16	16		8.12
4			lī	i			2			1	1						6	6	42	7.00
5		l			1	1	1							1		1	5	5	48	9 80
B		1					 .	 									1	1	2	2.00
7				1			 .	 				1			••		2	2	16	8 00
0				•••		••••									1		1	1	15	15.00
lge not							١.													
stated	1	••••	••••	••••	••••		••••		• •	••		•••	$ \cdot\cdot $	•••	••	<u></u>	1	1	1	1.00
otals	624	558	414	846	802	178	146	00	58		14	7	5	5	7	1	2,780	2,808	10,111	3.64

^{*}One of these mothers bore twins.

[†] Two of these mothers bore twins.

The reference marks in this table show the number of the first child in each case of plurality-birth; thus, at the age of 28, four mothers bore twins; in two cases, they were the first and second children; in one case the fourth and fifth; and in one the fifth and sixth; at the age of 36 two mothers bore twins; in one case the fourth and fifth children, and in one the eleventh and twelfth.

There was one mother 50 years old who bore her fifteenth child in 1881. The number of mothers, of 40 years of age and over, who bore children in 1881 was 128, the number of children borne by them during the year, 129, and the whole number of children borne by them 972, an average of 7.59 children to each mother. The record of the ages of the mothers has been kept since July 1, 1863, a period of eighteen years and six months, during which time there have been 1,924 mothers, 40 years old and over, who have borne, during these years, 1,946 children. Including all the children previously borne by these same mothers, we have a total of 13,906 children, an average of 7.23 children to each mother.

In making up the statistics on this subject for 1881, a review was made of Table VII., by which several mistakes were discovered, chiefly in the report for 1873. They have been rectified, and the figures, as given in the present report, are believed to be correct.

During this time, since July 1, 1863, the number of children borne by mothers whose age was stated, was 39,524; the number of children to mothers of each age, during the time, was as follows:—

ge of mother. Children.	Age of mother.	Children.	Age of mother. C	hildren
14 5	27	2,366	40	905
15 19	28	2,778	41	253
16 57	29	2,017	42	315
17 165	80	8,157	43	178
18 413	81	1,849	44	108
19 685	32	1,884	45	111
201,107	88	1,578	46	80
211,282	84	1,329	47	26
231,906	85	1,989	48	12
232,111	-36	1,209	49	8
24 2,341	37	990	50	10
232,651	80	1,068	• -	
262,437	89	693	All known ages.3	9,524

These children were divided as follows to mothers of different divisions of age:—

Under twenty years	2,451
Twenty, and under twenty-five	7,637
Twenty-five, and under thirty	
Thirty, and under thirty-five	
Thirty-five, and under forty	
Forty, and under forty-five	
Forty-five and over	
Total	M3,62

Of these 39,524 children, 22,337 or 56.52 per cent., were borne by mothers less than thirty years old; 15,241, or 38.56 per cent., by mothers between thirty and forty years old; and 1,946, or 4.92 per cent., by mothers forty years old and over.

PLURALITY BIRTHS.

The number of cases of plurality births in Providence in 1881 23, which was 4 less than in 1880.

The following statistics are given in regard to them:

Cases. 23; number of children, 46.

Locality. First ward, 3; Second, 0; Third, 3; Fourth, 0; Fight, 0; Sixth, 4; Seventh, 3; Eighth, 3; Ninth, 2; Tenth, 5.

Season. January, 2; February, 1; March, 0; April, 4; May, 0; June, 3; July, 3; August, 1; September, 2; October, 3; November, 3; December, 1.

Sex. In seven cases both children were males; in five, both were females; in eleven cases, there was one of each sex; total, 25 male and 21 female children.

Color. The parents were white in every case.

Number of the Children and Age of the Mother. From the reference marks in Table VII. may be found the number of plurally births to mothers at different ages; and the number of the children

in cach case. In five cases they were the first and second children of mothers 24, 25, 27, and two of 28 years of age; in two cases, the second and third children of mothers 23 and 24 years af age; in two, the third and fourth children of mothers 26 and 33 years old; in four, the fourth and fifth children of mothers 28, 32, 36 and 40 years old; in seven, the fifth and sixth children of mothers 28, 29, 31, 32, 35, and two of 30 years old; in two, the sixth and seventh children of mothers 37 and 39 years old; in one case the eleventh and twelfth children of a mother 36 years old. There were eleven cases in which the mothers were under thirty years of age.

Parentage. In nine cases both parents were American; in five, both were Irish; in one, both were English; in one, both were Scotch; in one, both German; in one, the father was English and the mother American; in one, the father American and the mother Swedish; in one, the father American and the mother English; in one, the father Irish and the mother American; in one, the father British American and the mother Irish; in one, the father English and mother Irish.

TWENTY-SEVEN YEARS' SUMMARY OF PLURALITY BIRTHS.

In the twenty-seven years, 1855-1881, the number of cases of plurality births in Providence was 633; of twins, 624; of triplets, 9; total number of children, 1,275.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children and one female; making 9 male and 18 female children in the nine cases.

Of the 624 cases of twins, there were 204 in which both children were males; 211 in which both were females; and 209 in which there was one child of each sex; total, 617 male and 631 female children. Of the 1,275 children of all the cases of plurality births, there were 626 males and 649 females; a proportion of 103.6 females to 100 males, or 50.90 male and 49.10 female children in each 100.

Color. In all the cases of triplets the parents were white. Of the 624 cases of twins, in 602 the parents were white, and in 22 they were colored.

Parentage. Of the triplets, in one case, the parents were both American; in five cases, both were Irish; in one case, both German;

in one, the father was English and the mother Irish; and in one, the father Scotch and the mother Irish.

In the following table, for the 624 cases of twin births, the nativity of both parents is given; the first letter representing the father's birth-place, and the second the mother's.

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British American; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish.

Parentage. No.	Parentage. No.	Parentage. No.	Parentage. No.
A. and A 207	E. and A14	8w. and I 1	I. and W 1
J. and I240	S. and A 2	It. and I 1	A. and B. A
E. and E 30	B. A. and A 6	P. and I 1	I. and B. A 5
H. and 8 7	G. and A 2	A. and E 4	F. and B. A 1
G. and G 16	8w. and A 1	I. and E 6	8. and B. A 1
B. A. and B. A 10	A. and I	S. and E 1	A. and Sw 1
D. and D 1	E. and I12	G. and E 1	
P. and P 9	8. and I 3	A. and 8 1	Total624
it. and It 1	G. and I 1	I. and 8 3	
I. and A15	B. A. and I 2	E. and S 1	

There were 247 cases of twin births in which the mothers were of American birth; this was 39.58 per cent. of all; in 274 cases, or 43.91 per cent., the mothers were Irish.

In the 633 cases of plurality births in the twenty-seven years, there were 248 American mothers and 385 foreign mothers. The whole number of American mothers bearing children during this time was 24,924; the 248 mothers in cases of plurality births were about 1.00 per cent. of this number; the whole number of foreign mothers bearing children in the same time was 27,909,—the 385 foreign mothers in cases of plurality births were therefore 1.38 per cent. of all the foreign mothers.

Proportion of plurality births to whole number. The following statement shows how the year 1881 compares with the twenty-six years previous, as to the number of plurality births, and the proportion of these to the whole number of births; also the same facts for the whole period of twenty-seven years, 1855-1881 inclusive:—

Single Births. Twin. Triplet.	While Na. While Na. of Births. of Children
Single Births. 1 Will. 1 Tiplet. 26 years, 1855—188049,4206019	
1881 2,757 230	
27 years' totals52,1776249	52.>1053,455
Proportion of cases of plurality births:—	
Twins.	Triplets.
1855–1880 1 in 83 2	1 in 5,35%
1881 1 in 120.8	•••••
1855–1881 1 in 84.6	
Proportion of children of plurality births:-	_
Twins.	Triplets.
1855–18801 in 42.1	· · · · · · · · · · · · · · · · · · ·
18811 in 60.9	
1855-1881 1 in 42.8	l in 1,979.7
Proportion of plurality births by parentage.	
in the twenty-seven years, 1855-1881 inclusi	
following figures show the number of births	. •
and mixed parentage, and the number and cases in each class:—	proportion of plurality
Whole No. Births. Plura	lity Cases. 1 in Per Ct.
American parents21,201	
Foreign parents24,939	
Am. father, foreign mother 2,958	. 25 118.3 or 0.85
Foreign father, Am. mother 3,712	. 40 92.8 or 1.08

Of the 633 cases of plurality births, the 208 of American parentage were 32.86 per cent.; the 360 of foreign parentage were 56.87 per cent.; the 25 with American father and foreign mother were 3.95 per cent.; and the 40 with foreign father and American mother were 6.32 per cent.

TABLE IX. MARRIAGES, 1881. Nativity of the Parties.

	BIRTH-PLACES OF BRIDES.									
BIRTH-PLACES OF GROOMS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number of grooms		
United States	675	40	28		43	•••••	8	794		
Ireland	48	107	4		8	•••••		162		
England, Scotland and Wales	51	17	41		12			121		
Germany	8	1		10			1	20		
British America	25	4	2		87			68		
Portugal and Western Islands	1	1	• • • • • ·	·		8		10		
Other Countries	10	5		ļ. .			12	27		
Whole number of brides	818	175	75	10	95	8	21	1,202		

Bride and groom born in the same country. There were 890 marriages, or 74.04 per cent. of all, in which both parties were natives of the same country; this was 57 less marriages, and 2.83 per cent. less than in 1880. In 675 marriages, or 56.16 per cent. of all, both parties were born in the United States; this was 75 less marriages, and 4.72 per cent. less, than in the previous year. There were 107 marriages, or 8.90 per cent. of all, where both were natives of Ireland; 12 less marriages, and 0.75 per cent. less, than in 1880. There were 41 marriages where both persons were English or Scotch; 1 more than in 1880; 37 where both were British American; 17 more than in the preceding year; and 10 where both were German; 6 more than in the preceding year. The 10 brides born in Germany all married their own countrymen; also the 8 brides born in Portugal or the Western Islands.

Bride and groom born in different countries. The inter-marriage of the different nations represented in our population, is shown by the preceding table. Thus there were 312 marriages in which the bride and groom were of different nationalities; this was 25.96 per cent. of all the marriages; 27 more than in 1880, and 2.83 per cent. more. Of these 312 marriages, there were 119, or 38.14 per cent., in which the groom was born in the United States; and 143, or 45.83 per cent., in which the bride was born in the United States. Of the 794 grooms who were natives of this country, the 119 who married natives of other countries were 14.99 per cent.; this was 3.33 per cent. more than in 1880. Of the 162 grooms born in Ireland, 55. or 33.95 per cent., married women who were natives of other countries than their own; of the 121, born in England or Scotland, 80. or 66.12 per cent.; of the 20 born in Germany, 10, or just 50 per cent.; of the 68, born in British America, 31, or 45.59 per cent. There were 818 brides born in the United States; of these, 143, or 17.48 per cent., married natives of other countries; this was 1.75 per cent. more than in the previous year. Of the 175 brides born in Ireland, 68, or 38.86 per cent., married natives of other countries: of the 75 born in England or Scotland, 34, or 45.33 per cent.; of the 95 born in British America, 58, or 61.05 per cent.

We find in this table 27 grooms who were born in other countries than those named; they were natives of the following countries: Italy, 7; Sweden, 5; Norway, 4; Denmark, Greece and Switzerland, 2 each; Austria, Belgium, Holland, Africa, and the West Indies, one each.

The 21 brides who were natives of "other countries," were as follows: Sweden, 7; Italy, 4; Norway, 2; France, Hungary, Holland, Denmark, Bermuda, and Burmah, one each; and one was born in the Indian Ocean; and one "at sea."

In the following table are given the number and percentage of American, foreign and mixed marriages, in each of the twenty-seven years from 1855–1881, inclusive. By mixed marriages are meant those in which one of the parties was of American, and one of foreign nativity:—

	Amer	rican.	For	elgn.	Miz	red.
Year.	Number.	Per Cent.	Number.	Per Cent.	Number.	Per Cent.
1855	. 808	48.78	265	41.98	59	9.84
1856	852	54.74	212	82.97	79	12.29
1457	. 884	63.54	175	28.50	55	8.96
1858	. 288	50.54	314	88.21	63	11.25
1859	. 883	54 50	['] 224	36 66	54	8.84
880	336	54.19	925	36.29	59	9 52
1861	. 304	55.56	193	88.27	61	11 15
1863	. 248	52.65	179	82.72	80	14.63
1868	849	56.07	199	81 47	76	12.46
1864	. 427	57 78	211	28.55	101	13 66
L865	. 417	56.81	903	28.49	90	12.69
1886	. 471	58.00	244	80 05	97	11.95
1867	485	56.78	260	81 46	101	11.81
1868	580	00.78	215	21.66	127	14.56
1869	530	56.99	286	80.75	114	12.26
1870	552	57 56	262	27.82	145	15.13
1871	528	55.46	378	29.48	143	15.06
1872	591	54.52	805	28 14	188	17.84
1878	652	56.69	817	27.67	181	15.74
1874	648	55.10	846	29.42	182	15.48
1875	586	54.81	320	29.66	173	16.08
1876	547	57.64	241	25.39	161	16.97
1877	545	57.19	288	24.45	175	18.36
1878	601	59.15	245	24.12	170	16.73
1879	643	60.97	217	90.98	201	18.77
1880	750	60.88	248	19.72	239	19.40
1881	675	56.16	265	92.04	202	21.80

The proportion of American marriages was less in 1881 than it had been since 1875; the proportion of foreign marriages was larger than in any year since 1878; while the proportion of mixed marriages has steadily increased, and was larger in 1881 than ever before. The whole number of marriages for the twenty-seven years, as given in this table is 23,115; in Table VIII. it is 23,121; for an explanation of this discrepancy, see the Report for 1879, page 22.

Marriages of colored persons. The number of marriages among the colored people in 1881 was 61; 4 more than in 1880, and the

largest runder that has ever decurred in Providence in the year. The appropriate and average age of the parties in these 31 macroages were as follows:—

61 maies	म्हण, ं,जी प्रवासिक
& females	Min. 1. 17. Transfer

In the twenty-six years, 1506–1551 inclusive, the number of marriages of colored persons was 350; the appropriate and average age of the parties as fellows:—

At the January session of the General Assembly in 1881 the law forbidding the intermarriage of colored and white persons was repealed.

In consequence of this we find on the records five marriages where one of the parties was white and the other colored. In one instance a white man married a colored woman; it was his second marriage, and her third. In the other four cases, the man was colored and the woman white; in no case was it the first marriage of both parties; but in one case it was the first marriage of the man and the second of the woman; in one the first marriage of the woman, and the second of the man. In one instance, where it was the second marriage of both parties; both had been divorced.

Age of persons married. Table X. gives the number and percentage of all the brides and grooms in each division of ages:—

TABLE X. MARRIAGES, 1881. Ages of the parties.

	1	AGES OF THE BRIDES.											
AGES OF THE GROOMS.	Under 20 years.	20 to 25.	25 to 30.	30 to 35.	35 to 40.	40 to 45.	45 to 50.	50 to 55.	55 to 60.	60 to 65.	65 to 70.	Whole number of Groome.	Percentage
Under 20 years	12	5										17	1
20 to 25	93	233	45	5	1						in	377	1
25 to 30	40	179	135	25	8					1		288	2
80 to 35	11	58	65	40	13	3	1					191	10
35 to 40	2	21	28	19	10	5	2			****		87	7
10 to 45	3	8	12	12	12	6	3					.55	1
15 to 50	1	1	5	6	3	10	5		. 3	****	1	25	E
50 to 55		1	2	2	5	5	3	2		****		20	ı
5 to 60			3		3	2	1	1	1		****	П	
0 to 65			2			2	8	3	1			11	
5 to 70					2		1			1	9	-	
0 to 75							1					1	
75 to 80							1			****		1	
30 to 85						1					****	1	
Whole number of Brides	162	506	207	109	57	34	21	6	5	-2	3	1,207	
Percentage of Brides	13.48	42.10	24.71	9.07	4.74	2.83	1.75	.50	.41	.16	.25		10

There were 446 marriages, or 37.10 per cent. of all, in which the bride and groom were in the same division of age. The number of grooms who were less than 25 years old was 394, or 32.78 per cent. the number of brides 668, or 55.58 per cent.; 162 brides were less than 20 years old. One man more than 80 years old married woman between 40 and 45; and one man between 25 and 30, married a woman more than 60 years old.

The aggregate and average age of all persons married during the year were as follows:—

1.303 malesAggregate	age, 35,283 yearsAverage age, 29.35 years	
1,303 femalesAggregate	age, 30,801 yearsAverage age, 25.62 years	

The following figures give the same facts for each of the twenty-six years, 1856-1881 inclusive, and for the whole period:—

	Males	Females.
Years.	No. Aggregate age. Average age.	No. Aggregate age. Average age.
1856	639 18,104 years28.33 years	637 15,559 years24.42 years.
1857	607 16,753 years27.59 years	606 14,427 years23.84 years.
1856	560 15,850 years28 29 years	560 13,702 years 24.47 years.
18 59	611 16,989 years27.61 years	611 14,616 years 23.92 years.
1880	620 17,135 years27.50 years	620 14,667 years23.66 years.
1861	547 15,448 years28.24 years	547 13,316 years22.33 years.
1962	547 15,773 years28.83 years	547 13,565 years24.80 years.
1863	610 17,208 years28.21 years	610 14,792 years24.24 years.
1864	739 21,361 years28.90 years	739 18,075 years24.45 years.
1865	709 20,539 years28.96 years	709 17,605 years24.83 years.
1866	812 22,801 years28.08 years	812 20,036 years24.67 years.
1867	855 24,443 years28.58 years	855 21,078 years24.65 years.
1868	872 24,546 years28.15 years	872 21,440 years24.59 years.
1869	990 96,548 years28.55 years	930 23,135 years24 88 years.
1370	959 27,343 years26.51 years	959 23,773 years24.79 years.
1871	943 26,743 years28.37 years	943 23,290 years24.70 years.
1879	1,084 31,110 years28.70 years	1,084 26,835 years24.76 years.
1873	1,150 \$2,879 years28.59 years	1,150 28,701 years24.96 years.
1874	1,176 33,927 years28.85 years	1,176 29,510 years25.09 years.
1875	1,079 30,848 years28.59 years	1,079 27,090 years25.11 years.
1876	949 27,496 years28.97 years	949 23,640 years 24.91 years.
1877	958 27,904 years29.28 years	953 24,092 years25.28 years.
1878	1,016 29,460 years28.99 years	1,016 25,796 years25.40 years.
1879	1,071 \$1,016 years28.96 years	1,071 27,103 years25.32 years.
1880	1,232 36,127 years29.32 years	1,232 31,015 years25.18 years.
1881	1,202 35,283 years29.35 years	1,202 30,801 years25.62 years.
% years.	22,472643,634 years28.64 years2	2,460557,659 years24.82 years.

The number of males and females does not agree in this table, because in 1856 the ages of 17 males and 19 females, and in 1857, the ages of 7 males and 8 females, were not given.

Number of the marriage. - Table XI. shows how many of the mar-

riages in 1881 were the first, second, third, etc., marriages of the parties:—

TABLE XI. MARRIAGES, 1881. Number of times married.

GROOMS.	BRIDES. Number of the Marriage.			Whole number of
Number of the Marriage.	ringe. First.	Second.	Third.	Grooms.
First	878	86	4	968
Second	138	68	5	311
Third	11	10	2	23
Whole number of Brides	1,027	164	11	1,202

For 1,027 of the brides, or 85.44 per cent. of all, it was the first marriage; also for 968 of the grooms, or 80.53 per cent. There were 878 cases, or 73.04 per cent., in which it was the first marriage of both parties; this was 46 less than in 1880, and 1.96 per cent. less. In 68 cases, or 5.66 per cent. of all, it was the second marriage of both.

Denominational. The 1,202 marriages of 1881 were solemnized by the clergymen of different denominations, or by civil authority, as follows:—

Roman Catholic387	Advent Christian	11
Calvinistic Baptist142	Evangelical Advent	11
Free Baptist 43	Latter Day Saints	1
Methodist191	Hebrew	2
Episcopalian154	Bethel Pastor	1
Congregationalist119	Lutheran	1
Unitarian 81	New Jerusalem	3
Universalist45	Free Religious	1
Presbyterian32	Denomination not specified	13
Christian 8	Justice of Supreme Court	1
Seamen's Mission	TotalI.	202

Nativity and residence. In 103 marriages, or 8.57 per cent. of all, both parties were born in the city of Providence; in 112 others, or 9.32 per cent., the groom was born in this city; and in 205, or 17.05 per cent., the bride only was born here; this makes a total of 523 persons, or 21.76 per cent. of all the persons married, who were natives of the city.

There were 195 marriages, or 16.22 per cent. of all, in which both parties were non-resident, 121 in the first six months of the year, and 74 in the second six months; of these 195 marriages there were 130, or 66.66 per cent., in which both parties came from outside of the State; and in 102 of these, both parties were residents of Massachusetts; in 13 others, one of the parties was from that State.

Marriage and education. The number of persons married in Providence in 1881 was 2,404, divided as follows as to sex and nativity:—

Born in United States	. Males, 794	Females, 818	Total, 1,612
Born in Foreign countries	.Males, 408	Females, 384	Total. 792

Of these, the number and proportion, who signed the marriage certificate with a mark, were as follows:—

1881.	Whole Number Married.	Signed with a Mark.	In Each 100.
Males, born in United States	791	83	4.16
Females, born in United States	818	89	4.77
Total, born in United States	1,612	72	4.47
Males, born in for ign countries	408	74	18.14
Females, born in foreign countries	384	97	25.26
Total, born in foreign countries	793	171	21.59
Totals	2,404	243	10.11

Of the 72 persons born in the United States, who signed the marriage certificate with a mark, 34 (16 males and 18 females) were colored persons; and 27 (10 males and 17 females) were children of

foreign parents; so that the number of persons of white American parentage, who signed with a mark, was only 11 (7 males and 4 females); this was 2 more than in 1880; 4 more males and 2 less females.

The number of persons who signed with a mark in 1881 was 27 less than in 1880; and the proportion of these to the whole number of persons married was 0.85 less. The decrease was entirely in the number of females; the number of males who used the mark, being much larger than in the preceding year. The decrease was very evenly divided between the American and foreign, as to numbers; but the proportion of the foreign class who signed with a mark decreased very much more than that of the Americans. The following are the figures: Of those born in the United States, the number was 14 less in 1881 than in 1880; males, 9 more; females, 23 less; the proportion, 0.48 less. Of those born in foreign countries, the number was 13 less in 1881 than in 1880; males, 10 more; females, 23 less; the proportion, 3.79 less.

The statistics on this point for the twenty-two years, 1859-1880, are as follows:—

	Whole Number	Signed with	In Each
1859-1880.	Married.		100.
1899-1880.	Married.	a Mark.	1004
Males, born in United States	12,443	585.,	4.70
Females, born in United States	12,696	856	
Total, born in United States	25,139	1,451	5.71
Males, born in foreign countries.	7,021	2,223	31.68
Females, born in foreign countrie	8 6,768	3,225	
Total, born in foreign countries	es13,789	5,448	
Totals	38,928	6,889	17.70

DEATHS.

There were 2,145 deaths in Providence in the year 1881. This was 65 more than in the preceding year; males, 52 more; females, 13 more; whites, 54 more; colored, 11 more; natives of the United States, 41 more; of foreign countries, 24 more; of American parentage, 79 more; of foreign parentage, 14 less.

The total number of deaths in the twenty-seven years, 1855–1881 inclusive, was 37,715; the average annual death rate, one in 50.73 of the population. In 1881 the death rate was one in 50.11, or 19.95 deaths in each thousand inhabitants. In the different classes of the population the rate was as follows:

	Estimated Population, 1881.	Deaths, 1881.	To population one death in	Deaths to each 1,000 of population.
Whole number	107,500	2,145	50.11	19.90
Males	51,041	1,047	48.75	20.51
Females	56,459	1,098	51.41	19.48
White	103,763	2,038	50.91	19.64
Colored	8,737	107	34.93	28.68
Born in United States	78,819	1,622	48.59	20.58
Born in Foreign countries	28,681	523	54.84	18.24
American parentage	49,525	999	49.57	20.17
Foreign parentage	57.975	1.146	50 50	10 77

The following general statistics are given in regard to the 2,145 deaths:—

Table XII. Deaths, 1881.

Sex, condition, color, nativity, parentage, locality, season.

													-
1881.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Whole Year.
Whole number of Deaths	190	164	173	200	195	162	196	224	160	162	153	166	2,165
Sex.—										0			1774
Males	85	82	111000		10000			122	72	89		81	1,00
Females	105	82	89	111	95	89	99	102	88	73	80	85	1,08
Condition.—													-
Married	78	1000	58		71	57	43					57	100
Single	86	S2 22	86		96	87	137			96		10	1,181
Widows	17	6	21	10	18	15	14	13	15	17	35	2 4	15
Divorced		0		9	1	1		1		1	1	-	16
	***						***			1			1
Color.—		754	101	100	200	***	105	ave	401	400	740	201	-
Whites	177	1000	1000	7.0	181	10	185	11	154	158	145	164	1,008
	13	10	9	10	19	10	**	111	0	i i			201
Nativity.—		100				200	***	***	100	200		***	
United States	131	126		10000	1000	1000	159	173	120	133	2000	119	1,5
Ireland	38	25	28	41	43	20	25 6	11	6	3	29	20	101
England and Scotland Germany	17	7	7	10	11	9	5	1	3	3	-	1	16
British America		4	4	2	2	3		2	2		1	7	21
Portugal & West'n Islands.	1	1]]]				1
Other countries	1	1		1	1						1	1	- 35
Parentage.—								m					
American	93	86	88	96	94	71	81	102	59	76	72	81	-900
Irish	63	51	62	75	72	59	78	83	72	60	57	60	755
English and Scotch	25	15	14	19	20	18	22	21	11	15	19	17	20
German	4	2	2	3	3	5	3	5	5	1	3	1	15
British American		7	4	6	3	5	8	10	5	7	5	5	65
Portuguese	2	1	2		1	1	1	1	4	1	2		16
Other foreign	3	2	1	1	2	3	3	2	4	2	2	2	27
Locality, Wards				1									
I	31	24	24	23	20	23	24	29	19	30	19	16	122
H	12	13	5	7	7	7	10	11	4	4	10	8	25
III	13	14	18	18	17	16	19	20	11	24	15	13	198
IV	14	14	7	5	7	14	6	15	15	8	9	*	120
V	18	12	16	12	13	13	20	23	12	13	17	21	190
VI	5	9	13	9	13	11	11	11	8	4	11	9	114
VII	16	10	21	18	23	15	21	15	17	7	11	11	185
VIII	27	12	20	25	16	17	15	20	19	12	9	23	215
IX	17 28	26	13	32	26	19	26	28	17	18	17	95	34
Y Public Institutions	9	7	9	14	16	6	8	11	8	33	25	20 8	115
Totals	190	164	173	200	195	162	196	224	160	162	153	100	2,16

Season. The largest number of deaths in 1881 was in the third quarter of the year, and the smallest in the fourth quarter. The number of deaths in each quarter of the year 1881, and of each of the five years preceding, is given in the following figures; also the quarterly aggregate and average numbers for the twenty-six years, 1855–1880 inclusive:—

2	1881.	1880.	1879.	1878.	1877.	26 years, 1855-1 1876. Aggregate. Av'	
January-March	527	560	514	541	411	383 8,593	330
April-June	557	533	439	423	440	419 7,881	303
July-September	580	525	493	520	557	57310,371	390
October-December	481	462	580	505	530	490 8,725	336

The number and percentage of deaths in each quarter of the year 1881, and for the period of twenty-six years, 1855-1880, were as follows:—

	188	1.	26 years, 1855-1880.		
D	eaths.	Per cent.	Deaths.	Per cent	
January-March	527	24.57	8,593	24.10	
April-June	557	25.97	7.881	22.12	
July-September	580	27.0t	10,371	29.10	
October-December	481	22.42	8,725	24.58	
-		_		_	
Totals	2.145	100.00	35 570		

The following figures show the number and percentage for each quarter of each of the twenty-seven years, 1855-1881 inclusive:—

				_	_	_	
	Januar	y-March.	Apri	l-June.	July	-Sept.	OctDec.
	Deaths.	Per cent.	Deaths.	Per cent.	Deaths.	Per cent.	Deaths. Peret.
1855	213	21.49	208	20.99	339	34.21	231
1856	308	28.92	244	22.91	296	27.80	21720.57
1857	214	23.14	173	18.70	286	30.92	252
1858	279	27.43	278	27.34	257	25.27	20319.96
1859	213	23.69	205	22,80	270	30.04	21122.67
1860	243	24.27	209	20.88	297	29.67	25225.18
1861	240	22.84	236	22.45	336	31.97	23921
1862	197	21.55	192	21.01	271	29 65	25427.77
1863	287	23.64	248	20.43	370	31.14	301 24.79
1864	302	23 58	273	21.31	403	31.46	30321.65
1865	299	24.69	240	19.82	372	30.72	300 \$4.77
1866	238	22.97	218	21.04	338	32.63	243 25.35
1867	258	26.88	202	21.04	286	29.79	21422.29
1868	231	20,81	244	21.98	346	31.17	28926.04
1869	331	26.35	250	19.90	363	28.91	31224.84
1870	331	26 20	271	21.46	371	29.37	290
1871	311	24.80	263	20.98	385	30.70	29591#
1872	330	20,59	304	22.71	518	32.31	3912130
1872	427	24.84	407	23.68	504	29.32	
1874	444	22.35	501	25 21	556	27 98	486 24.46
1875	488	25.48	401	20.94	531	27.73	495
1876	383	20.54	419	22.47	573	30.72	49026.27
1877	411	21.21	440	22.70	557	28.74	53027.25
1878	541	27.20	423	21.27	520	26.14	505
1879	514	25.37	439	21.67	493	21.33	58028.63
1880	560	26.92	533	25.63	525	25.24	462 22.21
1881	527	24.57	557	25.97	580	27.04	48122.0

In the first quarter of the year 1881 the number of deaths is smaller than in the previous year, and the percentage less than it has been since 1877. In the second quarter the number and percentage are larger than ever before at this season; though there was no epidemic of any particular disease, a large number of deaths occurred in these three months. In the third quarter the number is larger than ever before, but the percentage, though larger than for four years previous, is smaller than in many of the earlier years. In the fourth quarter the number and percentage are slightly larger than in 1880, but the percentage is less than for some years previous to that.

Sex and Parentage. The following statistics are given as to the sex and parentage of the 2,145 decedents in 1881:—

American parentage	473 males	526 femalestotal, 999
Foreign parentage	574 males	572 femalestotal, 1,146
Totals1	.047 males	1.098 femalestotal, 2,145

There were 51 more females than males; of American parentage, 53 more; of foreign parentage, 2 less. The number of foreign parentage was 147 more than that of American parentage; of males, 101 more; of females, 46 more. In the twenty-seven years, 1855–1881, the decedents were divided, as to sex, as follows:—

Condition. The number of married persons who died in 1881 was 669; 70 more than in 1880; 319 of American, and 350 of foreign parentage. The number of single persons who died was 1,181; 9 less than in 1880; 516 of American, 665 of foreign parentage. The 210 widows who died were 116 of American, and 94 of foreign parentage; the whole number being 22 more than in 1880. The number of widowers, 75, was 24 less than in the preceding year; 39 were of American, and 36 of foreign parentage. The number of divorced persons who died in 1881 was 10; 9 of American, and one of foreign parentage; 6 more than in 1880.

Color. The number of white decedents in 1881 was 2,038; 54 more than in 1880; the number of colored, 107, or 11 more than in 1880. The number of white children born in 1881 was 2,682; the proportion of births to deaths, therefore, was 100 to 75.99. The number of colored children born was 121; the proportion of births to deaths, 100 to 88.43. In the twenty-seven years, 1855–1881, the number of colored children born was 2,059; the number of colored decedents in the same time, 2,156; the proportion of births to deaths, 100 to 104.7.

Nativity and Parentage. The decedents are always registered according to nativity, and also according to parentage; and the great difference in the two classes of statistics shows the value of such a

distinction. The number of decedents of American nativity in 1881 was 1,622, or 75.62 per cent. of all; the number of American nativity was 41 more than in 1880, and the percentage 0.39 less the number of American parentage was 79 more than in the previous year; the percentage 2.34 more.

The number of decedents of foreign birth in 1881 was 523, w 24.38 per cent. of all; this was 24 more than in 1880, and 0.39 per cent. more. The number of foreign parentage was 1,146, or 53.45 per cent. of all; 14 less and 2.34 per cent. less than in the preceding year.

There were 10 decedents who were natives of "other countries" than those named in the table; they were as follows: native of Sweden, 4; of West Indies, 2; of France, Wales, Italy and Fas Indies, 1 each. The 27 of "other foreign" parentage were: Swedish, 2; Italian, 6; Polish, 3; French, 2; Russian, Austrian, Dutch, Swiss, African, East Indian, and West Indian, 1 each.

Locality. The number of deaths in each ward of the city and in Public Institutions is given in Table XII., for each month and for the whole year. The number of deaths in the first ward in 1881 was 29 more than in 1880; in the second ward, 33 more; in the third, 15 more; in the fourth, 32 more; in the fifth, 28 more; in the sixth 16 more; in the seventh, 2 less; in the eighth, 33 less; in the ninth, 17 less; in the tenth, 71 less; in Public Institutions, 35 more. The first six wards and Public Institutions show an increase of 188 over the preceding year; the remaining four wards, a decrease of 123.

The number of decedents in each ward, according to parentage was as follows:—

Wards.	American,	Foreign.	Wards.		-	Fores
I	119	163	VI		79	
11	81	17	VII		90	mar.
III	81	117			126	
IV	59	63			145	
V	82	108	X		79	
			Public !	Instit	ntions 58	evenin.
Who	le city					

The following figures show the number of deaths in different Pullic Institutions in 1881, and in each of the two years previous:-

	1881.	1:360.	1879
Rhode Island Hospital	53	#	3
Butler Hospital	18	8	t
Dexter Asylum			
Home for Aged Men	•	1	
Bome for Aged Women			
Children's Home			
Children's Hospital and Nursery			
R. C. Orphan Asylum			
R. C. Convent			
Reform School	1		
Little Sisters' Home for the Poor	1		
Friends' School	6		
Shelter for Colored Children	3		(
Hotels	4		
Police Station			
Total	115		7

Aggregate and average age. Table XIII. shows the number of deaths and the aggregate and average age of the decedents, according to parentage, for each month and for the whole year 1881:—

TABLE XIII. DEATHS, 1881.

Aggregate and average age, by parentage.

	Amer	ican Par	entage.	Fore	ign Pare	ntage.	Total American and Foreign.			
MONTHS.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	
January	93	3,562	38.30	97	3,345	84.50	190	6,907	36.3	
February	86	3,072	35.72	78	2,319	29.73	164	5,391	32 .8	
March	88	3,315	37.67	85	2,748	32.33	173	6,063	35.0	
April	96	3,460	36.04	104	3,249	31.22	200	6,709	83.5	
Мау	94	2,815	29.95	101	8,824	82.91	195	6,139	31.4	
Jane	ก	2 956	41.63	91	2,414	26.52	162	5,370	33 .1	
July	81	1,649	20.35	115	2,328	20.24	196	8,977	20.9	
August	102	2,492	24.47	122	3,033	24.86	924	5,595	24.6	
September	59	1,743	29.54	101	2,345	23.22	160	4,088	25.8	
October	78	2,614	84.39	86	1,908	22.19	182	4,522	27.9	
November	72	2,840	89.44	81	2,966	27.97	158	5.106	38.8	
December	81	8,608	44.54	85	2,903	84.15	166	6,511	89.9	
Whole Year	999	34,126	34.16	1,146	32,182	28.08	2,145	16,808	80.9	

The average age of all decedents in 1881 was 30.91 years; 1.34 years more than in 1880. Of the decedents of American parentage, the average age in 1881 was 34.16 years; 0.38 years less than in 1880; of those of foreign parentage, the average age was 28.08 years, or 2.45 years more than in the preceding year.

Comparing this table with the same for the preceding year, we find the following differences in the different months. Among the decedents of American parentage, there were six months in 1881 when the average age was greater than in 1880; namely: January, February, June, September, November, and December; in the remaining six months, the average age was less in 1881 than in the year before Among the decedents of foreign parentage, there were eight months in 1881 when the average age was greater than in 1880, January, February, March, April, May, July, August and November; in the other four months the average age of the decedents was less than in 1880.

Table XIV. gives the same statistics for the twenty-six years, 1856-1881, inclusive:—

Table XIV. Deaths, Twenty-Six Years, 1856-1881.

Recapitulation. Aggregate and average age by parentage.

TWENTY-SIX	Amer	ican Pare	entage.	Fore	ign Parer	ntage.	Total	American Foreign.	and i
YEARS. 1856-1881.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average
January	1,618	55,579	34.35	1,479	38,249	25.86	3,097	58,89	23
February	1.470	49,768	33.86	1,383	32,117	23.95	2,853	81,887	27
March	1,536	54,224	35.30	1,421	35,764	25.17	2,957	89,988	
April	1,500	51,379	84.25	1,498	40,120	26.78	2,998	91,499	
May	1,427	51,368	36.00	1,355	35,404	26.13	2,782	88,77	
June	1,178	41,011	34.81	1,272	31,855	25.04	2,450	72,500	
July	1,620	41,575	25.66	1,952	36,402	18.65	3,572	77,977	21.5
August	1,749	48,022	27.46	2,053	39,138	19.06	3,802	87,100	223
September	1,516	44,283	29.21	1,722	34,256	19.89	3,938	78,50	24.5
October	1,403	46,753	33.32	1,567	36,079	23.02	2,970	82,833	27.5
November	1,429	47,955	33.56	1,503	37,517	24.96	2,932	85,479	21
December	1,570	56,577	36.04	1,503	39,798	26.48	3,073	96,873	31.1
Totals	18,016	588,494	32.67	18,708	436,699	23.34	36,794	1,025,193	27.3

Both of these tables show great differences between the average age of the two classes. In the year 1881, (Table XIII.,) the highest average age of American decedents was 44.54 years in December; while the highest average age of the foreign class was 34.50 years in January; a difference of more than 10 years; the lowest average age of both classes was in July; for those of American parentage, 20.35 years; of foreign parentage, 20.24 years. The greatest difference between the two classes was 15.11 years in June; the least difference, 0.11 of one year in July. In the months of May and August the average age of the decedents of foreign parentage exceeded that of the American decedents, which very seldom happens.

In Table XIV. we find the highest average age of American decedents in the period of twenty-six years, 1856-1881 inclusive, was 36.04 years in December; of those of foreign parentage, the highest average age was 26.78 years in April; 9.26 years less than the highest average of the American class. Of both classes of decedents, the lowest average age was in July; of Americans, 25.66 years; of foreign decedents, 18.65 years. The average age of American decedents was less in the year 1881 than in the long period of years, only in the months of May, July, and August; in the other nine months it was greater. Of the decedents of foreign parentage, the average age was less in the single year than in the term of years, only in the month of October. For the year, the average age of American decedents was 1.49 years greater in 1881 than in the period of twenty-six years; of the decedents of foreign parentage, the average age for the year in 1881 was 4.74 years more than for the period of years.

The following table shows the difference between the two classes in each month, both for the single year 1881 and for the twenty-six years, 1856—1881. As already stated, in the months of May and August, 1881, the average age of American decedents was less than that of the foreign class; this is indicated by prefixing the sign — in the column of "difference" to the figures for those months. In the other months of 1881, and in every month in the long period of years, the average age of the American decedents was greater than that of the foreign.

Difference in Years between the Average Age of American and Foreign

Decedents.

		1881.		26 3	rears, 1866-1	881.
MONTHS.	Averag	e Age.		Averag	e Age.	
	American.	Foreign.	Difference.	American.	Foreign.	Difference
January	38.30	84.50	8.80	84.85	25.86	8 49
February	85.72	29.73	5.99	83.86	23.95	9.91
March	87.67	32.83	5.84	85.30	25.17	10.13
April	86.04	81.22	4.82	84.25	26.78	7.47
May	29.95	82.91	2.96	86 00	26.13	9.87
June	41.63	26.52	15.11	84.81	25.04	9.77
July	20.35	20.24	0.11	25.06	18.65	7.01
August	24.47	24.86	0.39	27.46	19.06	8.40
September	29.54	23.22	6 32	29.21	19.89	9.32
October	84.89	22.19	12.20	83 32	23.02	10.30
November	89.44	27.97	11.47	88.56	24.96	8.60
December	44.54	84.15	10 89	86.04	26.48	9.56
Whole Year	84.16	28.08	6.08	82.67	23.34	9.83

In June, October, November and December, the difference between the two classes is greater in the single year than in the term of years; in the other months it is the reverse.

In the year 1881 the largest number of deaths in one month among those of American parentage was 102, in August; the smallest number, 59, in September. Of decedents of foreign parentage, the largest monthly total was 122, in August; the smallest, 78, in February. In the period of twenty-six years, 1856–1881, the largest monthly aggregate for both classes was in August; American, 1,749; foreign, 2,053; the smallest monthly aggregate for both classes, in June; American, 1,178; foreign, 1,272. The percentages of all the deaths in the different quarters of the year, according to parentage, for the period of twenty-six years, 1856–1881, were as follows:—

	1st Quarter.	2d Quarter.	3d Quarter. 4th Quarter.
American parentage	95.67	22.79	27.1124.43
Foreign parentage	22.89	22.05	30.6124.45

Table XV. gives the number of decedents, and the aggregate and average age, according to parentage, in each of the twenty-six years, 1856–1881, inclusive, and for the whole period:—

TABLE XV. DEATHS, 1856-1881.

Recapitulation. Aggregate and average age by parentage.

:	Amer	ican Par	entage.	Fore	eign Pare	ntage.	Tota	American Foreign.	and
YEARS.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregate Years.	Average Years.
1856	541	14,554	26.90	524	8,258	15.75	1,065	22,812	21.4
1857	523	16,314	81.19	402	8,007	19.91	925	24,321	26.2
18 58	512	15,910	81.07	505	8.478	16.78	1,017	24,888	23.9
859	479	14,993	81.30	420	8,640	20.57	899	23,6 33	26.2
860	512	16,640	32.54	489	9,751	19 94	1,001	26,411	26.3
861	538	16,327	30.34	513	9,605	18.72	1,051	25,932	24.6
1862	460	14.589	31.71	454	10,994	24.21	914	25,583	28.0
1863	597	19,398	82.49	617	13,757	22.29	1,214	33,155	27.3
864	649	21,732	33.48	632	11,365	17.98	1,281	88,097	25.8
865	681	19,628	3 1.10	580	11,991	20.67	1,211	31,619	26.1
86	513	18,906	36 .85	523	12,762	24.40	1,036	81,668	80.5
867	528	19,258	8 6.47	432	12,149	28.12	960	81,407	82.7
868	596	21,779	36.54	514	12,539	24.89	1,110	34,318	30.9
869	627	20,490	82.68	629	14,776	23.49	1.256	85,266	28.0
870	642	21,078	32.83	621	14,393	23.18	1,263	85,471	28.0
871	643	22,725	85.34	611	17.005	27.83	1,234	89,73 0	81.6
873	832	27,572	88.14	771	19,160	21.88	1.603	46,782	29.1
873	862	28,675	83.27	857	19,061	22 24	1,719	47,736	27.7
874	885	27,588	81.17	1,102	24,802	22.51	1,987	52,390	26.8
875	. 914	3 0,419	33 2 8	1,001	24,802	24 28	1,915	54,721	28.5
876	875	29,025	83.17	990	23,959	24.20	1,805	52,984	28.4
877	908	28,353	81.22	1,030	25,410	24.67	1,938	53,768	27.7
878	889	26,724	· 8 0.06	1,100	25,178	22.89	1,989	51,902	26.0
879	941	29,896	81.77	1,085	28,44 6	26.22	2,026	58,342	28.8
880	920	31,775	84.54	1,160	29,729	25.63	2,080	61,504	29.5
861	999	34,126	34.16	1,146	82,182	28.08	2,145	66,808	80.9
8 years' totals.	18,016	588,494	82.67	18.708	436,099	23.54	36,724	1,025,193	27.9

The greatest difference between the average age of the two classes of decedents was 15.50 years in 1864; the least difference, 5.55 years, in 1879. The difference between them in the whole period was 9.83 years.

Number and percentage of decedents at different ages. Table XVI. gives the number and percentage, according to parentage, of decedents in different divisions of ages, in 1881; also for the twenty-six years, 1856–1881 inclusive:—

TABLE XVI. DEATHS, 1881.

Percentages at different ages, according to parentage.

		rican itage,		eign stage, 31.	Total / can an eign,		26 ye 1856-	
AGES.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Under 1 year	206	20.62	2.9	18.24	415	19.35	7,124	19.4
1 and under 2	51	5.10	95	8.29	146	6.81	8,183	8.67
2 and under 5	81	8.11	83	7.42	166	7.74	8,427	9.3
Total under 5	83×	33.83	889	88.95	727	88.90	13,784	87.40
5 and under 10	41	4.40	47	4.10	91	4.24	1,943	5.18
10 and under 15	14	1.40	22	1.93	86	1 68	749	2.04
15 and under 20	25	2.50	88	8.32	6:3	2.94	1,902	3.27
20 an d under 30	67	6.71	170	14.84	237	11.05	3,716	10.1:
80 and under 40	85	8.51	108	9.42	193	9.00	3,312	9,0
40 and under 50	67	6.71	91	7.94	158	7.86	2,784	7.56
50 and under 60	95	9.51	91	7.04	186	8.67	2,067	7.26
80 and under 70	87	8.71	88	7.68	175	8.16	2,621)	7.16
70 and under 80	95	9.51	61	5.82	156	7.27	2.473	6.73
60 and under 90	71	7.11	85	3.05	106	4.94	1.292	8.5
90 and over	11	1.10	6	.52	17	.79	2/13	.73
Total	994	100.00	1,146	100.00	2,145	100.00	86,724	100.00

The percentage of deaths of children of American parentage, under five years of age, was a little larger in 1881 than in 1880; the per-

centage of decedents between fifteen and twenty years of age was very much larger than in the previous year. This increase of deaths among younger persons reduced slightly the average age of all decedents of American parentage, as was shown by Table XIII. Of the decedents of foreign parentage, the percentage under five years old was 2.26 less than in 1880; also the percentages of decedents between five and twenty years of age were 4.29 less than in the previous year; while the percentage of foreign decedents in 1881, over 50 years of age, was 2.62 more than in 1880; this caused the average age of all decedents of foreign parentage to be 2.45 years more in 1881 than in 1880.

The following statement gives the percentage, according to parentage, of the decedents under five years old, in each of the twenty-six years, 1856-1881, inclusive:—

American.	Foreign. Total American & Foreign
Under five years.	
•	54.95 per cent45.91 per cent
185732.31 per cent	51.74 per cent40.75 per cent
185835.35 per cent	55.44 per cent45.33 per cent
185928.60 per cent	48.34 per cent37.82 per cent
186032.81 per cent	48.87 per cent40 66 per cent
186135.87 per cent	52.63 per cent 44.05 per cent
1862 31.08 per cent	42.07 per cent 36.54 per cent
1963 28.47 per cent	46.84 per cent37.81 per cent
1864 28.81 per cent	51.11 per cent 39.82 per cen
186529.95 per cent	46.37 per cent37.82 per cent
188623.78 per cent	40.53 per cent32.24 per cent
186724.05 per cent	36.35 per cent29.59 per cent
196827.35 per cent	43.19 per cent34.69 per cent
186930.78 per cent	43.24 per cent37.02 per cent
1870 31.00 per cent	42.67 per cent36.72 per cent
187128.63 per cent	36.01 per cent32.22 per cent
30.89 per cent	40.00 per cent35.27 per cent
•	43.05 per cent36.71 per cent
•	43.55 per cent39.31 per cent
	40.96 per cent36.66 per cent
•	43.13 per cent38.98 per cent
•	41.26 per cent87.36 per cent
•	42.64 per cent40.02 per cent
-	37.24 per cent
•	• 2
-	36.21 per cent
1881 33.83 per cent	33.95 per cent33.90 per cent

In the forty-two years, 1840-1881 inclusive, there were 49,869 decedents in Providence, of whom 19,469 were less than five years old; the number and percentage in each division of age under five years were as follows:—

Whole number of decedents		49,8	969			
Under one year	. 9,629	or	19.31	per c	ent.	of all.
One and under two years	. 4,966	or	9.96	per c	ent.	of all.
Two and under five years	4,874	or	9.77	per c	ent.	of all.
Total under five years	.19,469	or	89.04	per c	ent.	of all.

Table XVII. shows the number and percentage of decedents in different divisions of age, according to parentage, in the aggregate for the twenty-six years, 1856-1881 inclusive:—

TABLE XVII. DEATHS, 1856-1881.

Number and percentage at different ages, according to parentage.

		Parentage 1856-1881.		arentage, 1856–1881.	American Parentage 1856	& Foreign , 26 years, -1881.
AGES.	Number of Deaths.	Percentage in each division of age.	Number of Deaths.	Percentage in each division of age.	Number of Deaths.	Percentage in each division of age.
Under 1 year	8,159	17.58	8,965	21.19	7,124	19.40
1 and under 9	1,169	8.49	2,014	10.77	8,183	8.67
2 and under 5	1,896	7.75	9,081	16.86	3,427	9.33
Total under 5	5,794	81.77	8,010	42.82	18,784	87.40
5 and under 10	981	5.17	972	5.20	1,903	5.16
10 and under 15	891	2.17	858	1.91	749	2.04
15 and under 20	609	3.38	598	3.17	1,202	8.27
20 and under 30	1,632	2.06	2,084	11.14	3,716	10.12
30 and under 40	1,559	8.65	1,758	9.87	8,819	9.02
40 and under 50	1,859	7.54	1,425	7.62	2,784	7.58
50 and under 60	1,466	8.14	1,201	6.49	2,667	7.26
60 and under 70	1,565	8.63	1,074	5.74	2,629	• 7.16
70 and under 80	1,655	9.19	818	4.87	2,478	6.78
80 and under 90	964	5.85	328	1.75	1,292	3.53
90 and over	171	.95	92	.49	268	.73
Totals	18,016	100.00	18,708	100.00	86,794	100.00

The percentage of foreign decedents under five years old in 1881, (Table XVI.,) was the smallest ever recorded, and was only 0.12 per cent. larger than the percentage of American decedents of that age, which was quite unusual. We may see from Table XVII. that generally the proportion of the foreign far exceeds that of the American; in the twenty-six years, 1856-1881, the number of decedents of foreign parentage under five years old was 2,286 more than that of American parentage; and the percentage 11.05 more. the age of five years, the percentages of American decedents increase with advancing age; the largest percentage being 9.19, between 70 and 80 years of age; while among the decedents of foreign parentage, the percentage between 20 and 30 years is much the largest; the proportion steadily decreasing after that age. The number of American decedents in the twenty-six years who lived beyond the age of 50 years was 5,811; the percentage, 32.26; of foreign decedents, the number was 3,513; the percentage, 18.77. These facts accord with the statements under Tables XIII. and XIV. that the average age of American decedents is greater than that of the foreign class; in the twenty-six years, 1856-1881, it was 9.33 years more.

From Tables XVI. and XVII. we find that the percentage of children of American parentage under five years old, who died in 1881, was 2.06 more than the proportion of that class in the twenty-six years, 1856–1881; the proportion of decedents of 60 years and over in the single year was 2.31 per cent. more than in the term of years. Of foreign decedents, the proportion under five years of age in 1881 was unusually small, and was 8.87 per cent. less than in the period of twenty-six years; the proportion in 1881 who lived beyond the age of 60 years was 4.22 per cent. more than in the long number of years.

Deaths by age and parentage in each month and quarter of the year. The comparative mortality of the two classes of decedents at different ages and seasons is shown in Table XVIII. It gives the number of deaths at each age and at all ages, according to parentage, in each month and quarter of the year 1881. The whole number of deaths during the year was 2,145; of American parentage, 999; of foreign parentage, 1,146; divided as follows in the different seasons:—

•	1st Quarter.	2d Quarter.	3d Quarter.	4th Quarter.
American Parentage	26.73	26.18	24.22	22.93
Foreign Parentage	22.69	25.83	29.49	

 Deaths by age and parentage in each month and quarter, 1881.

IJ				7.5				***		-	IFE.				-	
R	Under I.	1 to 2.	2 to 5.	5 to 10.	10 to 15.	15 to 20.	20 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.	S0 to 90.	90 to 100	100 % ov'r	All ages.
	15 15 18	9 6 6	7 4 2	6 1	2 2	1 4 4	11 3 6	14 7 6	4 8 7	6 6 9	6 12 10	18 3 13	5 5	· i		88
	48	14	13	9	5	9	20	27	19	21	28	34	19	1		26
	15 16 7	4 6 4	11 11 4	283	1 1	4 1 1	11 6 4	7 10 9	3 7 7	9 7 10	10 10 7	9 8	9 3 5	1 'i		9 7
	38	14	26	13	3	6	21	26	17	26	27	25	17	2	•••	20
	25 39 17	8 6 3	14 8 4	4 7 8	i	3 1 1	6 2 5	2 5 6	3 6 9	4 9 4	5 2	1 6 5	5 6 4	1		10
	81	17	26	14	2	5	13	13	11	17	12	12	-15	4		24
	17 9 13	3 2 1	7 6 8	3 2 3	3	4 1	5 4 4	6 8 5	6 4 10	9 7 15	8 7 5	5 7 12	6 7 7	. 0102		778
1	39	6	16	8	4	5	13	19	20	31	20	24	20	4		90
	206	51	81	44	14	25	67	85	67	95	87	95	71	11		90
	15 11 14	4 6 5	6 6 3	2 4 3	i	9 5 5	19 11 14	9 6 9	4 6 6	13 11 7	12 4 7	7 4 7	4 3 4			200.00
	40	15	15	9	2	12	44	24	16	31	23	18	11			20
	11 14 12	8 5 9	973	4 2 6	3 2 6	3 2 3	18 16 18	9 14 9	12 11 8	978	7 10 2	6 8 4	4 8 3	1	2 24	10
	37	22	19	12	11	8	52	82	31	24	19	18	10	1	**	25
	43 27 23	12 13 9	7 9 9	3 5 6	3 1 2	4 4	12 17 12	7 11 12	11 8 6	2 10 5	897	3 5 3	4 2 2	ï	 'i	11 13 16
	93	31	25	14	В	8	41	30	25	17	24	11	8	1	1	31
	15 9 15	14 8 2	8 12 6	5 3 4	i	3 4 9	14 11 8	7 5 10	7775	3 6 10	5 7 10	374	5	1111		2000
9	39 200	95	20 85	12	3 22	10	33 170	22 108	19	19	22 88	14 61	35	3 5	1	1,1
				_							_			_	_	

.8.

7. 2.

6. 25. 25. 25. 25. 25.

Ratio ot deaths from each cause: to whole num-ber from known ber seuses. 7 Total. Disease, sex, age, parentage, and percentage from known causes. PARRNTAGE. Foreign. пвэ1тэт Л 90 and .19vo '06 O1 08 .08 of 07 LIFE. .07 03 08 .09 o1 05 9 .06 01 04 PERIODS '07 01 08 30 to 30. DIVISIONAL 12 to 30. 10 to 12. 5 to 10. .6 of 2 ban I grader 2. Ē l'nder l yenr. 2 **DEATHS**, 1881. .[gioT SEX. Females. Males. Drowning..... Bladder, Disease of...... Congestion of..... Accidents, (various)...... Inflammation of..... Burns and Scalds... Asthma..... Brain, Discase of..... TABLE XIX. CAUSES OF DEATH. Railroad.... Atelectasis, Pulmonum. Bones, Disease of...... Polsoning. A poplexy..... Ansemia..... Aneurism of Aorta. Abscesses. :

		BKX.) à	1810	MAL	DIVISIONAL PERIODS OF	100	6	LIN	.			-	PAR	PARENTAGE	9	-wnu
CAUSES OF DEATH.	Males.	Females.	Total.	Under I year.	bna i 2 Teban	.ê oj ŝ	.01 03 0	10 to 15.	15 to 90.	.08 01 08	30 to 40.	40 to 20.	.0n cs 08.	.07 03 00	70 10 80.	.08 03 06	90 & OYer.	American.	Foreign.	.fatoT	rom each o
Bronchitis	3	8	3	8	8	1	1-	i •	1 .	 	1:	100	1-	10	10	100	<u> </u> :	इ	38	3	8.49
Cancer (various)	=	11	88	-	:	Ė	i	:		:	ĠŦ.	63	00	t-	-	=	:	2	2	88	1.81
of Breast	:	9	2	Ì	i	÷	:	÷	÷	÷	÷		•	:	-	_	_	•	•	95	7.
" of Stomach	-	*	Ξ	:	:	:	•	:	i	:	Ø1		63	ø	91	$\frac{\cdot}{\cdot}$.	۲۰	•	C	20.
" of Uterus	:	36	38	i	:	:	÷	-	÷	:	_	09	00	63	;	÷		a	-	16	57.
Carbuncle	_	:	-	:	:	:	:	:	÷	<u>:</u>	:	-	:	$\frac{\cdot}{\cdot}$	÷	÷	:	=	:		9.
Cellulitis Pelvio	:	-	_	i	:	:	•	i	÷	:	÷	÷		÷	÷	÷	<u>:</u>	-	:	~	8
Child-birth	:	13	22	:	:	i	•	÷	:	8	t-	÷	:	<u>:</u>	÷	÷	•	30	ю	18	.
" Puerperal Convuls		01	<i>9</i> 1	:	•	:	•	÷	:	-	÷	÷	:	<u>:</u>	$\frac{\cdot}{\cdot}$	÷	•	91	:	6 4	ş.
" Puerperal Fever		7	7	•	:	÷	:	:	:	90	4	01	$\frac{\cdot}{\cdot}$	$\frac{\cdot}{1}$	÷	÷	-	67	11	7	<u>s</u>
Cholera Infantum	38	\$	20	8	35	•	Ť	:	i	÷		÷	:	<u>:</u>	<u>:</u>	÷	<u>:</u>	22	8	102	4.79
Cholera Morbus	•	GN.	•	:	:	_	•	÷	Ì	÷	:	*	<u>:</u>	÷	$\frac{\cdot}{\cdot}$	-	:	-	•>	•	8.
Colfe	Ø9	83	10	64	_	:	÷	÷	İ	:	\div	÷	$\frac{\cdot}{\cdot}$:	_	=	:	63	61	۰.	si si
Consumption	157	187	7	•	•	G1	69	-	8	8	ß	ä	83	9	۵	<u>;</u>	:	110	83	344	16.15
Convalsions	8	11	37	88	ю	93	:	=	÷	÷		$\frac{\cdot}{\cdot}$	$\frac{\cdot}{\cdot}$	÷	÷	:	.:	8	39	31	1.74
Croup	\$	क्ष	9	Ø.	15	S	4	Ä	÷	÷	\vdots	÷	<u>:</u>	:	÷	÷	-	11	85	48	8.9
Debility		33		2	-:	•	-	=	_	-	- :	_	_	_	=	-	- -	10	2	11	8.

uses.	UMOU UMOU	Ratio of de from each of to whole i her from ki	8.	si	3.8	8	5.45	8į	s.	s.	8	8	si	8.	* :	1.78	8	ä	8.
known causes.		Total.	E	1-	\$	61	118	-6	~	7	=	-6	10	1	60	8	-	0	- -
Vuo Vuo	PARENTAGE.	Foreign.	1	63	8	69	57	*	:	1-	•	64	60	00	~	2	-		=
from	PAR	American.	2	•	8	i	25	64		1-	6 0	90	91	a	01	11	:		
percentage		.T9V0 & 08F.	1:	:	_:	:	:	:	:	:	:	:	:	:	:	:	:	:	_
5		.06 c3 08	1:	:	_	_	:	:	:	69	_	:	:	:	:		:	*	_:
		.06 co 07		:	84	:	:	81	:	:	:	:	:	:	:		:	01	
Disease, sex, age, parentage, and	LIFE.	.07 oz 08	I i		7	:	_	:			<i>a</i>	:	69	•	_	_	•	:	
196,		.08 c3 08	:	94	1	:		1	_:	01	:	-	(PI	æ	:	93	:	:	
ea c	DIVISIONAL PERIODS OF	.00 to 50.			<u>:</u>	:	:	:	:		<u>:</u>	:	:	on	:	•	_	<u>:</u>	_:
par	RIOD	.04 nt 08	1 :	:	:	:	:	:	_	_		<u>:</u>	:		<u>:</u>		<u>:</u>	<u>:</u>	_:
zge,	F.	.08 01 02	<u> :</u>	64	<u>:</u>	<u>:</u>	•	64		<u>:</u>	~	<u>:</u>	-	_	_	2	<u>:</u>	<u>:</u>	_:
#	MAL	15 to 90.	<u> </u>	<u>:</u>		_:	*	_:_	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>			<u>:</u>	10	<u>:</u>	<u>:</u>	_:
6,9	71810	10 to 15.	<u> </u>			<u>:</u>	-6	<u>:</u>	<u>:</u>	<u>:</u>	_	_		<u>:</u>	<u>:</u>		<u>:</u>	<u>:</u>	
Se	Ā	5 to 10.	<u> </u>		_:_	_:	ಪ	_:_		<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	_ <u>:</u> _	:	*	<u>:</u>	<u>:</u>	_i
ă		\$ to 2.	<u>L:</u>	_i_		<u>:</u>	\$	<u>.:</u>	<u>:</u>	90	_	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>		<u>:</u>	<u>:</u>	_:
<u>.</u> :		ban I .g Tobau	l !	<u>:</u>	4	<u>.</u> :	8	<u>:</u>	<u>:</u>		:_		<u>:</u>	<u>:</u>	_			<u>:</u>	_:
1881		I Tebri year.	=	<u>:</u>	<u> </u>		· •	<u>:</u>				- <u>-</u>	<u>:</u>	10	<u>:</u>	:	<u>:</u>	<u>:</u>	_
		Total.	12	-	Ŧ	61	116	•		74	*	60	10	11	60	28	-	4	_
ВЕАТН В,	SEX.	Females.	6	_	18	-	25	93	7	80	7	7	1	10	64	2		64	-
_		Males.	F	6	82	O1		64		·	-	-	•	1-	-	38	~	9	
TABLE XIX.—CONTINUED.		CAUSES OF DEATH.	Delillity, Birth Premature	Diabetes	Diarrhea	" Chronic	Diphtheria	Dropsy	" of Chest	Dysentery	Enteritis	" Gastro	Epilepsy	Erysipelas	Fever, Malarial	" Typhoid	Gangrene	Senlle	Glands, Discase of

Ratio of de from each of to whole ber from kr	5.30	.38	90.	.88	.14	00.	11.	.05	11.	1.88	1.50	.75	48	83	114	.00	20
Total.	113	00	-	ţ=	90	0.5	613		50	40	60	16	6	5	02	1	-
Foreign.	25	*	П	7	00	П	01	-	Q1	16	6	4	00	77	01	7	***
American.	19	4		00	-	-	1	****	-	10.4	23	19	1	1	-	:	-
90 & OVer,	-	-	*	:	:	:	:	:	:	:	:	1	:	:	:	1	
'06 01 08	10	1	:	:	:	:	1	1	1	:	:	:	:	:	:	:	-
.08 01 02	88	-	-		-	:	:	-	1	1	:	-	01	:	-	1	
.07 01 08	25	0.1	-	-	1	1	1	1	:	1	1	4	1	-	1	:	-
.09 of 0a	15	04	1	1	-	-	-	1	:	1	1	01	C)	1	1	1	
40 to 50.	120	-	-	:	-	:	69	1	1	1	-	20	00	-	+	1	•
30 to 40.	=	-	***	-	-	1	:	:	:	:	1	9	1	O1	1	:	,
20 to 30.	-	:	7	1	1	1	:	1	00	:	:		1	-	1	1	0
12 10 50	-	:		:	:	-	:	-	1	1	:	-	-	:	:	1	
10 10 12	Q4	:	:	1	:	:	:	:	:	:	:	:	:	:	:	1	
2 io 10.	00	:	***	:	:	:	:	1	:	*	CA	:	:	:	:	:	*
2 (0 5.		***		1	:		1	:	:	9	6	•	1	:	:	:	
I and under 2.	-		:	:	:	:	:	:	:	13	9	:	:	:	-	:	
Under 1	3	:	1	7	-			:	10	17	15	:	:	:	1	1	
Total	3118	00	1	-	60	01	00		00	40	25	16	6	10	60	1	7.0
Lemujes.	99	2	*****	93	O.		1	-		30	16	11	10	01	-	1	4
Males.	150	93	-	*	-	C1	01	-	00	30	16	AQ.	4	90	04	1	10
CAUSES OF DEATH.	Heart, Discuses of	" Hypertrophy of	Teat	Hemorrhage	" from Lungs	" from Stomach	Hernia	Hip-Joint, Disease of	Homicide and Murder	Hooping Cough	Hydrocephalus	Insanity	Intemperance	" Delirium Tremens	Intussusception	Jaundice	Cidnava Disausa of

		SEX.	1				DIVI	DIVISIONAL PERIODS	AL F	ERIC		OF 1	LIFE.	1			P	PARENTAGE.	AGE,	-um
CAUSES OF DEATH.	Males.	Females.	Total	Luder 1	I and under 2.	3105.	5 to 10.	10 to 15.	15 to 20.	30 to 40.	40 to 20.	50 to 60.	.07 01 00	70 to 80.	.00 to 90.	30 & OVEL.	American.	Foreign.	Total.	Ratio of de from each es to whole n to whole n
Kidneys, Bright's Disease of	13	17	30	1	1:	1:	1	1	-	100	1	10	100	19	100		15		15 30	0 1.40
Laryngitis	-		1	:	:	:	:	:	:	:	1	-	1	:	:	:			1	10.05
Liver, Disease of	12	9	18			:	:			01	1	9	-	10	.:	:	-	0 1	13 1	18
" Inflammation of	.01	60	10	+:	:	:	-	:	:	*:		1	2	01	.:	-		- 01	00	10
Locomotor Ataxia	00	-	00	:	:	:	:		:	:	H	-	1:	:	:	:	-			00
Lungs, Disease of	1	1	64	:	:	:	1	:	1:	:	:	:	:	:	:	***			Oi.	60.
Malformations (all kinds)	90	7	15	15	:	:	:	:	:	*	-	:	:	:	:	:	-	11		15 .70
Malnutrition		01	04	04	:	**	-	:	:	:	:	:	:	:	:	:	04			00.
Marasmus	30	14	34	27	***		-	:	:	:	1		- 2	-		:	17		17 84	-
Measles	11	14	25	0	1-	*	- 04	-	-	-	100		:	:	;	-	10		13	1.17
Meningitis, Cerebro Spinal	-	t-	11	-	:	0.5	0	1	-	01	-01		-	:	-	- 1	-	-		20. 11
" Spinal	1	01	00	:	:	-		:	:	1	*	1	:	:	:	-			- 69	31.
Metritis	*****	1	-			-	-	**		-	1	:	:	:	-	***		-		00.
Old Age	27	99	7.18	****				** **	-			7	3		18	14	900		7 2	7.0 8.43
Ophthalmis Seonstorum		100	1 27			*		1		-	1	-	-	1	-				-	20.

edias eana: eana:	Total. Ratio of dofter to whole to whole bet from ker com ker	33.	160 7.51	14	1 .06	3 .14	1 .05	8 .14	1 ,05	8 .14	12 .56	45 2.11	4	8 .14	6 .28	4	6
PARENTAGE	Foreign.	80	86	8	_	01	i	60	П	64	7	38		-		69	-
PA	лвогісал	8	8	90	Ī		_	i	i		4	11	60	61	10	69	•
	90 and over.	1:		:	:	:	:	:	:	:	:	:	- <u>-</u>	:	:	:	
1	.08 cJ 08	000	00	-	:	:	:	- :	:	:		:	:	:	i	i	•
1	.08 o1 07	 :	=	_	_	:	:	•	:	:	01	:	:		64		•
Lire	.07 03 08	6	æ				:	:	:	:	_	:	:	i	:	•	_
	.00 01 05	-	8	64	:	:	:	:	:	-	:	:	:	_		:	۰
9	40 to 50.	1:	=	:	:	1	1	:	. :	1	9	:	:	:	09	:	•
DIVISIONAL PERIODS OF	.01 01 08	1	14	:	:	<u>:</u>	:	:	:	_	:	:	:	:	:		c
1 2	30 to 30.	<u> :</u>	7	:	:	i	:	G9	_	:	_	:	:	:	:	:	•
MAL	.0g cot d.I.	1	1	:	:	:	•	<u>:</u>	<u>:</u>	:	<u>:</u>	<u>:</u>	:	<u>:</u>	<u>:</u>	_	
181	.č1 ot 0f	:	01	:		:	:	_	:	:	:	_	:	_:	_:	<u>:</u>	
ă	5 to 10.		_	:	:	:	:	<u>:</u>	:	:	_	7	_	:	_	<u>:</u>	_
	2 to 5.	<u> </u>	2	•	<u>:</u>	:		<u>:</u>	<u>:</u>	:	<u>:</u>	- ま - ー	<u>:</u>	<u>:</u>	<u>:</u>	•	
1	l and Lader S.	<u> </u>	18	04	<u>:</u>	:	<u>:</u>		<u>:</u>	<u>:</u>	:	*	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	
	Under I year.	1:	3	•	<u>:</u>		<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	61	~	_	<u>:</u>	<u>:</u>	•
	.latoT	8	160	*	-	∞		89	-	•	13	3	•	80	8	•	•
SEX.	Females.	1	88	•		64		1	_	2	7	18	+	83	2	64	ç
	Males.	64	1	20	7			69	:	_	90	25	:	:		61	G
	CAUSES OF DEATH.	Pleurisy	Pneumonia	" Congestion of Lungs.	Prostate, Disease of	Pupura Hemorrhagica	Pustule, Malignant	Pyæmia	Quinsy	Rectum, Disease of	Rheumatism	Scarlatina	Scrofula	Skin, Disease of	Spine, Disease of	Stomach, Disease of	it Indommetion of

Ratio of deaths from each oanse to whole num-ber from known causes. 8 8 8 4 3 100.00 Disease, sex, age, parentage, and percentage from known causes. 9,145 Total. PARENTAGE. 1.146 Foreign. 8 **Атре**Новп. 1 90 % OVEr. 156 106 .06 ot 08 10 00 90 237 198 158 186 175 LIFE. .07 oz 08 .08 of.08 ò 40 to 20° DIVISIONAL PERIODS 30 to 40° .08 or 08 8 12 to 30. 8 10 10 12. 8 .01 oz 8 146 166 .ð oð <u>£</u> Under 1 year, I and under 2. TABLE XIX.—CONTINUED. DEATHS, 1881. 12 2.145 Total. SEX. Females. 10. 9 Males. Totals Suicide Surgical Operation..... Congenital..... CAUSES OF DEATH. Tabes Mesenterica,..... Tetanus and Tris. Nascen Teething..... Throat, Disease of..... Syphilis Tumor Unknown Tuberculosis.

Whole Year.	55	7.	*	25	23	53	105	*	344	37	69	43	116	14	19	11	88	60-	121	40	389
Fourth Quarter.	15	16	22	13	13	4	7	C)	88	9	20	14	98	2	4	1	10	*	24	4	4
December,	-	9	00	6	+	******	-	-	255	*	-	01	9		F	-	9	64	13	1	1
November,	100	1-	9	00	4	4	+	1	19	O.	14	C/s	16	00	00	-	04	-	Į+	1	
October.	100	90	10	-	9	-	60	F	22	:	9	10	14	G4	-	1	-01	-	*	G1	80
Third Quarter.	16	12	15	30	13	00	96	01	90	6	7	36	30	9	11	-	10	****	30	64	7
September.	9	4	-	4	00	O1	18	*****	88	01	7	02	15	1	Q8	:	7	1	-	10	69
August.	00	10	00	*	9		35	-	36	00	-	14	=	60	64	-	00	1	12	12	00
July.	90	1	9	******	4	-	43	01	36	4	69	6	-	G)	-	-	00	-	11	2	01
Second	11	83	56	18	17	8	-	***	96	13	1-	-	84	-	-	10	11	i	36	6	12
nue-	01	00	7	7	6	1	1		88	9	01	:	01	:		00	63	-	10	4	00
May.	-	10	11	90	60	00	*****	1	33	7	80	-	=	-	1	-	G1		14	7	9
April.	0	2	11	9	9	9	-	-	58	90	03	-	15		H	-	8	-	12	1	60.
First Quarter.	138	18	17	14	61	13	H		90	0	15	G4	81	01	00	10	I.e.	1	31	2	6:
March.	7	1-	ic.	00	00	OI	-	:	123	10	01	:	00		-	7	1	i	19	4	*
February.	0	+	-	10	9	*	-	-	81	64	9	-	1-	Q1	-	99	60	1	6	1	-
January.	-	10	10	9	00	E-			41	4	1+	1		-	01	63	80	-	10	***	7
CAUSES OF DEATH.	Accidents (all kinds)	Apoplexy	Brain, Diseases of	Bronchitts	Cancer (all kinds)	Child-birth	Cholera Infantum	Cholera Morbus	Consumption	Convulsions	Croup	Diarrhoa	Diphtheria	Dysentery	Enteritis	Erysipelas	Fever, Typhoid	" Malarial	Heart, Diseases of	Hooping Cough	Hydrocephalus

Table XX.—Continued. Deaths, 1881. Season and Disease.

																	_		_		
Whole Year.		2	7	\$	2	ಷ	8	=======================================	15	ä	2	11	2	\$	23	*	•	*	2	101	12
irth narter.	Eva	•	49	2	*	1-	-	91	81	•	•	9	63	•	•	_	6 1	4	•	4	3
.Ted atte	Dec		_	•	0 1	6 1	i	_	2	-0	•	~	:	_	:	:	_	:	*	Ĕ	Ē
rem ber.	MOV	-	99	•	:		Ä	:	1-	•	:	ă	*	*	F4	:	_	+	:	Ē	2
oper.	æ0 —	-	~	Ö	_	•		_	۰۰-	i	24	2	:	_	-m-		-	-	20	1	≝
hi narter.	TPI	•	8	1-	1-	13	a.	61	8	10	•	2	94	10	•	60	**	91	-	3	Ž
.nedmet	dəg	7		_oi	•	- 01		-	*	93	**	04	01	:	61	~	i	Ī	•	2	٤
traç.	Bay	-	-	- et .	_ 04	່ ພັ	i	Ī	9	:	04	•	i	=	_	-	*	=	Ī	ž	3
•4	լաբ	*	/~4	61	~	•	8 0		9	en	-	61		•		_	_	_	**	23	薑
ond narter.	10 10	'-	+	=	1-	2	*	6	2	-	10	€	:	2	10	4	04	2	_	2	ź
.0	unr	Ī		15	60	- es-	•	•	1-	-	_	Ē	•	+	_	*	_	64	:	35	Ē
٠.	учву	m		-4-	•••	•	-		*	•	64	_ &	•	80	-	i	_	•	_	38	Ę
.11.	τqΛ	<u></u>	_e,	:			•		- • -	Ol	-04	*		-1-	**	_	:	*	:	2	8
t Brior.	Fire Qu		- 61	13		- iä	- -		7	•	*	9	1-	E	2	43	01	•	*	ţ	8
ch.	TaM	-		-Ţ-	- 6		:		•	-	60	35	61	-		64	-	~	69	2	2
·Vraury.	Lep1			-₩			- : -	~	▼	64	:	88	•	•	_	64	:	•	:	13	<u>इ</u>
.V.IAL	uvr		_	~	-	01	-		Ğ.	83		84	_	-0	*	_				9	Ē
CAUSES OF DEATH.		nasnity	ntemperance	Kidneys, Diseases of	Liver, Diseases of.	Marasmut	Heasles	Meningitis, Corebro Spinal	Old Age and Gangrene Senile	Paralysis	Peritonitis	Pneumonia and Congestion of Lungs	Rheumatism	Scarlatina	Stomach, Diseases of	Sulcide	Teething	Tuberculosis	Unknown	Other Causes as given	Totale

NOTES ON THE CAUSES OF DEATH,

DURING THE YEAR 1881.

These notes explain some of the figures of Tables XIX. and XX.; there are also additional statistics given here, and these are compared with similar statistics of other years.

The percentage of deaths from the different causes is always reckoned on the number of deaths from known causes.

There were 2,145 deaths in Providence during the year 1881; of these there were 15, or 0.70 per cent., of which the cause was not stated; the number from known causes, therefore, was 2,130, and on this number all percentages are reckoned. The number of deaths for which no cause was stated was 2 less, and the percentage 0.12 less, than in 1880.

During the twenty-six years, 1856-1881, the whole number of deaths was 36,724, of which 565, or 1.54 per cent., were from unknown causes; 36,159 from known causes. In the forty-two years, 1840-1881 inclusive, the total number of deaths was 49,869; 48,302 from known causes; and 1,567, or 3.14 per cent., from unknown causes.

Table XX. shows the prevalence of different diseases at different seasons of the year. Of diseases of the respiratory system, we find a large number in the first and second quarters of the year; of diarrhoeal diseases, 77.47 per cent. were in the third quarter; more than half of the deaths from hooping cough, also, were in the third quarter. There was no particular epidemic during the year; diphtheria, from which there were nearly twice as many deaths as in 1880, was very evenly distributed through the year.

Abscesses. The number of deaths from abscesses in 1881 was 7, or 0.33 per cent. of all from known causes; this was 4 more than in

1880, and the per cent. 0.18 more. Three were of the pelvis, one of the parotid gland, and one of the hip; two were not specified. Other abscesses are reckoned under the head of disease of the organ affected.

Males, 2; females, 5; total, 7; or 0.33 per cent, of all. American parentage, 5; foreign parentage, 2.

The statistics for the twenty-six years, 1856-1881, are:-

Number of deaths from all known causes, 36,159.

From abscesses, 101; or 0.28 per cent. of all.

Males, 51; females, 50.

American parentage, 47; foreign parentage, 54.

Accidents. There were 52 deaths from accidents in 1881; 13 less than in 1880, and 2.44 per cent. of all. There were 7 deaths from burns or scalds, 4 of which were of children under five years old; 8 from drowning; 8 from falls, all of whom were over 40 years old, and 3 of them were women over 80; 5 from poisoning, 2 of which were by leakage of illuminating gas; one by acid fumes; one by belladonna (given by the mistake of a druggist's clerk); and one from an overdose of morphine, self-administered; 9 deaths from railroad accidents. There are 18 accidents under the heading of " various"; they were as follows: one had the "skull fractured by a horse's hoof"; one the "horse ran away with him"; one was "thrown from a horse"; two were "thrown from a wagon or carriage"; three were "run over", one by a dray, one by an ice-cart, one by a furniture wagon; one was " suffocated by the bed-clothes"; one was "overlaid in bed"; one died from "injury at birth"; one died from "concussion of the brain"; one from "injury of lumbar vertebra"; one from the "fall of a pulley-block on his head"; and one from a "pistol shot in the bowels"; the latter was a child a year and a half old, who was killed by the accidental discharge of a pistol in the hands of its mother.

The number of each sex and parentage were :-

Males, 37; females, 15; total, 52; or 2.44 per cent. of all. American parentage, 19; foreign parentage, 33.

For the twenty-six years, 1856-1881:-

Males, 776; females, 262; total, 1,038; or 2.87 per cent. of all.
American parentage, 401; foreign parentage, 637.

Ancurism. There was but one death from this cause in 1881; one American female died of ancurism of the aorta. The statistics for this cause for the twenty-six years, 1856-1881, are:—

Males, 22; females, 5; total, 27; or 0.07 per cent. of all.
American parentage, 10; foreign parentage, 17.

Apoplexy. The number of deaths reported from this cause in 1881 was 74, or 3.47 per cent. of all from known causes; the number was 26 more, and the per cent. 1.15 more than in 1880; it is a much larger number of deaths from this cause than ever occurred before in one year; the largest number before recorded being 56 in 1879. The statistics as to sex and parentage for 1881 are as follows:—

Males, 38; females, 36; total, 74; or 3.47 per cent. of all.
American parentage, 52; foreign parentage, 22.

For the twenty-six years, 1856-1881:-

Males, 392; females, 406; total, 798; or 2.21 per cent. of all.
American parentage, 558; foreign parentage, 240.

For further particulars on this subject, see "Paralysis."

Brain, Diseases of. The number of deaths in 1881 from various diseases of the brain was 88, or 4.14 per cent. of all; 8 more deaths, and 0.27 per cent. more than in the preceding year. Eight deaths were from "congestion," and 53 from "inflammation of the brain." The statistics of sex and parentage are:—

Males, 43; females, 45; total, 88; or 4.14 per cent. of all.

American parentage, 42; foreign parentage, 46.

For the twenty-six years, 1856-1881, the figures are:—
Males, 758; females, 650; total, 1,408; or 3.89 per cent. of all.
American parentage, 734; foreign parentage, 674.

Bronchitis. This was the cause of 53 deaths in 1881, or 2.49 per cent. of all. More than half of the decedents were less than two years old, and 28.30 per cent. were over 60. The deaths were 3 less than in 1880, and the rate 0.22 per cent. less.

The particulars as to sex and parentage for the year were as follows:— Males, 33; females, 20; total, 53; or 2 49 per cent. of all-American parentage, 27; foreign parentage, 26.

For the twenty-six years, 1856-1881:-

Males, 234; females, 278; total, 512; or 1.42 per cent. of all.

American parentage, 231; foreign parentage, 281.

Cancers. The number of deaths from cancers in 1881 was 65, or 3.05 per cent. of all; in 1880 the number was 68, or 3.30 per cent. of all. In Table XIX. 10 are given as cancer of the breast; 8 more than in 1880; 11, cancer of the stomach; 3 more than in 1880; and 16 as cancer of the uterus; the same number as in the preceding year. Of the 28 under the heading "various" there were two of which the location was not specified; the others were: 11 of the liver; 3 of the face; 3 of the pancreas; 2 of the bowels; and 1 each of the eye, throat, colon, pylorus, kidney, bladder, and foot.

The statistics of sex and parentage are as follows:-

Males, 18; females, 47; total, 65; or 3.05 per cent. of all. American parentage, 37; foreign parentage, 28.

For the twenty-six years, 1856-1881:-

Males, 248; females, 651; total, 894; or 2.47 per cent. of all.

American parentage, 562; foreign parentage, 332.

Of the 894 decedents in twenty-six years, 72.82 per cent. were females; and 62.86 per cent. were of American parentage.

Child-birth. There were 29 deaths from child-birth in 1881, including 2 from puerperal convulsions, and 14 from puerperal fever. This was 3 more deaths, and 0.10 per cent. more than in the previous year. The 29 deaths were 1.36 per cent. of all from known causes, and 1.04 per cent. of the 2,780 mothers who bore children during the year; the decedents were 13 of American, and 16 of foreign parentage. In the twenty-six years, 1856–1881, the number of deaths from this cause was 434; 185 of American, and 249 of foreign parentage. This was 1.20 per cent. of all deaths from known causes in that time, and 0.85 per cent. of the 51,222 mothers bearing children during the same period.

Cholera Infantum. There were 102 deaths in 1881 from this disease; 2 more than in the previous year, and more than in any year

since 1877; the percentage, however, was not quite so large as in 1880. The following statistics are given:—

Males, 58; females, 44; total, 102; or 4.79 per cent. of all.

American parentage, 85; foreign parentage, 67.

The following are the statistics of sex and parentage for each of the twenty-six years, 1856-1881, and for the whole period:—

	SE	x.	Paren	TAGE.	Whole
1	fales.	Females.	American.	Foreign.	Number. Percentage
1856	85	22	16	41	575.5
857	27	23	28	24	505.56
858	27	22	18	86	494.9
859	14	19	18	20	83
860	35	83	26	42	687.00
861	29	82	25	36	616.09
1861	25	19	17	27	444.90
1863	36	80	19	47	665.5
864	29	29	18	40	584.65
1865	20	80	18	87	504.9
1886	80	17	15	82	474.6
1867	24	25	23	96	49
1868	38	89	23	48	706.4
1869	85	80	27	88	655.2
1570	48	45	87	56	937.4
871	80	83	26	86	62
1872	72	79	58	93	1519.5
1873	42	50	89	53	925.4
874	73	62	44	90	1346.8
875	62	74	50	86	136
876	66	56	48	79	122 6.5
877	70	52	43	79	1226.8
1878	85	85	26	44	70 8. 5
879	80	29	28	81	59
1880			41		1004.8
1861	RQ.		85	67	1024.7

Cholera Morbus. The number of deaths reported under this head is always small. The number in 1881 was 4; the same as in the preceding year.

Males, 2; females, 2; total, 4; or 0 19 per cent. of all. American parentage, 1; foreign parentage, 3. For the twenty-six years, 1856-1881, the whole number was 120, or 0.33 per cent of all.

Consumption. The number of deaths from this cause always far exceeds that from any other, and has increased the past few years. The number in 1880 was 322, and the percentage 15.60; in 1881 the number was 344, and the percentage 16.15. The decedents were divided as follows as to sex and parentage:—

American parentagemales,	49females,	, 67total, 116
Foreign parentage	108females,	, 120total, 228
		
Totalsmales,	157 females,	. 187total, 844

For the twenty-six years, 1856-1881, the statistics are:-

Whole number of deaths from known causes, 36,159. Deaths from consumption, 6,034; or 15.69 per cent. of all. Annual average, 233.

American parentage......males, 1,218..........females, 1,573.......total, 2,791

Foreign parentage......males, 1,529........females, 1,714.......total, 3,243

Total......females, \$,267......total, 6,034

Having estimated the population of the city at 107,500, in the middle of the year 1881, we find the death rate from consumption to be one in 312 of the inhabitants, or 3.20 in each thousand persons. The average annual rate for the fifteen years, 1840–1854, was one in 209.9; for the twenty-seven years, 1855–1881 inclusive, it was one in 317.1.

The figures already given show that, for the whole period, considerably more than half of all the decedents from consumption were of foreign parentage. This has been true since 1874; but in every year previous to that date, with one exception, there were more decedents from consumption of American than of foreign parentage. The following figures show the percentage of each class of decedents from consumption for each of the last eight years:—

De	aths from		•		
Con	sumption.	American.	Per cent.	Foreign.	Per cent.
1874	270	106	89.26	164	60.74
1875	299	114	88.13	185	61.87
1876	284	120	42.25	164	57.70
1877	294	116	44.22	178	55.78
1878	805	107	35 08	198	64.92
1879	293	116	39.59	177	60.41
1880	822	111	84.47	211	65.58
1881	344	116	83.72	228	66.28
Total					

In 1873, there were 117 decedents of American and 114 of foreign parentage, from consumption; in 1874, the decedents of foreign parentage from consumption were 164, an increase of 50, and in no year since that date has the number of decedents of foreign parentage from consumption been less than 164. On the other hand, the number of decedents from consumption, of American parentage, has not increased, and has but slightly changed for many years, notwithstanding the increase of population. There were 120 decedents from consumption, of American parentage, in 1857; 123 in 1860; 122 in 1870; 117 in 1872 and 1873; 116 in 1877; and 116 in 1881. The sudden increase of decedents from consumption, of foreign parentage, in 1874, was mostly owing to the annexation in that year of the Tenth Ward, with a large foreign population.

The following shows the number and percentage, according to parentage, of the decedents from consumption in Providence, in periods of five years each, from 1856 to 1880 inclusive, and also for the year 1881:—

50 77		43.55	Consumption. 930
50 77	=		
	480	49.23	97
50.57	520	49 48	
44.78	682	55.22	
38.05	928	61.95	1,49
33.72	228	66.28	844
		- -	33.72 22866.28

In 1856, the living population of the city was 59 of American and 41 of foreign parentage in each 100; the decedents from consumption, in the same year, were 58 of American and 42 of foreign parentage in each 100. In 1881, the living population was 46 American and 54 foreign in each 100; but the decedents from consumption were 33.4 American and 66.6 foreign in each 100.

But the true rate of mortality from consumption, in the two classes of the population, is best shown by a direct comparison of the number of deaths with the actual population. At four different periods in Providence we have had censuses showing the parentage of the population; and at the same periods we have had full and accurate reports of the decedents from consumption, by parentage. Taking these periods and comparing the decedents from consumption, by parentage, with the population by parentage, we find the following results for the city of Providence:—

POPULATION OF AMERICAN PARENTAGE.

1856-One death from consumption in 268.2.

1865—One death from consumption in 310.1.

1875—One death from consumption in 415.0.

1880-One death from consumption in 485.2.

POPULATION OF FOREIGN PARENTAGE.

1856 -One death from consumption in 262.6.

1865—One death from consumption in 248 8.

1875—One death from consumption in 288.4.

1880—One death from consumption in 268.0.

TOTAL POPULATION.

1856—One death from consumption in 266.

1865-One death from consumption in 280.

1875—One death from consumption in 886.

1880-One death from consumption in 826.

These figures show several most important facts:-

1. In the population of American parentage in Providence there has been a very remarkable change in the mortality from consumption, during the last twenty-five years. In 1856 there was, in this

ation, one death from consumption in 268.2; in 1880 there was one death in 435.2. This decrease, as seen from the figures, een continuous during the whole period.

In the population of foreign parentage, in Providence, the f mortality from consumption has changed only very slightly in st twenty-five years. In 1856 the rate was one death in 262.6 population; in 1880 it was one in 268.0; and during this it has been both slightly higher and lower than these rates. rate of mortality from consumption, among the population of n parentage, in 1880, was almost precisely the same as the rate population of American parentage in 1856.

me of the effects of this change in the rate of mortality from mption are as follows:—

There were 111 deaths from consumption in 1880, in the popon of American parentage, in Providence; if the rate, or the per in proportion to population, had been the same in 1880 as in , there would have been 180 deaths from consumption instead of

There were 211 deaths from consumption in 1880, in the popon of foreign parentage, in Providence. If the rate in this popon had been the same as in the American population, the numof decedents of foreign parentage, from consumption in 1880, ld have been 130 instead of 211.

ther illustrations might be given showing the very great change has taken place in twenty-five years in the mortality from conption, and it would be interesting to inquire into the causes of the oge, but this must be deferred to a future time.

he following table shows numerous and important facts in regard he decedents from consumption in Providence, in each of the aty-six years, from 1856 to 1881, inclusive:—

DEATHS PROM CONSUMPTION IN PROVIDENCE, TWENTY-SIX YEARS, 1856-1881.

Sex; parentage; season; age; whole number, and percentage from known causes.

Sec Sec
137 588 588 514 590 544 590 544 590 545
125 100 107 108 108 108 119
19 12 11 17 15 22 20 10 15 14 25 18 18 18 18 18 18 18 1
14 10 14 10 18 8 18 11 12 13 14 10 10 19 10 10 10 10 10 10 10 10 10 10 10 10 10
20 15 18 18 18 13 21 14 19 18 15 16 19 20 15 26 18 27 38 39 39 38 14 50 15 18 20 15 20 15 20 15 20 15 20 15 20 15 20 15 20 15 20 15 20 15 20 20 20 20 20 20 20 20 20 20 20 20 20

DEATHS FROM CONSUMPTION IN PROVIDENCE, TWENTY-SIX YEARS, 1856-1881.—CONTINUED. Sex; parentage; season; age; whole number, and percentage from known causes.

Total	2011 11415 1415 1415 1415 1415 1415 1415 1415 1415 1415 1415 1415 1415 1
1881.	8888 888 44001-68688800
1880.1	\$188 E8E 4440x8E848800
	\$29% H&8
878.	5555 ESH -4-4-8758888H
1876, 1877, 1878, 1879	ENTER END 4-04-0850-85-
1876.	8286 859 44-00,88658890
	ECT E 588 0 - 0 888 688 88 1
1873, 1874, 1875.	8588 888 010 010 04-21:64880
1873.	2248 285
1872.	622 828 64-068 828 8800
1871.	8448 624
1868, 1860, 1870, 1871	2882 885 www.ww.8148984
1869.	5848 PLW word-28348684
	## ## ## ## ## ## ## ## ## ## ## ## ##
1866, 1867,	2144 686 4u section
1866.	58840 5822 Leura 65228 Ergin
1865.	6448 888 0 000 4 0 0 0 0 0 0 0 0 0 0 0 0
1863, 1864, 1865.	7444 558 88486614844
	111111111111111111111111111111111111111
1862.	8886 888 arenar8648900
1860, 1861.	64428 688 4404445548488
	6888 E125 C84 +48848421
1857, 1858, 1859.	\$480 CBD .80-800000000.
1858.	5004 640 H44044
	12 12 12 12 12 12 12 12 12 12 12 12 12 1
1856.	64440 6654 u L 646155496 nu
CONSUMPTION.	Pirst quarter Second quarter Third quarter First 4 months First 4 months AGR: Under 1 year Linder 1 year 2 and under 2 2 and under 2 3 and under 3 5 and under 3 6 and under 3 70 and under 3 80 and over 90 and over
0	First Secont Third Fourt Fourt Fourt Fourt Fourt Third Third 1 a 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2

Convulsions. The number of deaths reported from convulsions in 1881 was 37, or 12 less than in 1880; 28 of the decedents, or more than 75 per cent., were less than a year old. The particulars, as to sex and parentage, are:—

Males, 20; females, 17; total, 37; or 1.74 per cent. of all. American parentage, 18; foreign parentage, 19.

For the twenty-six years, 1856-1881:-

Males, 515; females, 419; total, 934; or 2.58 per cent. of all. American parentage, 389; foreign parentage, 545.

Croup. There were 49 deaths from croup in 1881, which was 20 more than in the preceding year, and was 2.30 per cent. of all deaths from known causes. They were divided in the different seasons of the year as follows:—

1st quarter15	3d quarter
2d quarter 7	4th quarter

There were 9 decedents less than a year old; 12 between one and two years; 23 between two and five; 4 between five and ten; and I who was more than ten. The following are the statistics of sex and parentage:—

Males, 24; females, 25; total, 49; or 2.30 per cent. of all. American parentage, 17; foreign parentage, 32.

For the twenty-six years, 1856-1881:-

Whole number of deaths, 816; annual average, 31.

Males, 409; females, 407; total, 816; or 2.26 per cent. of all.

American parentage, 331; foreign parentage, 485.

The following table shows the number of deaths from croup for each of the twenty-six years, 1856–1881 inclusive; also the percentage each year, of the whole number from known causes; and the particulars as to sex, age, parentage, season, locality and color:—

Total	818	400	331	======================================	22 25 25 25 25 25 25 25 25 25 25 25 25 2
1881.	2.30	28	17.00	t-00101000101-401-1-	
1880.	1.1.29	× 01	1-81	9 :01 ::::::::::::::::::::::::::::::::::	8 .4-8 .000
1879, 1880	2.13	16	8181	\$100 m m m m m m m m m m m m m m m m m m	00-1000mg, 10, 00
	1.93	18	808	**************************************	2-2-03-4-I 2
1877. 1878	2.07	188	14	0000000	F-84 - 01 - 01 00 02
1876.	8.48	00 00	1288	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	∞-∞4×0×0×5×5×
1875.	2.95	25.53	19	OE	6 01 4 1445
1874, 1875, 1876,	1.48	10	52	0,000	4
1873.	1.23	15	15	0.00	44444 .000
1870, 1871, 1872,	1.71	13	22	00 01 00 01 01 01 4 0	4 1-01444000 - 4
1871.	25.85 25.85	178	15	00001-046	∞-∞-∞-∞
1870.	2.16	118	25	. *************************************	SHED . C1010100 . 0
1869.	1.53	10	13	mamio,	00
1868.	1.01	6.0	41-		D
1867.	9.01	10	10		40100-000-0-00
1866.	20,00	11	123	0-01 00 m	
1865.	2.44	118	e 0	************	poi 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2 - 2
1864.	3.59	88	19	# # # # # # # # # # # # # # # # # # #	20 01 00 04 L 20
1863.	90 T.	13	=31	\$1 50 50 50 51 1 1 51 1 1 1 1 1 1 1 1 1 1	@ -+ w to g 00
1862,	3.51	14	15	0000044 0000	88000
1861.	40.00	13	15		40100001-4
1860.	1,88	11	44	81-81-4-10	wat to da wing 4
1829.	2 24	18	22	F :00000	01 :00 00 00 10-
1838.	201	15	515	444000	000000000000000000000000000000000000000
1857.	8.11	16	138	8-4-8-4-8998	00 400 00 -0 10 · · · ·
1800	2.05	16	. S	**************************************	64-1016-0-4-0
	uber e		B:		

The statistics of the five diseases for the twenty-six years, 1856-1881, are:—

Number of deaths, 3,962; annual average, 153.

Males, 1,978; females, 1,984; total, 3,962; or 10.96 per cent. of all.

American parentage, 1,609; foreign parentage, 2,353.

The proportions between the two classes of decedents in 1881 were as follows: 145.9 decedents of foreign parentage to each 100 of American parentage; or 59.34 of foreign and 40.66 of American parentage in each 100 decedents. Of decedents from all causes during the year the proportions were 114.7 of foreign to each 100 of American parentage; or 53.43 of foreign, and 46.57 of American parentage in each hundred.

Of the 2,803 children born in 1881, 51.30 per cent. were of foreign parentage, and 48.70 per cent. of American parentage; the fact that there are so many more children of foreign than of American parentage, and the additional fact that these diarrheal diseases prevail mostly among young children, make the disproportion between the two classes of decedents less striking; of the 177 decedents from these causes in 1881, 145, or 81.92 per cent., were children less than two years old.

The following table shows the number of deaths from each of these causes, with the number from the whole class of causes, and their percentage of all from known causes, reported in Providence during the period from 1840 to 1854, inclusive; during each year, from 1855 to 1881, inclusive, and in the aggregate during the whole period of 42 years, 1840 to 1881, inclusive:—

NOTES ON THE CAUSES OF DEATH.

1st quarter22	3d quarter
2d quarter28	4th quarter

Of the 116 decedents, 105, or 90.52 per cent., were less than ten years old. The statistics of sex and parentage are:—

Males, 65; females, 51; total, 116; or 5.45 per cent. of all.

American parentage, 59; foreign parentage, 57.

During the four years' epidemic of diphtheria, 1876–1879, inclusive, the decedents of foreign parentage were greatly in excess of those of American parentage, the proportion being 56.39 of foreign, and 43.61 of American parentage in each 100; previous to the epidemic the reverse of this was true, the figures being 58.68 of American, and 41.32 of foreign parentage in each 100 decedents. In 1880 and 1881 the American class is again in excess; in 1880, 57.38 per cent. of American, and 42.62 of foreign parentage in each 100 decedents; in 1881, 50.86 of American, and 49.14 of foreign parentage in each 100.

The first cases of diphtheria in Providence were reported in 1858; for the twenty-four years, 1858–1881, inclusive, the statistics are as follows:—

Deaths from known causes in 24 years, 34,231.

From diphtheria, 1,270; annual average, 52.

Mules, 640; females, 630; total, 1,270; or 3.71 per cent. of all.

American parentage, 621; foreign parentage, 649.

The number of deaths in 1881 was more than twice the annual average.

In the following table are the statistics as to sex, parentage, season and locality, for each of the twenty-four years, 1858–1881, and for the whole period:—

Dropsy. This was stated as the cause of 5 deaths in 1881; 2 males, 3 females; 2 of American, and 3 of foreign parentage; total, 5; or 0.23 per cent. of all.

Erysipelas. There were 17 deaths from this disease in 1881; 8 more than in 1880, and 0.36 per cent. more.

Males, 7; females, 10; total, 17; or 0.80 per cent. of all.

American parentage, 9; foreign parentage, 8.

For the twenty-six years, 1856-1881, the statistics are:—
Males, 94; females, 102; total, 196; or 0.54 per cent. of all.
American parentage, 114; foreign parentage, 82.

Fever, Typhoid. The number of deaths from this cause in 1881 was 38; 14 less than in 1880, and 1.78 per cent. of all. The particulars of sex and parentage are:—

Males, 18; females, 20; total, 38; or 1.78 per cent. of all.

American parentage, 17; foreign parentage, 21.

The same statistics for the twenty-six years, 1856-1881, are as follows:—

Whole number, 1,113; annual average, 43.

Males, 571; females, 542; total, 1,113; or 3.08 per cent. of all.

American parentage, 621; foreign parentage, 492.

There have been a few deaths reported simply as fever; some as "bilious fever," and about 30 from typhus fever. There was also in 1869 one death from "intermittent," and one from "malarial" fever; in 1880 there was one death from malarial fever; these are all included in the foregoing figures for the twenty-six years; we have not, however, included the three deaths from malarial fever which occurred in 1881, as they indicate what appears to be a new development of the disease in this city, and it seems desirable to keep a separate account of them. The figures for the twenty-six years are found in the following table, which gives all the details of sex, parentage, color, season, age, locality and percentage from known causes, for each of the twenty-six years, 1856–1881, and for the whole period. This table has not been published before, and in making it up some errors in previous reports have been corrected, and the figures, as here given, are believed to be correct:—

Sex; parentage; color; season; age; locality; whole number, and percentage from known causes.

4.00		200000000000000000000000000000000000000	
96 yrs Total	207 263 643		888888888
1881	212		
1880.	25.80	H	828040000
1879.	128		
1878.	138		
1877.	222		****************
1876.	19.75		64 10 14 10 10 10 10 10 10 10 10 10 10 10 10 10
1875.	Bass		200000000000000000000000000000000000000
1874.	0 m 4		4-5100000000
1873.	111	. Land Collegan . L	
1873.	120	1 .92 121 25	- 2000 2000 :
1871.	212	H :00048F-001 :1	0,0114100000
1870.	348		: @ @ # # # # # # # # # # # # # # # # #
1869.	101		: www.manaa.
808	16.26	01	F-0101-0001-401-1
1867.1	100	L 0.4-0100 L	540000000
1866.	85.38	4 2 20 4 2 10 10 20 20 4	8100004
1865.	14 8	H .00000 H H 000000	@70 Ka G G G G G G G G G G G G G G G G G G
1864.	13		08000000
1863.	212		4040000
1862.	133.5		H .001040
1861.	272		4014010014
1860.	10		w-00444
1859.	10.00		41-6-6000
1858.	108	94KB8HHH	4 .010004
1857.	401-		F-01400-44
1856.	16.8	H .24 .00040H	20 00 00 00 00 00 00 00 00 00 00 00 00 0
YER.	s	884884888	11111111111
FE	months 4 month months	unde unde unde unde	E-HER-DEEN
поп	frst 4 m econd 4 hird 4 n	nder and mader	ard I I V
TYP	Firs Seco Thir	AGE 55 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	WW

This table shows that there is very little difference between the number of males and of females who die from typhoid fever. Of the two classes of parentage, the proportions are 126.2 decedents of foreign parentage to each 100 of American parentage; or 55.79 of foreign, and 44.21 of American parentage in each 100.

As to the season at which this disease chiefly prevails there can be no question, as is shown by the following figures, giving the number and percentage for each quarter:—

	Number.	Per cent.	Number.	Per cent
1st quarter	161	14.47	3d quarter305	
2d quarter	162	14.55	4th quarter485	43.35

In the first and second quarters the numbers are almost the same, and both together do not equal the number in the fourth quarter of the year. Dividing the year in three sections the numbers and proportions are as follows:—

	Number.	Per cent.
First four months	207	
Second four months	263	
Third four months	643	

The decedents are of all ages from the oldest to the youngest, but by far the larger number are between 15 and 40 years of age; the following table shows the number and percentage in each division of age, for 1881, and for the twenty-six years, 1856-1881:—

PERIODS.	Under 1 year.	land under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	70 and under 80.	So and ander 50.	90 and over.	Total.
1881.			13		77										
Number		1	1	2	1	5	13	6	4	2	1	1	1		25
Per cent		2.63	2.63	5.26	2,63	13.16	34.22	15.79	10.53	5.26	2.63	2.63	2.63		1(0.00
1856-1881.	-														
Number	8	12	41	79	77	173	306	152	95	83	51	28	7	1	1,115
Per cent	.72	1.08	3.68	7.10	6.92	15.54	27.49	13.66	8.53	7.46	4.58	2.52	.63	.09	100.60

The number of deaths in each year, and the aggregate and average age of the decedents were as follows:—

Number of deaths; aggregate and average age	Number	of	deaths	:	aggregate	and	average	age
---	--------	----	--------	---	-----------	-----	---------	-----

	er of ths.	Aggregate Age.			Average Age.		
YEARS.	Number of Deaths.	Years.	Mos.	Days.	Years.	Mos.	Days.
856	30	788	5	3	26	8	11
957	80	781		18	26		12
568	24	829	5	28	84	6	22
59	87	1,252	7	18	83	10	9
16 0	90	706	6	19	35	3	j 28
61	23	723	8	1 7	81	4	25
02	21	540	9	11	25	9	
63	47	1,456	11	28	80.	11	29
84	42	1,360		27	39	8	18
5	82	2,262	4	14	27	7	2
8	46	1,420	9.	17	80	10	15
7	83	654	4	26	19	9	28
8	24	692	1 7	8	28	10	9
9	85	1,214	4	18	84	8	11
0	52	1,491	1	7	27	3	28
1	53	1,610	11	15	80	4	22
ž	68	2,161	8	19	81	9	12
8	61 -	1,672	6	10	27	5	1
4	60	1,563	11	24	25	10	24
5	71	1,938	5	19	27	-8	19
6	83	922	1 8	17	27	11	16
7	43	1,248	11	23	28	11	5
8	47	1,151	9	97	24	6	9
9	40	1,206	l ë	20	80	2	ī
0	58	1,505	ğ	~~	28	4	24
1	38	1,156		. 18	80	5	2
26 years	1,118	82,227		29	28	11	14

Heart, Diseases of. The number of deaths from diseases of the heart in 1881 was 121, including 8 specified as hypertrophy; this was 17 more than in 1880, and the per cent. 0.64 more. The statistics of sex and parentage are:—

Males, 57; females, 64; total, 121; or 5.68 per cent. of all.

American parentage, 65; foreign parentage, 56.

In the twenty-six years, 1856-1881, the number of deaths from diseases of the heart was 1,703, or 4.71 per cent. of all; annual average, 65.

The particulars for each year are given in the following table :-

	SE	ix.	PARENTAGE.	Whole
	Males.	Females.	American. Foreign.	Number. Percentage
1856	8	13	16 5	21
1857	. 13	18	22 9	31
1858	. 13	23	28 8	36
1859	. 13	11	17 7	24
1860	. 15	16	21 10	31
1861	. 23	18	29 12	41
1862	. 26	23	33 16	49
1863	23	19	30 12	42
1864	26	32	44 14	58
1865	. 18	29	33 14	47
1866	19	21	27 13	40
1867	26	23	35 14	49
1868	. 25	27	34 18	52
1869	. 33	29	35 27	62
1870	28	31	33 26	59
1871	40	37	46 31	77
1872	. 43	49	63 29	92
1873	. 46	54	59 41	100
1874	. 46	58	62 42	104
1875	. 43	45		
1876	41	47	51 37	88
1877				9348
1878			nen agestallinger agences	83
1879				111
1880			too describing excession	
1881		**********		
		01		-

Hemorrhage. There were 12 deaths from this cause in 1881; 6 less than in the previous year.

Males, 7; females, 5; total, 12; or 0.56 per cent. of all. American parentage, 4; foreign parentage, 8

Of these deaths, 3 were hemorrhage from the lungs; 2 from the stomach; 2 from the bowels; and 3 were umbilical hemorrhage; the others were not specified.

Hernia. This was given as the cause of 3 deaths in 1881; 2 less than in 1880.

Males, 2; female, 1; total, 3; or 0.14 per cent. of all. American parentage, 1; foreign parentage, 2.

Hooping Cough. The number of deaths from this cause in 1881 was 40, or 1.88 per cent. of all; in 1880 there were only 11 deaths, and the rate 0.53 per cent. The decedents were 17 under one year old; 13 between one and two; 6 between two and five; and 4 between five and ten.

Males, 20; females, 20; total, 40; or 1.88 per cent. of all.

American parentage, 24; foreign parentage, 16.

For the twenty-six years, 1856-1881:--

Whole number, 437; annual average, 17.

Males, 190; females, 247; total, 437; or 1.21 per cent. of all.

American parentage, 189; foreign parentage, 248.

Hydrocephalus. This was reported as the cause of 32 deaths in 1881, or 3 more than in 1880; all the decedents were less than ten years old. The statistics of sex and parentage are as follows:—

Males, 16; females, 16; total, 32; or 1.50 per cent. of all. American parentage, 23; foreign parentage, 9.

For the twenty-six years, 1856-1881:—

Whole number, 794; annual average, 30.

Males, 430; females, 364; total, 794; or 2.20 per cent. of all.

American parentage, 405; foreign parentage, 389.

Insanity. The number of deaths from insanity in 1881 was 16; twice as many as in 1880.

Males, 5; females, 11; total, 16; or 0.75 per cent. of all.

American parentage, 12; foreign parentage, 4.

For the twenty-six years, 1856-1881, the figures are:—

Whole number, 272; annual average, 10.

Males, 127; females, 145; total, 272; or 0.75 per cent. of all. American parentage, 194; foreign parentage, 78.

Of the 272 decedents, 46.69 per cent. are males; 53.31 per cent. females; 71.32 per cent. of American, and 28.68 per cent. of foreign parentage.

A large majority of these deaths occur at Butler Hospital for the Insane, where the patients come, of course, from many different places.

Intemperance. The number of deaths from intemperance and delirium tremens, in 1881, was 14; 5 more than in 1880; and 0.66 per cent. of all from known causes.

Males, 7; females, 7; total, 14; or 0.66 per cent. of all. American parentage, 2; foreign parentage, 12.

For the twenty-five years, 1857–1881, the statistics are:

Whole number from known causes, 35,131.

From intemperance, 252; annual average, 10.

Males, 179; females, 73; total, 252; or 0.72 per cent. of all.

American parentage, 70; foreign parentage, 182.

Of the 252 decedents, 71.03 per cent. were males; and 72.22 per cent. were of foreign parentage.

Kidneys, Diseases of. There were 48 deaths from diseases of the kidneys, including 30 from "Bright's Disease," and 1 from "Addison's Disease"; the number of deaths in 1880 was 46. The statistics for 1881 are:—

Males, 25; females, 23; total, 48; or 2.25 per cent. of all.

American parentage, 23; foreign parentage, 25.

For the twenty-six years, 1856-1881:-

Whole number, 574; annual average, 22.

Males, 332; females, 242; total, 574; or 1.59 per cent. of all.

American parentage, 319; foreign parentage, 255.

Liver, Diseases of. The number of deaths in 1881 from disease of the liver was 23, or one more than in 1880.

Males, 14; females, 9; total, 23; or 1.03 per cent. of all. American parentage, 7; foreign parentage, 16.

The statistics for the twenty-six years, 1856-1881, are:

Whole number, 403; annual average, 15.

Males, 207; females, 196; total, 403; or 1.11 per cent. of all.

American parentage, 193; foreign parentage, 210.

Malformations. The number of deaths reported from malformations in 1881 was 15; four more than in 1880. Three cases were "spina bifida"; 7 malformation of the heart; 1 of the anus; 1 of the rectum; and in three cases it was not specified.

Males, 8; females, 7; total, 15; or 0.70 per cent. of all.

American parentage, 4; foreign parentage, 11.

For the twenty-six years, 1856-1881, the statistics are:

Males, 133; females, 78; total, 211; or 0.58 per cent. of all.

American parentage, 130; foreign parentage, 81.

Of the 211 decedents, 61.61 per cent. were of American parentage.

Marasmus. There were 34 decedents from this cause in 1881, of whom 27, or 79.41 per cent., were less than a year old.

Males, 20; females, 14; total, 34; or 1.60 per cent. of all. American parentage, 17; foreign parentage, 17.

For the twenty-six years, 1856-1881, the statistics are:—
Whole number, 687; annual average, 26.
Males, 368; females, 319; total, 687; or 19.00 per cent. of all.
American parentage, 346; foreign parentage, 341.

Measles. This disease, which has appeared in Providence very irregularly, prevailed to some extent in 1881, chiefly in the second quarter of the year; there were 25 deaths from it during the year, of which the particulars are as follows:—

Males, 11; females, 14; total, 25; or 1.17 per cent. of all.

American parentage, 10; foreign parentage, 15.

The following table shows the number of deaths from measles in each year since 1856:—

1877 9	187016	186316	1856 1
1878 50	1871 1	1864 7	1857 2
1879 0	1872 7	1865 7	185840
1880 3	1873 28	1866 9	1859 0
1881 25	1874 4	1867 0	18600
	1875 0	1868 4	1861 6
Total, 26 yrs285	1876 0	1869 6	1863 1

The statistics for the twenty-six years, 1856-1881, are:

Males, 111; females, 124; total, 235; or 0.65 per cent of all.

American parentage, 70; foreign parentage, 165.

Of the 235 decedents, 70.21 per cent. were of foreign parents

Meningitis, Cerebro-Spinal. There were 11 deaths from this can in 1881; 7 more than in 1880.

Males, 4; females, 7; total, 11; or 0.52 per cent. of all.

American parentage, 3; foreign parentage, 8.

The first cases of this disease in Providence were reported in 186 the number in each year since then is as follows:—

187112	1878
187210	1879
187339	1880
187410	1881
1875 6	-
1876 4	Total, 18 years18
1877 7	
	187210 187339 187410 18756 18764

Whole number from known causes in eighteen years, 29,122.

From cerebro-spinal meningitis, 123.

Males, 67; females, 56; total, 123; or 0.42 per cent of all.

American parentage, 56; foreign parentage, 67.

The number and percentage of deaths, in each division of age, according to sex and parentage, were as follows for the eighter years, 1864-1881, inclusive:—

AGES.	Under 1 Year.	1 and under 2.	2 and under 5.	5 and and and	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	Males.	Females.	American.	Foreign.	Torani.
No.	6	8	17	26	13	15	15	12	00	1	-	10	99	56	44	3100
PER CENT.	7.32	4.88	13.89	21,14	10.57	12.20	12.30	9.75	6.50	18.	18.	54.47	45.53	45.53	54.47	100-000

Of the 123 decedents, 66, or 57.72 per cent., were less than 15 years old.

Murder. There were 3 deaths in 1881 under this head; 2 more than in 1880. All the decedents were males; 1 of American, and 2 of foreign parentage; all were between 20 and 30 years of age.

Old Age. The number of deaths for which this cause was assigned was 73, the same as in 1880; fourteen of the decedents were more than 90 years old; also the same number as in the previous year:—

Males, 27; females, 46; total, 73; or 3.43 per cent. of all.

American parentage, 39; foreign parentage, 34.

For the twenty-six years, 1856-1881, the statistics are :--

Whole number, 1,400; annual average, 54.

Males, 474; females, 926; total, 1,400; or 3.87 per cent. of all. American parentage, 850; foreign parentage, 550.

Of the 1,400 decedents, 66.14 per cent. were females; and 60.71 per cent. were of American parentage.

Only a small proportion of those dying at an advanced age are reported under this head. Thus in 1881 there were 279 decedents who were more than 70 years old, 177 of American, and 102 of foreign parentage; this was 13.10 per cent. of all decedents from known causes during the year.

In the twenty-six years, 1856-1881, the number of decedents of more than 70 years of age was 4,028; 2,790 of American, and 1,238 of foreign parentage; 11.14 per cent. of all from known causes.

Paralysis. This is generally a symptom of disease rather than a distinct disease itself, and in many cases the cause of it might be given; but many deaths are reported every year under the head of "paralysis". In 1881 the number was 27, which was 3 less than in 1880.

Males, 9; females, 18; total, 27; or 1.27 per cent. of all American parentage, 12; foreign parentage, 15.

For the twenty-six years, 1856-1881, the figures are:—
Whole number, 465; annual average, 18.
Males, 239; females, 226; total, 465; or 1.29 per cent. of all.
American parentage, 339; foreign parentage, 126.

Pneumonia. The number of deaths from pneumonia in 1881 was 160; ten less than in 1880; from congestion of the lungs, 14; or 9 less than in 1880. They occurred in the different seasons of the year as follows:—

1st quarter	8d. quarter10
2d quarter48	4th quarter

Of the 174 decedents, 60, or 34.18 per cent., were less than ten years old; and 76, or 43.68 per cent., were over fifty years of age.

Males, 85; females, 89; total, 174; or 8.17 per cent. of all.

American parentage, 88; foreign parentage, 86.

The general statistics for the twenty-six years, 1856-1881, are as follows:—

Whole number, 2,683; annual average, 103.

Males, 1,371; females, 1,312; total, 2,683; or 7.42 per cent. of all.

American parentage, 1,341; foreign parentage, 1,342.

The following table gives the particulars as to the deaths from pneumonia for each of the twenty-six years, 1856-1881, giving the sex, parentage, season, age, whole number, and percentage of all deaths from known causes:—

PNEUWONIA AND Condestion of Lunas in Providence, Twenty-Six Years, 1856–1862.

Annal Section of Lunas in Providence of the Section of the Section of the Section Court of the Section Court of the Section of the Sectin DM' parentage; season; age; whole number, and percentage from known causes.

											1				١	I	١	ł			1					1
PREUMONIA AND CONG. OF LUNGS.	1856.	1867.	1866.	9281	1990.	1981	88	88	964.	1865.18	888	967. 134	888	1870.		1872.	1873.	1.1874.	1875.	.1876.		1877. 1878.	1879	1880.1	.188	Total
AGE: Under 1 year 1 and under 5 2 10 10 15 11 15 20 40 20	ស្តីជីក្សា (សក្ខេស DF) (សុខ្មា)	**************************************	85500 -4456000	<u> </u>	Notes and an an an an an an an an an an an an an	**************************************			84F4883H486F4	<u> </u>	Har : ***********************************		<u> </u>	8000::0008=	₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩₩	<u> </u>	######################################		4350-48FF88899	521 201 201 201 201 201 201 201 201 201 2	2020 : 867 5087 44	8420-44-80004	: : : : : : : : : : : : : : : : : : :	85000005485884-	888-84-748885eu	282222411F2222224
First quarter Second quarter Third quarter Fourth quarter	88°2	2208	200 ac	47.08	2478	2202	2508	8208	1212	23°8	8 8 x 5	2000	8828	2222	27788	8278	3300 B	82728	8453	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	8822	2822	2828	8573	5458	1,018 886 878 878
First 4 months Second 4 months Third 4 months	128	252	228	858	88.55	985	828	253	31 18 18	2 18	856	3 ∞ ∷	988	828	& 84 80 84 80	### ###	2382	887	55 10	87	88	583	881	884	288	35 45 138 138 138 138 138 138 138 138 138 138

Rheumatism. There were twelve deaths ascribed to this cause in 1881.

Males, 8; females, 4; total, 12; or 0.56 per cent. of all. American parentage, 5; foreign parentage, 7.

Scarlatina. For two years, 1879 and 1880, scarlatina prevailed as a severe epidemic, causing 252 deaths in 1879, and 243 in 1880; in the latter part of the year, however, it began to decline, and the number of deaths in 1881 was 45, or less than in any year since 1877; 36 of these deaths, or 80 per cent., were in the first half of the year. The decedents were 2 under one year old; 4 between one and two years; 24 between two and five; 14 between five and ten; and 1 between ten and fifteen.

Males, 27; females, 18; total, 45; or 2.11 per cent. of all.
American parentage, 17; foreign parentage, 28.

For the twenty-six years, 1856-1881, the general statistics are:

Whole number, 2,047; annual average, 79.

Males, 1,005; females, 1,042; total, 2,047; or 5.66 per cent. of all

American parentage, 839; foreign parentage, 1,208.

The following table gives the particulars of sex, parentage, and locality, for each of the twenty-seven years, 1855-1881:—

DEATHS FROM SCARLAITHA IN PROVIDENCE, TWENTY-SEVEN YBARS, 1855-1881.

I. Total	2,089	1,027	28.23	488 888 888 888 888 888 888 888 888 888	23.2 29.2 28.2 28.2 28.2 29.2 20.2 20.2 20.2 20.2 20.2 20.2 20
1881.	2.11	28	128	7004F0044H H04H	: 800 4000045
1880.	248 11.77	192	88.53	288885331-01-0	11-2-2528823
1879.	252 19 49	186	101	88890440884E	8-852522
1870, 1871, 1872, 1878, 1874, 1875, 1876, 1877, 1878.	2.89	22	23	9991995	
1877.	12 0.62	400	80 80		
1876.	85 1.87	16	7.5	400.40	4
1875.	88.4	\$3	83	990000000400c0	805-884588
1874.	268 13.66	140	F	8222244248	
1878.	182	£ 5	28.25	23.00.120.100.000	8 4 2 4 4 9 1 2 8
1872.	1.20	81	81		
1871.	1.66	, 100 m	ө М	0) 1-4-01-	H 0.4 H 00 01 00 00
1870.	2.80	28	82	5004 : 0 - · · · · ·	H466.06
1860	138	48	25 88	81.0184410148	\$c 8 c c C c c s
1867, 1868.	88	22	34	H48H48844414	ö ⊕ α α α α α α α α
1867.	58.		917		400
1846.	. S				
1864. 1865.	9.08	28	68	28 x 20 x 2 x 2 x 2 x 2 x 2 x 2 x 2 x 2 x	@ # B # C B #
	12.11	86	83		# 1 8 4 8 8 6 · · ·
1968.	85.55	808	116		
1969.	1.56		*:		
1961	84 5.	55	77		484 - 864 H
1860	1.17	10	2.0	808894884	&-800c0 · ·
1859.	6.18	22	153		
1867. 1858. 1859.	5.83	28	83	000000000000	D041401
	28 8 29 8	88	82	======================================	
1855. 1856.	14.00	558	25	91 01 05 91 11 11	875005
1855.	4.87	23	28		
SCARLATINA.	Whole number Percentage	SEX : Males Females	PARENTAGE: American Foreign	January. January. February March Jane June June Geptember. November.	Ward II.

Scrofula. There were 4 deaths from this cause in 1881; 3 more than in 1880, and 0.19 per cent. of all; all were females; 3 of American, and 1 of foreign parentage.

Skin, Diseases of. There were 3 deaths from diseases of the skin in 1881; this was 2 more than in 1880; all the decedents were females; 2 of American and 1 of foreign parentage.

Small-Pox. The city continues to escape this disease, which has prevailed extensively in many other large cities; and there has been no death from it since 1875. There was one case sent to the hospital at Field's Point in the summer of 1881; the patient recovered, and no other cases arose from it.

For the twenty-six years, 1856-1881, the statistics are:—
Males, 58; females, 21; total, 79; or 0.22 per cent. of all.
American parentage, 48; foreign parentage, 31.

The particulars for each year are as follows:-

SMALL-Pox.

•	S	EX.	PAREN	TAGE.	Whole	
	Males.	Females.	American.	Foreign.	Number.	Percentag
856	7	2	5	4	9	
858				· · · · · · · · · · · · · · · · · · ·		
			5			
860	 4	1	8	2	5	
861	· · · · · · · · · · · · · · · · · · ·					
862	2	2	1	8	4	a
863						
864	7		8	4	7	.
965	9		8		11	۵۵
						•••••
867	1		1		1	
868	1	· · · · · · · · · · · · · · · · · · ·	1	· • • • • • • • • • • • • • • • • • • •	1	
			1			
			9			
			8			
			· · · · · · · · · · · · · · · · · · ·			
876			•••••			
1877				· · · · · · · · · · · · · · · · · · ·	• • • • • • • • • • • • • • • • • • • •	
1878						
						• • • • • • • • • • • • • • • • • • • •
1880			· · · · · · · · · · · · · · · · · · ·			•••••
881			· · · · · · · · · · · · · · · · · · ·			

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Spine, Diseases of. The number of deaths reported from diseases of the spine, during 1881, was 6; twice as many as in the preceding year, and 0.28 per cent. of all.

Male, 1; females, 5.

American parentage, 5; foreign parentage, 1.

Stomach, Diseases of. There were 23 deaths reported in 1881 under this head, of which 19 were called "inflammation"; the whole number in 1880 was only 10. In 1881:—

Males, 8; females, 15; total, 23; or 1.08 per cent. of all.

American parentage, 6; foreign parentage, 17.

Suicide. There were 14 cases of suicide in 1881, the largest number that has ever been recorded in the twenty-six years, 1856–1881. The manner of each was as follows: "cutting the throat", 3; hanging, 2; drowning, 1; shooting, 1; by opium, 1; by laudanum, 2; by morphine, 2; by arsenic, 1; and 1 jumped from a third story window, while intoxicated.

Males, 12; females, 2; total, 14; or 0.66 per cent. of all.

American parentage, 8; foreign parentage, 6.

The statistics for the twenty-six years, 1856-1881, are:-

Males, 115; females, 46; total, 161; or 0.45 per cent. of all.

American parentage, 89; foreign parentage, 72.

Teething. The number of deaths from this cause in 1881 was 9, or 7 less than in 1880.

Males, 3; females, 6; total, 9; or 0.42 per cent. of all. American parentage, 3; foreign parentage, 6.

Tuberculosis. This was reported in 1881 as the cause of 25 deaths; 16 more than in 1880.

Males, 12; females, 13; total, 25; or 1.17 per cent. of all.

American parentage, 16; foreign parentage, 9.

If all the diseases of a similar nature were included under this

class, the percentage of it would exceed that of any other; the diseases have never been so classified in these reports. The heading "tuberculosis" was not specified separately until the year 1867, since when the deaths have been as follows:—

Whole number from known causes in fifteen years, 24,867.

From tuberculosis, 206; annual average, 14.

Males, 91; females, 115; total, 206; or 0.83 per cent. of all.

American parentage, 136; foreign parentage, 70.

Tumors. There were 6 deaths from tumors in 1881; the same number as in 1880; all the decedents were females of American parentage; 0.28 per cent. of all. Three were specified as "ovarian" tumors; 2 "abdominal"; and 1 "fibroid".

Unknown. There were 15 deaths in 1881 for which no cause was reported; 0.70 per cent. of all deaths during the year; the number was 2 less than in 1880, and the per cent. 0.12 less.

Males, 10; females, 5; total, 15; or 0.70 per cent, of all deaths.

American parentage, 9; foreign parentage, 6.

Ten of the decedents were less than a year old. The statistics of deaths from unknown causes for the twenty-six years, 1856-1881, are as follows:—

Whole number, 565; annual average, 22.

Males, 294; females, 271; total, 565; or 1.54 per cent. of all.

American parentage, 194; foreign parentage, 371.

Of the 565 decedents, 65.66 per cent. were of foreign parentage.

STILL-BORN.

The statistics in regard to the still-born children in Providence in 1881 were as follows:—

Number. The number of still-born in Providence in 1881 was 161; 40 more than in 1880, and 5.74 per cent. of the 2,803 children born during the year, or one in 17.41.

Sex. There were 101 males; 25 more than in 1880; and 60 females, or 15 more than in the preceding year.

Color. White, 147; colored, 14.

Ward VII18	Ward I25	Locality.
VIII18	II 4	
IX20	III13	
X20	IV11	
	V12	
Whole city 16	VI 9	

Season. January, 10; February, 18; March, 9; April, 19; May, 15; June, 11; July, 10; August, 13; September, 13; October, 12; November, 16; December, 15.

The number and percentage in each quarter of the year were :-

	Number.	Per cent.	Number.	Per cent.
lst quarter		22.98	3d quarter36	22.36
3d quarter	45	27.95	4th quarter48	96.71

Parentage. The following table shows the parentage of the still-born children in 1881; the first letter indicating the birth-place of the father; the second, that of the mother:—

A. and A76	E. and A 1	It. and I
I. and I40	8. and A 1	A. and E
E. and E	Af. and A 1	S. and E
G. and G 1	W. I. and A 1	A. and B. A
B. A. and B. A 2	A. and I 8	E. and S
P. and P 3	E. and I 1	I. and S
It. and It 1	S. and I 1	-
I. and A 3	P. and I 1	Totallii

A.—American; I.—Irish; E.—English; G.—German; B. A.—British American; P.—Portuguese; It.—Italian; Af.—African; W. I.—West Indian.

The number of children born in Providence in 1881 was 2,803; of these, 161, or 5.74 per cent., were still-born. Of the 2,803 children born, 1,428, or 50.95 per cent., had American mothers; 809, or 28.86 per cent., had Irish mothers; and 566, or 20.19 per cent., had mothers who were natives of other foreign countries.

Of the 161 still-born children, 83, or 51.55 per cent., had American mothers; 52, or 32.30 per cent., had Irish mothers; and 26, or 16.15 per cent., had mothers of other foreign countries.

The proportion of still-born children to the whole number of children borne by mothers of each class was as follows: of the 1,428 children of American mothers, 83, or 5.81 per cent., were still-bon. that is one in 17.20; of the 809 children of Irish mothers, 52 were still-born; or 6.43 per cent., or one in 15.56; of the 566 children of other foreign mothers, 26 were still-born; or 4.59 per cent., or one in 21.78. The proportion in 1881 was largest among the Irish mothers. The whole number of children borne by American mothers in 1881 was 96 more than in 1880; and the proportion of these to all the children born was 0.25 per cent. more; of still-born children of American mothers in 1881 the number was larger, but the proportion less, than in 1880. Of children of Irish mothers in 1881, the number was slightly larger, but the proportion less than in 1880; of still-born of Irish mothers the number was 16 more, and the proportion 2.55 per cent, more than in the preceding year. other foreign mothers, the number and proportion were greater in 1881 than in 1880; of still-born children in this class, the number and proportion were also larger than in the preceding year.

For the twenty-six years, 1856-1881, the statistics on this subject are as follows:—

Whole number of children born, 51,852.

Number of still-born, 3,009, or 5.80 per cent. of all; or one in 17.23. Of these 3,009 still-born children, 2,815 were white, and 194 colored; 1,742, or 57.89 per cent., were males; and 1,267, or 42.11 per cent., females.

Reckoned according to the fathers' birth-places, 1,309, or 43.50 per cent., were American; 1,285, or 42.71 per cent., were Irish; and 408, or 13.56 per cent., were children of other foreign fathers; of 7, the parentage was not stated.

The following statement shows the whole number of children borne by mothers of different nativities, during the twenty-six years, 1856–1881, inclusive; also the number of still-born children to mothers of each class; and the percentage of the latter to the whole number of children:—

	Whole number	Still-born	Percentage	Or 1
	of children.	children.	of still-born.	in
American mothers	24,463	1,307	5.84	18.79
Irish mothers	19.853	1,360	6.85	14.60
English, Scotch or Welsh mot	hers 3,851	188	4.88	20.48
German mothers	1,048	80	2.88	84.48
British American mothers	1,754	72	4.10	24 . 36
Other foreign mothers	888	52	5.88	17.08

GENERAL SUMMARY.

The following recapitulatory table will give the general results of registration in Providence during the 27 years, 1855 to 1881 inclusive:—

TABLE XXI. BIRTHS, MARRIAGES, AND DEATHS.

YEAR.	Population.	Births.	To population, one birth in	Marriages.	Of population, one person married in	Deaths.	Of population, one death ta
1855	47,785	1,000	29.86	636	37,56	991	48.22
1856	48,727	1,675	29.09	656	37.14	1,065	45.73
1857	49,457	1,688	29.30	614	40.27	935	33.40
1858	49,457	1,724	28.68	560	44.15	1,017	2.0
1859	50,100	1,593	31.45	616	40 00	809	63.71
1860	50,666	1,648	30.74	633	40.02	1,001	50.61
1861	50,666	1,725	29.37	547	46.31	1,051	48.21
1862	52,238	1,529	34.16	547	47.75	914	37.35
1863	53,024	1,307	40.56	610	43.46	1,214	45.57
1864	53,810	1,344	40.04	739	36.40	1,281	42.00
1865	54,595	1,298	42.06	709	38.50	1,211	- 62.08
1866	55,700	1,632	34.13	812	34.30	1,036	68.77
1867	56,824	1,624	34.99	855	33.23	900	-59.19
1868	64,138	1,866	34.37	872	36.77	1,110	57.76
1869	66,522	1,819	36,57	930	35.76	1,256	52.9
1870	68,904	1,889	36.48	959	35.93	1,263	54.50
1871	72,011	1,960	36.74	943	38.13	1,254	50.40
1872	75,118	2,206	34.05	1,084	34.63	1,603	45.8
1873	78,225	2,128	36.76	1,150	34.01	1,719	45.50
1874	96,452	2,780	34.69	1,176	41.01	1,987	45.5
1875	100,675	2,664	37.79	1,079	46.65	1,915	52.3
1876	101,511	2,631	38.58	949	53.48	1,865	54.4
1877	101,000	2,543	39.72	953	52.99	1,908	52.11
1878	102,286	2,627	38.94	1,016	50.34	1,989	STA
1879	103,572	2,522	41.07	1,071	48.35	2,026	51.11
1880	104,857	2,627	39.92	1,234	42.49	2,080	50.41
1881	107,500	2,803	38.35	1,202	44.71	2,145	50.12
Average	70,875	1,980	35.79	857	41.35	1.397	50.73

BIRTHS AND DEATHS BY PARENTAGE.

The following recapitulatory table shows the number of births and deaths, by parentage, in Providence, in each of the twenty-six years, from 1856-1881 inclusive. In the three years, 1863, 1864, and 1865, there was an excess of deaths over the births, in the population of American parentage:—

TABLE XXII.

ļ	AMERIC	AN PARI	INTAGE.	Forei	ON PARE	NTAGE.
YEAR.	Deaths.	Births.	Excess of Births.	Deaths.	Births.	Excess of Births.
1856	541	759	218	524	916	392
1857	5 23	701	178	402	987	585
1856	512	771	259	505	953	448
18 59	479	659	180	420	934	514
1880	519	731	219	489	917	428
1861	538	740	202	518	965	479
1962	460	663	203	454	866	412
1863	597	528	69	617	779	162
1864	649	579	—70	632	765	133
1865	631	546	85	580	752	172
1866	513	726	218	523	906	383
1867	528	725	197	482	899	467
1868	596	882	286	514	984	470
1869	627	926	299	629	893	264
1870	642	890	248	621	999	878
1871	643	980	837	611	980	369
1872	832	1,035	203	771	1,171	400
1873	862	1,003	141	857	1,125	268
1874	885	1,217	832	1,102	1,563	461
1875	914	1,180	286	1,001	1,484	483
1876	875	1,169	294	990	1,462	472
1877	908	1,238	830	1,030	1,305	275
1878	889	1,215	826	1,100	1,413	312
1879	941	1,228	287	1,085	1,294	209
1880	920	1,252	832	1,160	1,875	215
1881	999	1.365	366	1,146	1,488	293
26 years	18,016	23,708	5,692	18,708	28,144	9,486

-		
		•

Vital Statistics, Providence, R. I.


1881.

CITY REGISTRAR'S
TWENTY-SEVENTH ANNUAL REPORT.

1883 | PITT DOULARRATE No. 14.

BIRTHS, MARRIAGES, AND DEATHS





TWENTY-EIGHTH ANNUAL RI

UPON THE

BIRTHS, MARRIAGES, AND

THE NEW YORK PUBLIC LIBRARY.

CITY OF ASTOR, LENOX AND TILDEN FOUNDATIONS.

IN THE

PROVIDENCE:

FOR THE YEAR

1882,

Br EDWIN M. SNOW, M. D.,

SUPERINTENDENT OF HEALTH AND CITY REGISTRAN.



PROVIDENCE PRESS COMPANY, PRINTERS TO THE CITY.



THE CITY OF PROVIDENCE.


JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved April 23, 1883.]

RESOLVED. That the twenty-eighth annual report of the City Registration for the year ending December 31, 1882, be received, and that six hundare copies thereof, and of the communication accompanying the same. I printed in pamphlet form for the use of the City Council, under direction of the joint committee on printing.

A true copy: witness,

HENRY V. A. JOSLIN, CITY CLERK.



INTRODUCTION.

CITY REGISTRAR'S OFFICE, PROVIDENCE, April 16, 1883.

To the Honorable the City Council:

The statistics of births, marriages, and deaths in the city of Providence, during the year ending December 31, 1882, are contained in the enclosed twenty-eighth annual report upon this subject.

During the year 1882, the number of births in the city was 2,788, or 15 less than in 1881; the number of persons married was 2,344, or 60 less; and of deaths 2,242, or 97 more.

The population of the city, in 1880, was 104,857, and in January of the present year, it was 116,755. According to the rate of increase between these two periods, the population of the city, in the middle of the year 1882, was 114,377. We find, then, the proportion of births, marriages and deaths to population, in 1882, to be:

Births, 2,788, or one in 41.02; or 24.38 in each 1,000.

Persons married, 2,344, or one in 48.80; or 20.49 in each 1,000.

Deaths, 2,242, or one in 51.02; or 19.60 in each 1,000.

The report herewith presented contains the usual tables of previous reports, with additional tables illustrating the characteristics of some important diseases, especially of typhoid fever. It also contains the statistics of the population of the city as obtained by the census of the first of January, 1883.

EDWIN M. SNOW, M. D.,

Superintendent of Health and City Registrar.

CONTENTS.

Віятня	PAGE. 1-20
Wards, months, sex and color, (Table 1.)	. 1
Births in connection with season, sex and locality, (Table 11.) Proportion of the sexes born at different seasons	. 2-3 . 4
Number and proportion of the sexes, 29 years	. 5
Color:—Births, marriages and deaths in 27 years	. 7
Parentage of children born in each ward, (Table III.)	. 9-10
Parentage:-Mixture of different nations, (Table IV.)	ii
Average number of children to each mother, 28 years, (Table V.)	13
Color:—Number and proportion of the seven, 2/ years. Color:—Births, marriagos and deaths in 27 years. Parentage of children born in each ward, (Table IN.). Number and percentage of American and of foreign children, 27 years. Parentage:—Mixture of different nations, (Table IV.). Number of ist, 21, etc., children in 1882, and 1835–1881, (Table V.). Average number of children to each mother, 29 years, (Table VI.). Number of the child, with age of the mother, (Table VII.). Plurality births, attastica for 1882.	14-16
Plurality births, statistics for 1882 Plurality births, 28 years' summary	17-20
Marriages	20-32
MARRIAGES. Marriages at different seasons, 23 years, (Table VIII.). Birthplaces of parties married in 1882, (Table IX.). Bride and groom born in same country, etc. Proportions of American, foreign, and mixed marriages, 28 years	21-22
Bride and groom born in same country, etc	23-24
Proportions of American, foreign, and mixed marriages, 28 years	. 25 26
Marriages of colored persons. Age of persons married in 1882, (Table X.). Average age of persons married, If years. (Table). Number of times married, (Table XI.) Denominational.	27
Average age of persons married, 27 years. (Table)	28 29
Nativity and residence	80-31
Marriage and education; certificate signed with a mark	31-32
Deaths to population in 1882; proportion in each class	83-108 28
Deaths by sex, condition, color, nativity, parentage, and locality,	
(Table XII.)	34
Sex. condition, color, nativity, parentage, and locality of decedents.	
Aggregate and average age by parentage, 1882. (Table XIII.)	87-39 40
Aggregate and average age by parentage, 1882, (Table XIII.)	41 48
Difference between average age of American and foreign decedents,	#1_#0
Percentages at different ages by parentage, 1862. (Table XVI.)	47
Deaths by age and parentage in each month and quarter, 1882, (Table	48-49
XVIII.)	49-51
Causes of death in connection with season, (Table XX.)	50-6 0
NOTES ON THE CAUSES OF DEATH IN 1882	31-103
Cholera Infantum; decedents by sex and parentage, 27 years, (Table)	66 67
Consumption; sex and parentage	-
Consumption in Providence, 27 years, (Table)	69-70 71-79
Diarrhoa and dysentery; recapitulation for 42 years	78-75
Consumption in Providence, 27 years, (Table). Croup; statistics for 1882, and 27 years, (Table). Diarrhosa and dysentery; recapitulation for 42 years. Diphtheria; statistics for 1882, and 25 years, (Table). Fever, Typhoid, 1882, and 27 years, (Table). Fever, Typhoid, 1882, and 27 years, (Table). Fever, Typhoid, Epidemic in 1882.	75-77 78-83
Fever, Typhoid, Epidemic in 1882	83-84
Meningitis. Cerebro spinal; deaths at each age. 19 years	89-90
Pneumonia; statistics for 1882 and 27 years, (Table)	93-94
Small-pox; recapitulatory table, 27 years	86
Fever, Typhoid, Epidemic in 1882. Heart, discusses of; decedents by sex and parentage, 27 years, (Table). Meningitis. Cerchro spinal; deaths at each age, 18 years. Pneumonia; statistics for 1882 and 27 years, (Table). Scarlatina; statistics for 1882 and 28 years, (Table). Small-pox; recapitulatory table, 27 years. Unknown causes. Still born children in 1882. Still born children. 1855–1882.	100 102
Still-born children, 1856–1882	103
GENERAL SUMMARY:-	
Births, Marriages and Deaths; numbers and proportions, 28 years, (Table XXI.)	104
(Table XXI.) Births and deaths by parentage; excess of births, 27 years, (Table XXII.)	105
Census of Providence, 1880, sex, nativity, parentage, and color	106
Census of Providence, 1883, houses, tenements, familles, sex and total population	106
E - 5	

CITY REGISTRAR'S REPORT.

1882.

BIRTHS.

There were 2,788 children born in the city of Providence, during. the year 1882.

In the following table we find the number of children born in each ward of the city in each month and quarter of the year; also, for the whole city, the number of each sex, and the number and sex of the colored children born in each month and quarter:—

TABLE I. BIRTHS, 1882. Wards, Months, Sex and Color.

MONTHS.	WARDS.									Whole City.		Colored children in- cluded in the preceding.				
	1.	п.	m.	ıv.	v.	VI.	VII.	VIII	ıx.	x.	Males,	Females.	Total.	М.	F.	Total.
January February Mareh	36 45 34	3 1 13	18 20 23	14 11 13	13 23 14	6 10 9	28 22 24	25 24 27	43 35 47			105	226	3 3	3 3	6
Ist Quarter	115	17	61	38	50	25	69	76	125	119	357	338	695	9	7	16
April May June	43 29 25	8 7 1	23 15 25	8 18 8	17 18 19	12 9 8	19 19 20	27 22 33	36 33 40	53 53 45	107	121 116 102		6 9 4	6 5 2	19 14 6
2d Quarter	97	16	63	34	54	29	58	82	109	151	354	339	693	19	13	32
July August September	26 25 28	3 5	17 24 24	11 16 6	19 18 25	12 10 7	18 16 30	14 29 21	38 36 27	48 46 60	110		223	5 1	8 2 1	107
14 Quarter	79	12	65	33	62	29	64	64	101	154	354	309	663	8	11	19
October November December	35 28 17	7 1 4	25 29 23	5 15 14	10 14 21	12 12 11	21 19 25	26 21 27	51 35 52	58 66 53	113	123 127 114	250 240 247	3 2 3	6 8 7	10 10
tth Quarter	80	12	77	34	45	35	65	74	138	177	873	364	737	8	21	29
Whole Year	371	57	266	139	211	118	256	296	478	601	1,438	1,350	2.788	44	59	96

The number of births, 2,788, was 15 less than in the year 1881; but with that exception was larger than in any previous year. In the first ward the number of births in 1882 was 17 more than in 1881; in the fourth ward, 11 more; in the seventh ward, 13 more; in the ninth ward, 49 more; in the remaining six wards the number of births was 105 less than in the previous year. A census of the city was taken in January, 1883; an estimate, based on the figures of this census, gives a population of 114,377 for the middle of the year 1882. The ratio of births, therefore, for the year 1882, was one in 41.02 of the population, or 24.38 births to each thousand of the inhabitants.

Season. In the first four months of the year 1882, and in December, the number of births was larger than in the previous year; in each of the remaining months the number in 1882 was less than in 1881. In the first quarter of 1882 the number was 61 more than in the corresponding season of 1881; in the second quarter it was 37 less; in the third quarter, 43 less; in the fourth quarter, 4 more. In the first half of the year 1882 the number of births was 1,388, or 24 more than in the first half of 1881; in the last half of the year the number was 1,400, or 39 less than in 1881.

The following figures show the aggregate number of births for each quarter for the twenty-six years, 1856-1881, inclusive; the number for each quarter of 1882, and the corresponding numbers for the twenty-seven years, 1856-1882:—

	Jan. March.	April. June.	First 6 mos.	July. Sept.	Oct. Dec.	Second 6 mos.	Whole Year.
Twenty-six years.	12,564.	12,918	25,502	12,905	13,445	26,350	51,852
1882	695.	693	1,888	668	787	1,400	2,788
Twenty-seven year	rs13,279.	18,611	26,890	13,568.	14,182.	27,750	54,610

For the period of twenty-eight years, 1855-1882, inclusive, the aggregate monthly totals of births are as follows; the daily average, also, is given for each month and for the whole period:—

Births in	January. F	ebruary.	March.	April.	May.	June.
28 years, 1855-1882	4,419	4,814	4,903	4,381	4,645	4,948
Daily average	5.09	5.45	5.65	5.22	5.35	5.89
Births in	July.	August.	Sept.	Oct.	Nov.	Dec.
28 years, 1855-1882	4,609	. 4,711	4,663	4,891	4,763	4,993
Daily average	5.81	5.43	5.55	5.63	5.67	5.75
Whole number o	f birthe in 28 3	ears, 56,24	0; daily av	erage, 5.5	0.	

These figures have for a number of years shown the same relative position of the months as to the number of births, namely, that the largest aggregate number of births in any month is in December; the smallest, in February; the highest daily average is in June; the lowest, in January. There were six months in which the daily average was above the general average of the whole period, namely: March, June, September, October, November and December; in the other months the daily average was less than the general average.

Sex and Locality. The number of children of each sex, born in each ward of the city, in each quarter of the year, and in the whole year, is shown by the following table:—

TABLE II. BIRTHS, 1882. Wards, Sex, and Season.

	1st Quarter.			2d	2d Quarter.			3d Quarter.			4th Quarter.			Whole Year.		
WARDS.	M.	F.	Total.	м.	F.	Total.	м.	F.	Total.	М.	F.	Total	м.	F.	Total.	
L	63	52	115	43	54	97	37	42	79	41	39	80	184	187	871	
П	8	9	17	12	4	16	9	-3	12	3	9	12	32	25	57	
III	31	30	61	35	28	63	35	30	65	41	36	77	142	124	200	
IV	19	19	38	14	20	34	16	17	33	13	21	34	62	77	139	
V	24	26	50	28	26	54	34	28	62	24	21	4.5	110	101	211	
VI	16	9	25	18	11	29	17	12	29	19	16	35	70	48	118	
VII	39	30	69	22	36	58	37	27	64	27	38	65	125	131	250	
vIII	42	31	76	43	39	82	31	33	64	37	37	74	153	143	296	
XX	57	68	125	59	50	109	58	43	101	84	54	138	258	2.5	473	
X	58	61	119	-80	71	151	80	74	154	84	93	177	302	200	601	
Whole City	357	338	695	354	339	693	354	309	663	373	364	737	1,438	1,350	2,788	

Proportions of the Sexes. In the months of March, May, August and November, there were more female than male children born; in the first, fourth and seventh wards, the number of female children exceeded that of the male children; but of the whole number, 2.788. 1,438 were males and 1,350, females; 88 more males than females; a proportion of 106.5 boys to 100 girls; or 51.58 boys, 48.42 girls, in each 100 children born.

The number and proportion of each sex in each quarter of the year, were as follows:—

Males. Females. Males. Females. Males. Females.

Jan.-March....357 and 338, or 105.6 to 100, or 51.37 to 48.63 in each 100 born.

April-June....354 and 339, or 104.4 to 100, or 51.08 to 48.92 in each 100 born.

July-Sept.....351 and 309, or 114.6 to 100, or 53.39 to 46.61 in each 100 born.

Oct.-Dec.....373 and 364, or 102.5 to 100, or 50.61 to 49.39 in each 100 barn.

Whole year...1,438 and 1,350, or 106.5 to 100, or 51.58 to 46.42 in each 100 born.

The proportion of males is much larger in the third quarter than in either of the others; the excess of male children in 1882 was rather less than in the two preceding years. The following figures show the same statistics for the twenty-seven years 1856–1882. The whole number of children born in this time was 54,640:—

Males. Females. Males. Females. Males. Females.

Jan.-March.....6,879 and 6,400, or 107.5 to 100, or 51.80 and 48.20 in each 100 born.

April-June.....7,003 and 6,608, or 106.0 to 100, or 51.45 and 48.55 in each 100 born.

July-Sept......6,909 and 6,659, or 103.8 to 100, or 50.92 and 49.08 in each 100 born.

Oct.-Dec......7,322 and 6,860, or 106.7 to 100, or 51.63 and 48.37 in each 100 born.

Whole period.28,113 and 20,527, or 106.0 to 100, or 51.45 and 48.55 in each 100 born.

In the twenty-nine years 1854-1882 inclusive, the number of children born in the city was 57,609; 29,631 males and 27,978 females; the number and proportion in each year were as follows:—

```
Males.
                  Females. Males. Females. Males. Females.
1854..... 711 and
                     658; or 108.0 to 100; or 51.91 and 48.09 in each 100 born.
1855..... 807 and
                     798; or 101.7 to 100; or 50.41 and 49.56 in each 100 born.
1656..... 891 and
                     784; or 113.6 to 100; or 53.19 and 46.81 in each 100 born.
1857..... 872 and
                     816; or 106.8 to 100; or 51.66 and 48.34 in each 100 born.
1858..... 890 and
                     834; or 106.7 to 100; or 51.62 and 48.38 in each 100 born.
          825 and
                     768; or 107.4 to 100; or 51.79 and 48.21 in each 100 born.
1860..... 809 and
                     839; or 96.4 to 100; or 49.09 and 50.91 in each 100 born.
1861 ..... 936 and
                     789; or 118.6 to 100; or 54.26 and 45.74 in each 100 born.
                     738; or 107.2 to 100; or 51.78 and 48.27 in each 100 born.
1862..... 791 and
                     618; or 111.4 to 100; or 52.71 and 47.29 in each 100 born.
1863..... 689 and
1864..... 663 and
                     681; or 97.8 to 100; or 49.33 and 50.67 in each 100 born.
1865 ..... 691 and
                     607; or 113.8 to 100; or 53.24 and 46.76 in each 100 born.
1866..... 849 and
                     783; or 108.4 to 100; or 52.02 and 47.98 in each 100 born.
1867..... 830 and
                     794; or 104.5 to 100; or 51.11 and 48.89 in each 100 born.
1868..... 1944 and
                     922; or 102.4 to 100; or 50.59 and 49.41 in each 100 born.
1869..... 945 and
                     874; or 108.1 to 100; or 51.95 and 48.05 in each 100 born.
1870 ..... 967 and
                     722; or 104.8 to 100; or 51.19 and 48.81 in each 100 born.
1871...... 956 and 1,004; or 95.6 to 100; or 48.78 and 51.22 in each 100 born.
1872..... 1,079 and 1,127; or 95.7 to 100; or 48.91 and 51.09 in each 100 born.
1873..... 1,109 and 1,019; or 108.8 to 100; or 52.11 and 47.89 in each 100 born.
1874..... 1,408 and 1,372; or 102.6 to 100; or 50.45 and 49.85 in each 100 born.
1875..... 1,390 and 1,274; or 109.1 to 100; or 52.18 and 47.82 in each 100 born.
1876..... 1,359 and 1,272; or 106.8 to 100; or 51.65 and 48.35 in each 100 born.
1877..... 1,302 and 1,241; or 104.9 to 100; or 51.20 and 48.80 in each 100 born.
1878..... 1,358 and 1,289; or 107.0 to 100; or $1.69 and 48.31 in each 100 born.
1879..... 1.297 and 1,225; or 105.9 to 100; or 51 43 and 48.57 in each 106 born.
1880...... 1,363 and 1,264; or 107.8 to 100; or 51.88 and 48.12 in each 100 born.
1881..... 1,462 and 1,341; or 109.0 to 100; or 52.16 and 47.84 in each 100 born.
1883..... 1,438 and 1,850; or 106.5 to 100; or 51.58 and 48.42 in each 100 born.
29 years..29.631 and 27,978; or 105.9 to 100; or 51.43 and 48.57 in each 100 born.
```

Color. There were 96 colored children born in Providence in 1882; this was 25 less than in the preceding year. Estimating the colored inhabitants to bear the same proportion to the total population as at the time of the census in 1880, the number of colored persons in the city in 1882 was 3,976. The ratio of births in this class, therefore, was one in 41.42 of the population, or 24.14 births to each thousand of the colored population. The ratio of all births during the year to the total population was one in 41.02, or 24.38 in each

The births, marriages and deaths among the colored population of Providence in the twenty-seven years, 1856–1882, inclusive, were as follows:—

Twenty-six years, 1856-18812	2,015 births 989 marriages2,1	02 deaths.
1882	96 births 53 marriages	93 deaths.
Twenty-seven years, 1856-1882 2	2,1 births1,042 marriages2,1	— 197 deaths.

The proportion of deaths to births among this class in 1882 was 99.0 to 100; for the whole period the proportion was 104.1 deaths to 100 births.

The number of births, marriages and deaths among the colored people of Providence, in each year since 1856, is as follows:—

1856 50 births 14 marriages	. 48 deaths.
1857 20 marriages	
1856 51 births 17 marriages	
1859 9 marriages	
1860 15 marriages	
1861 15 marriages	
1862 61 births 12 marriages	
19-3 32 births 55 marriages	
1864 38 births 12 marriages	
1865 46 births 23 marriages	
1896 86 marriages	
1867	
1868 90 births 58 marriages	
1869 65 births 60 marriages	
1879 104 births 52 marriages	
1871 91 births 46 marriages	
1872 99 births 54 marriages	
1873 49 marriages	
1874 109 births 60 marriages	104 deaths.
1975 48 marriages	111 deaths
1876 108 births 38 marriages	110 doothe
1877 104 births 46 marriages	105 douths
1878 104 births 54 marriages	106 deaths.
1879 43 marriages	110 deaths
1880 57 marriages 58	of deaths.
1881 121 births 61 marriages	
1882 96 births 53 marriages	• Avi ucailla.
—— — — — — — — — — — — — — — — — — — —	. ~ uestins.
97 years 9 111 bloths 1 040 manufactor	0.107.4004
27 years2,111 births1,042 marriages	z,iri ueatds.

PARENTAGE.

The statistics as to the parentage of the children born in the city in 1882 are given in the following tables, and show how many different nationalities are represented in our population and the relative proportions of the different classes.

Table III shows the number of children, according to parentage, born in each ward, and the number and proportion of each class in the whole city. Where the parents were of different foreign countries the parentage is reckoned according to the birth-place of the father:—

TABLE III. BIRTHS, 1882. Wards and Parentage.

				1	WAE	RDS.					Clty.	he 100
PARENTAGE.	ı.	11.	111.	ıv.	v.	VI.	VII.	vin	ıx.	x.	Whole City	In each
American	155	42	90	66	66	83	120	153	231	140	1,146	41.1
Irish	81	2	65	17	57	9	55	56	92	179	618	21.9
English, Scotch and Welsh	21	1	18	10	19	5	10	26	32	106	243	8.7
German	8	1	5	2	12	2	8	1	6	15	55	1.8
British American	24	9	11	4	18	1	7	10	16	23	116	4.1
l'ortuguese			30						••••		80	1.0
Other Foreign	17		! 	18	8	8	11	••••	20	11	83	2.9
Total Foreign	151	6	124	51	109	20	86	98	166	314	1,140	40.9
American father and Foreign mother	32	5	17	10	16	9	20	22	40	58	224	8.0
Foreign father and American mother	88	4	85	12	20	6	80	28	86	74	278	9.9
Ward totals	371	57	266	139	211	118	256	296	478	601	2,788	100.0

In the Third, Fifth and Tenth wards the number of children of purely foreign parentage was larger than that of American parentage; in the whole city the number of American parentage was six more than that of foreign parentage. The "other foreign" men-

tioned in this table were as follows: in the First ward, 10 of Russian parentage; 4 of Swedish, and 1 each of Austrian, Bohemian and French; in the Fourth ward, 15 of Italian parentage, and one each Russian, Swiss and Swedish; in the Fifth, 3 Swedish; in the Sixth, 1 each Swiss, Austrian and Italian; in the Seventh, 10 Italian and 1 Swedish; in the Ninth, 11 Swedish, 3 each Norwegian and French, 2 Swiss and 1 Danish; in the Tenth, 6 Swedish, 3 Italian and 1 each Norwegian and South American. In the whole city, there were 11 of Russian parentage; 29 Italian; 26 Swedish; 4 Norwegian; 4 French; 4 Swiss; 2 Austrian; 1 Bohemian; 1 Danish; 1 South American; total, 83.

The following table gives the number and percentage of children of American, foreign and mixed parentage, in each of the twenty-seven years, 1856--1882; and the number and percentage for the whole time:—

			Amer. father.	For. mother.
,	American.	Foreign.	For. mother.	Amer. father.
Year. Nu	mber. Per cent. Nu	mber. Per cent. N	umber. Per cent	. Number. Per ct
1856	70542.09	85751.17	. 558.22	583.5
1857	65088.51	98755.51	. 518.02	502.9
1858	71241.80	886 51.39	. 583.36	683.9
18 59	59937.60	88755.69	. 598.70	48 3.0
1860	66140.11	85752.00	. 694.19	613.7
1861	65437.91	91653.10	. 864.99	694.0
1862	58988.53	79952.26	. 744.84	674.8
1863	46785.73	71154.40	. 624.74	675.1
1864	52038.69	69952.00	. 604.47	654.8
1865	47636.67	68452.70	. 705.89	685.9
1966	656 40.20	82150.30	. 704.29	855.2
1867	64539.72	79148.70	. 804.98	1086.6
1868	77941.75	80846.51	. 1035.52	1166.9
i8 69	82245.19	77642.16	. 1045.72	1176.4
870	80042.85	86245.64	. 904.76	1377.9
871	87944.85	82842.24	. 101	1527.7
879	90741.12	1,00245.42	. 1285.80	1697.6
1878	86940.84	94144.22	. 1846.29	1848.6
1874	.,06888.42	1.826 47.70	. 1495.36	2878.5
875	99987.50	1,25647.15	. 1816.79	2288.5
1876	1,00788.27	.28046.75	. 1636.16	2828.8
1877	,05441.45	1,06741.98	. 1847.23	2389.8
878	,03589.40	1,17444.69	. 1806.85	2389.0
879	,02940.80	1,08541.04	. 1997.89	25910.2
1880	,04839.90	1,09141.53	. 2047.77	28410.8
1881	,14140.71	1,15141.06	. 2247.99	28710.9
863	,14641.10	1,14040.90	. 2248.03	2789.9
_		_		

Of children of American parentage, the number is greater than in any previous year; and the percentage greater than in any year since 1877. Of children of foreign parentage, the number was 11 less than in 1881, and the percentage less than ever before; of children of American fathers and foreign mothers, the number is the same as in the preceding year, but the percentage, slightly larger; and larger than in any previous year; of children of foreign fathers and American mothers, the number and percentage are less than for several years previous.

The percentage of children born in Providence of parents of each nation, is as follows, for each of the twenty-seven years, 1856-1882, inclusive:—

		-		Other.
			German. American. Portu	•
			Per cent. Per cent. Per	
			2.27	
185783.51	44.32	5.92 .		1,54
185841.80				
185987.60	45.58	5.84	3.26	1.5
188040.11	40.96	6.55	2.85	1.66
186187.91	42.61	6.03	2.49	1.97
186288.52	41.58	6.02	8.07	1.6
18 6385 .78	48.06	6.35	2.98	1.99
186 4	41.86	6.55	2.75	1.3
186586.67	42.99	5.98	1.85	1.9
i:::::::::::::::::::::::::::::::::::::	89.52	6.25		23
186739.72	87.81	5.66	8.09	2.2
1868 41.75	36.01	6.48	1.881.07	1.0
186945.19	84.14	4.01	1.811.82	1.3
187042. 85	84.68	6.19	1.96 1.91	9
187144.85	82.25	4.54	1.632.09	1.7
187241.12	83.46	5.67	9.779.31	2.3
187340.84	30.50	7.19	1.882.49	2.10
1874	32.68	8.18	2.84	231.19
87587.50	81.58	7.69	2.852.67	981.4
87638.27	81.05	7.75	2.592.891.	921.2
87741.45	28.08	6.98	1.962.561.	101.3
87889.40	29.65	6.74	2.55 2.86	181.7
			1.98	
			1.67	
			2.088.001.	
188341.10				

The British American were included in the *cother foreign" until 1868, and the Portuguese until 1874. The highest percentage of American parentage was 45.19 in 1869; the lowest, 35.73 in 1863; of Irish parentage the highest percentage was 45.58 in 1859; the lowest, 21.99 in 1882; the percentage of this class has steadily decreased for the past five years.

The following table shows more exactly the parentage of the children born in 1882, by giving the birth-places of both parents:—

	TABLE	IV.	Віктня, 1882.	Parentage.
--	-------	-----	---------------	------------

	ВІ	s.	according birth-					
BIRTH-PLACES OF FATHERS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands	Sountries.	Whole number accort to the father's birth place.
United States	1,146	117	53	6	40	ļ	. 8	1,370
Ireland	135	- 564	85	1	12		1	748
England, Scotland and Wales	70	87	182		23		2	813
Germany	18	5	8	43			4	78
British America	85	13	11	1	91			151
Portugal and Western Islands	4					29	1	84
Other Countries	16	6	2	2	2	1	70	99
Whole number according to the mother's birth-place	1,424	743	286	53	167	30	86	2,788



per cent. more than the children of the same class in the previous year. Of the children of mixed parentage, the whole number was 502, or 18.00 per cent. of all; 9 less children and 0.23 per cent less than in 1881. The number of children of purely foreign parentage in 1882 was 1,140, or 40.90 per cent. of all; this was 11 less children and 0.16 per cent less than in 1881. Of these 1,140 children. 979, or 85.88 per cent., had parents who were natives of the same foreign country; and 161, or 14.12 per cent., had parents who were natives of different foreign countries.

NUMBER OF THE CHILD.

Table V. shows how many of the 2,788 children born in 1882 were the first, second, third, etc., children of their mothers; also the same facts in regard to the 53,452 children born during the twenty-seven years preceding, 1855-1881, inclusive:—

NUMI	BER	OF	CHILD.	1892.	27 years, 1855-1881.	NUME	BEI	OF	CF	IILD.	1882.	27 years, 1853-1861.
First oli	ild (of th	ne mother	689	18,060	Trocifik	ch'	ld of	the	moth'r	17	213
Second	44	••	61	507	10,920	Thirteen	4Å	**	••	44	9	67
Third	••	16	44	8 85	8,527	Fourtees	nth	44	44	44		45
Fourth	14	**	**	814	6,465	Fifteenti	A	66	44	**	1	23
FI/th	**	44	44	250	4,742	Sixteent	A	44	46	44	1	7
Sixth	**	41	+4	901	8,397	Seventes	mLA	**	**	66		7
Seventh	44	••	**	140	2,414	Eighteen	ιLA	**	44	44		1
Eighth	44	44	.4	102	1,546	Nineteen	ιtλ	**	86	**		
Ninth	44	44	44	56	998							
Tenth	**	•	"	29	647	Whole				chil-		
Elevent)	."	64	64	23	847	dren.	•••	• • • • •	••••	•••••	2,788	58,452

TABLE V. BIRTHS, 1882. Number of the Child.

Of the 2,788 children born in 1882, 689, or 24.71 per cent., were the first children of their mothers; this was 2.45 per cent. more than the same class of children in 1881. Of the 53,452 children born in the twenty-seven years, 1855–1881, inclusive, there were 13,060, or 24.43 per cent., who were the first children their mothers had borne.

Average number of children to each mother. The number of children born in 1882 was 2,788; but as the number of twin births was 29, the number of mothers bearing children during the year was 2,759. Table VII shows that these mothers had borne in all, 9,974 children, which gives an average to each mother of 3.62 children.

The next table shows the average number of children to each mother, in each of the twenty-eight years, 1855–1882, inclusive. This table shows the number of children the mothers included in the figures had borne, previous to the end of the year 1882; but it does not show the average number of children the same mothers may have in their life-time:—

TABLE VI. BIRTHS, 1855 TO 1882.

Average number of children to each mother.

	No. of	No. of	Aggregate.
Years.	Mothers.	Children.	No. of Children. Average.
1855	1,588	1,600	
1856	1.638	1,675	5,563 5,563 5,563
1857	1,664	1.688	5,7368.44 children to each mother.
1859	1,708	1,724	5,6853.33 children to each mother.
1859	1,570	1,593	5.6183.58 children to each mother.
1860	1,627	1.648	5,8413.59 children to each mother.
1861	1,694	1,725	6,0173.55 children to each mother.
1862	1,519	1,529.	5,322 3.52 children to each mother.
1863	1,296	1,307	4,7173.64 children to each mother.
1864	1,327	1.844.	4,8873.68 children to each mother
1865	1,279	1,298	
1866	1,614	1.632	5,7448.56 children to each mother
1867	1,595	1.624.	5,4973.50 children to each mother.
1868	1,841	1,866.	6,3383.44 children to each mother
1889	1.797	1.819.	5,9543.30 children to each mother
1870	1,859	1,889.	6,1458.81 children te each mother
1871	1,937	1,960.	6,4203.81 children to each mother
1872	2,174	2,206.	7,3463.36 children to each mother.
1873	2,104	2,128.	6,7313.20 children to each mother.
1874	2,738	2,780.	8,9603.27 children to each mother.
1875	2,631	2,664	8,9773.41 children to each mother.
1876	2,604	2,631.	8,914 8.42 children to each mother
1877	2,516	2,543.	8,705
1878	2,597	2,627.	9,4173.63 children to each mother.
	2,499		
1880	2,600	2,627.	9 386 8.61 children to each mother.
1861	2,780	2,803.	10,1113.64 children to each mother.
			9,9743.62 children to each mother.
			192,7858.47 children to each mother.

Table VII. gives the number of the child in connection with the age of the mother; shows how many first, second, third, etc., children were borne in 1882 by mothers at the ages given; the number of mothers of each age, and the aggregate and average number children they had borne:—

TABLE VII. BIRTHS, 1882. Number of child with age of mother.

<u>-</u>			-		KUM	BER	OF	СНІ	LD.	- =			-	_	d d	8	orne.	children
AGROFTII MOTHERS IN YBARS	•	2.1.	3d.	tth.	Sth.	6th.	7tb.	8th.	91).	10th.	Talli.	13th.	Idh.	13th.	Number of mothers at each	Children they bore in 1882.	Total children they had borne	Average number of chil
15	. 1		i.							.		.			1	1	1,	1.0
16	. 6	2.5		ļ	 .	ļ				٠. .		٠.			5	5		1.00
17	. 13	2	-141-				 .	ļ	1. 1	.	. .	٠,٠		٠.١.	15	15	17	1.13
18	. 10	3	1	ļ. .		ļ		ļ		.		.			20	20	25	1 2
19	. 35	30			ļ. .		ļ	ļ		.	. .	٠			44	44	53	1.9
20	. 38	20	8			 .	 	ļ		.	. .				66	66	102	1.35
81			5	4			٠			.	. .		۱.	٠.	92	92	134	1.46
22	. 80	3.7	12	3	2			١			.].,	.J.		٠	134	131	212	1.58
3	. 68	15	24	5	3	2		 .	.		٠.	. .			145	145	267	1.14
4	. •5;	140	321	15	2	2		ļ	ļļ.	١.	<u>.ļ.</u> .	J. I	!		144	148	800	
5	. †71	65	*35	19	9	4		•••		. .	٠.				190	193	426	2.24
6 .	37	+3.5	45	*23	13	8				. .	٠.	ļ	٠.إ.		139	141	368	2.65
7	48	45	34	20	20	4	2	1			.			١	174	174	466	
8	43	1.5	26	•31	15	8	1	•••		. .	٠. .		ب.		177,	179	489 2	2.76
9	†31	80	23	29	21	+17	8	1	1.	. .	٠١	.	٠.¦.		160	163	554 3	3.46
0	26	31	28	41	26	+25	11	6	1	1	.	!	٠. .	.1	195	196	734 8	3.87
1	14	12	1:	20	17	13	4	8	1	1	104	104	409 3	3.SI
2	20	17	20	28	23	15	12	8	1	٠. و	.		.		152	152	626 4	1.12
8	3	13	16	22	+21	,17	8	4	21.	ď	 	!			105	106	478 4	1.50
4	8	•11	1	-16	19	7	•14	9	4.	. 1	ıl		١.	.	100	103	477 4	1.77
5	6	8	ь	*17	•22	24	•17	•19	9	3 2	2 1		2.	.	133	138	1	5.91
3	4	6	14	•14	22	17	8	12	•4	2 4	2	1	٠. .		108	110	602 5	.57
7	l]	6	7	10	8	13	19	•7	- 1	5 1	1	ı.	١.	.ll	77	78	475 6	3.17
3	1	1	+3	21	12	14	17		10.	1		2	1.	.1	91	93	595 6	.54
9		9	3	5	9	6	9	7		3 8					60	60	431 7	.18
0	1	31		5	5	2	E	8		8 8		2.		.[.]	43	42		.07
1]	2		2	8 .		2	3		2, 3		1.	J.		20	20	1	.60
2		1	6	ı	6	4	2	4	5	. 2				. I i	81	84		.09
l		1			2	2	4	i	11.	2	1		1		14	14	- 1	.57
	1	16		1	2	1	2		1		li		.].		10.	10	- 1	.70
	1	T.			1	ıl.			1	Ι.					3	8		.66
		. 1]	1			li		Π.	1		2	9	18 9.	
		- 10	0.74								$ \cdot $	٦.	.].		1	1	13 13.	
)			예.	.								.			1	1	3 8.	
					!-	201	140			-		1-	- 1				.974. 3.	

[•]One of these mothers bore twins. †Two of these mothers bore twins.

The reference marks in this table indicate the number of the first child in each case of plurality birth; thus, at the age of 24, four mothers bore twins; in one case they were the first and second children; in two cases, the second and third, and in one case the third and fourth. At the age of 29, three mothers bore twins; in two cases they were the first and second children, and in one case, the sixth and seventh.

The oldest mother in 1882 was 50 years of age and bore her third child in 1882. The number of mothers of 40 years and over, bearing children in 1882, was 128; the number of children borne by them during the year, 128; the total number of children borne by them, 927; an average of 7.24 children to each mother. In the period of nineteen years and six months, during which the record has been kept of the ages of the mothers, there have been 2,052 mothers of 40 years and over, who have borne 2,074 children in these nineteen years and a half. Including all the children previously borne by the same mothers, we have a total of 14,833 children; an average of 7.23 children to each mother.

During this nineteen years and six months, since July 1, 1863, the number of children borne by mothers whose age was stated, was 42,312; the number of children to mothers at each age, was as follows:—

Age of mother.	Children.	Age of mother.	Children.	Age of mother. Ch	ildren
14	5	. 27	2,540	40	918
15	20	28	2,957	41	278
16	62	29	2,180	42	349
17	180	30	3,353	43	187
18	433	31	1,453	44	118
19	729	32	2,036	45	114
20	. 1,173	3 3	. 1,679	46	32
21	. 1,374	34	1,432	47	27
22	,037	3 5	2,127	48	12
23	2,256	36	1,319	49	8
94	2,489	87	1,068	50	11
25	2,844	38	1,161	_	
26	2,578	39	753	All known ages42	,312

These 42,312 children were divided as follows, to mothers of different divisions of age:—

	Number of mothers.	Per cent
Under twenty years	1,429	3.3
Twenty, and under twenty-five	9,829	22,0
Twenty-five, and under thirty	18,099	30.96
Thirty, and under thirty-five	9,953	23,5
Thirty-five, and under forty	6,428	15.19
Forty, and under forty-five	1,875	4.48
Forty-five and over		0.47
		-
Total	42,812	100.00

PLURALITY BIRTHS.

The number of cases of plurality births in Providence in 1882 was 29, which was 6 more than in 1881.

The following are the statistics concerning them :-

Cases. 29; number of children, 58.

Locality. First ward, 4; Second, 0; Third, 2; Fourth, 3; Fifth. 2; Sixth, 4; Seventh, 2; Eighth, 3; Ninth, 2; Tenth, 7.

Season. January; 2; February, 2; March, 1; April, 3; May. 3; June, 1; July, 2; August, 3; September, 2; October, 1; November, 2; December, 7.

Sex. In 11 cases both children were males; in 10, both were females; and in 8 cases there was one child of each sex; in all, 30 male and 28 female children.

Color. The parents were white in all cases.

Number of the Children and Age of the Mother. The reference marks in Table VII indicate the number of the children and the age of the mother in each case of plurality birth. In five cases they were the first and second children; the mothers being, one of 24 years, two of 25 years, and two of 29 years; in six cases they were the second and third children of mothers, two of whom were 24 years old, and the others, 26, 28, 34 and 35 years; in three cases, the third and fourth children of mothers 24, 25 and 38 years old; in five cases, the fourth and fifth children of mothers 26, 28, 34, 35 and 36 years old; in two cases, the fifth and sixth children of mothers 33 and 35 years old; in two, the sixth and seventh children of mothers 29 and 30 years old; in two, the seventh and eighth children of mothers 34 and 35 years old; in two, the eighth and ninth children of mothers 35 and 37 years old; in one case, the ninth and tenth children of a mother 36 years old; and in one, the tenth and eleventh children of a mother 38 years old.

Parentage. In thirteen cases both the parents were Americans; in seven, both were Irish; in one, both German; in one, both Norwegian; in one, both Italian; in one, both British American; in two, the father was American and the mother Irish; in one, the father Irish and the mother American; in one, the father Irish and the mother Scotch; and in one, the father British American and the mother English.

TWENTY-EIGHT YEARS' SUMMARY OF PLURALITY BIRTHS.

In the twenty-eight years, 1855-1882, there have been 662 cases of plurality births in Providence; 653 of twins and 9 of triplets.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children and one female; making 9 male and 18 female children in the nine cases.

Of the 653 cases of twins, there were 215 cases where both children were males; 221 in which both were females, and 217 in which there was one male and one female child; total, 647 male and 659 female children. Of the 1,333 children of all the cases of plurality births, there were 656 males and 677 females; a proportion of 103.2 females to 100 males, or 50.79 girls and 49.21 boys in each 100 children born.

Color. In all the cases of triplets the parents were white; of the 653 cases of twins, in 602 the parents were white, and in 51 they were colored.

Parentage. Of the triplets, in one case the parents were both American; in five cases, both were Irish; in one case, both German; in one, the father was English and the mother Irish; and in one, the father Scotch and the mother Irish. There was but one American mother in the nine cases.

The nativity of both parents is given in the following table for the 653 cases of twin births; the first letter indicating the father's birth-place and the second the mother's.

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British American; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish; N., Norwegian.

Parentage. No.	Parentage. No.	Parentage. No.	Parentage. No.
A. and A 220	I. and A16	B. A. and I 2	I. and S
1. and I247	E. and A14	Sw. and [1	E. and S 1
E. and E 30	S. and A 2	It. and I1	I. and W
8. and 8 7	B. A. and A 6	P. and I	A. and B. A6
G. and G 17	G. and A 2	A. and E4	I. and B. A
B. A. and B. A 11	Sw. and A 1	I. and E	F. and B. A
D. and D 1	A. and I15	S. and E1	S. and B. A
N. and N 1	E. and I12	G. and E1	A. and Sw
P. and P 0	S. and I 8	B. A. and E1	_
It. and It 2	G. and I 1	A. and 81	Total653

The number of American mothers in these 653 cases was 261, which was 39.97 per cent.; the number of Irish mothers was 283, or 43.34 per cent.; of mothers of others foreign nations, there were 109, or 16.69 per cent.

In the 662 cases of plurality births in the twenty-eight years, there were 262 American mothers, and 400 foreign mothers. The whole number of American mothers who bore children during these years was 26,323; of this number, the 262 mothers in the cases of plurality births, were about 1.00 per cent.; the whole number of foreign

mothers during the same time was 29,246; and the 400 mothers in the cases of plurality births were 1.37 per cent. of this number.

Proportion of plurality births to whole number. The following figures compare the year 1882 with the twenty-seven years previous, as to the number and proportion of plurality births:—

Single Births. Twin. Triple 27 years, 1855–188152,1776249 1882 2,730 29	52,81053,452
28 years' totals54,9076539	55,56956,240
Proportion of cases of plurality births:	_
Twins.	Triplets.
1855-1881 1 in 84.6	1 in 5,867.8
1882 1 in 95.1	••••
1855-1882 1 in 85.1	1 in 6,174.8
Proportion of children of plurality birth	ns :
Twins.	Triplets.
1855-1861 1 in 42.8	1 in 1,778.6
18821 in 48.1	*****
1855–1882 in 43.1	1 in 2,082.9

Proportion of plurality births by parentage. The whole number of births in the twenty-eight years, 1855–1882, inclusive, was 55,569. The following figures show the number of births of American, foreign and mixed parentage, and the number and proportion of plurality cases in each class:—

	Whole No. Births.	Plurality	Cases.	1 in	Per C
American parents	22,334	221	• • • • • • •	101.1.	or 0 9
Foreign parents	26,066	878	••••	69.8.	or 1.4
Am. father, foreign n	nother. 8,180	27		117.7.	or 0.n
Foreign father, Am. n	nother. 3,989	41		97.2	or 1.0
		-	_		_
Total	55,569	662	• • • • • • •	83.9	• or 1.1

Of the 662 cases of plurality births during this period, the 221~nl American parentage were 33.38 per cent.; the 373 of foreign parentage were 56.35 per cent.; the 27 with American father and foreign mother were 4.08 per cent.; and the 41 with foreign father and American mother were 6.19 per cent.

MARRIAGES.

During the year 1882 there were 1,172 marriages in the city of Providence.

The following table shows the number of marriages in each month and quarter of the year 1882, and in each of the six years preceding; also the aggregate and average number for each month and quarter of the twenty-eight years, 1855–1882:—

TABLE VIII. MARRIAGES. Season.

MONTHS.	1882.	1881.	1880.	1879.	1878.	1877.	1876.	28 years, 1855–1882.	28 years' average.
January	105	117	105	102	76	68	79	2,128	76
February	82	112	100	85	73	65	93	1,905	68
March	55	81	53	47	78	88	44	1,246	45
lst Quarter	242	810	258	234	223	171	216	5,279	189
A pril	109	95	98	83	77	88	70	1,985	69
May	91	114	109	98	68	85	80	2,098	75
June	101	181	111	96	104	88	94	2,201	79
2d Quarter	301	840	818	277	249	251	244	6,234	228
July	70	88	97	59	94	25	56	1,833	65
August	80	69	75	88	62	76	62	1,593	57
September	111	81	115	106	93	87	93	2,216	79
3d Quarter	261	238	287	253	249	248	211	5,612	201
October	125	134	108	103	99	99	95	2,488	89
November	149	114	148	128	115	102	109	2,701	96
December	94	66	118	76	82	82	74	1,949	70
4th Quarter	368	314	869	807	296	283	803	7,188	255
Yearly totals	1,179	1,202	1,282	1,071	1,016	958	949	24,295	968

There were 30 less marriages in 1882 than in 1881, but 304 more than the average for the twenty-eight years. Having estimated the population of the city in 1882 at 114,377, the 2,344 persons who were married during the year give a ratio of one in 48.80 of the population, or 20.49 persons married in each thousand. The ratio of persons married, to births, was 84.07 to 100; of persons married, to deaths, 104.5 to 100.

Season. The larger part of these marriages occurred in the last part of the year; the largest quarterly total being 368, in the fourth quarter; the smallest quarterly total was 242, in the first quarter. The largest monthly total was 149, in November, and the smallest, 55, in March.

The number and percentage of marriages in each quarter of the year 1882 are given below; also the same statistics for the twenty-seven years, 1855–1881:—

	180	32.	1855-1861.			
¥	larringes.	Per Cent.	Marriages.	Per Cent		
January-March	242	20.65	5,037	21.73		
April-June	801	25.68	5,938	25.60		
July-September	261	22.27	5,381	23.3		
October-December	868	81.40	6,770	29.2		
Total	1.179	100.00	98.191			

The birth-places of both parties in each of the 1,171 marriages in 1882 are given in the following table:—

TABLE IX. MARRIAGES, 1882. Nativity of the parties.

	BIRTH-PLACES OF BRIDES.								
BIRTH-PLACES OF GROOMS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number of grooms	
United States	216	52	89	8	39		4	753	
Ireland	3 8	124	3		6		1	172	
England, Scotland and Wales	38	15	59		7			119	
Germany	10	4	8	14	ļ			81	
British America	21	8	2		28			54	
Portugal and Western Islands	3	ļ ,		 .		7		10	
Other Countries	7	1		1	<u> </u> 	 -	24	88	
Whole number of brides	783	199	106	18	80	7	29	1,179	

Bride and groom born in the same country. Of the 1,172 marriages during the year, there were 872 in which the bride and groom were natives of the same country; this was 74.40 per cent. of all; this was 18 less marriages but 0.36 per cent. more than in 1881. There were 616 marriages, or 52.56 per cent. of all, in which both parties were natives of the United States; this was 59 less marriages and 3.60 per cent. less than in the previous year. There were 124 marriages, or 10.58 per cent., in which both parties were born in Ireland; this was 17 more marriages than in 1881. The 59 marriages where both were natives of England or Scotland, were 18 more than the number of the same class the year before; there were 14 marriages where both were natives of Germany; 4 more than in 1881; and 28 where both were natives of British America; 9 less than in 1881.

Bride and groom born in different countries. There were 300 marriages in 1882, or 25.60 per cent. of all, in which the parties were natives of different countries; 12 less marriages and 0.36 per cent. less than in 1881. Of these 300 marriages there were 137, or 45.67

Mixed.	oreign.		American.
Kumber. Per Ce	Per Cent.	ent. Numi	Sumber. Per (
59 8	41.93	265.	30848.7
7912	32.97	212	352 54.7
55 8	28.50	175	38402.5
6311	38.21	214	2×350.5
54 8	36.66	224	33354.56
59 9	36.29	225	33654.1
6111	33.27	182	30455.5
8014	82.72	179	28852.6
7619	31.47	193	34256.0
10118	28,55	211	42757.78
9012	28.49	202	41758.8
9711	30.05	244.	47158.00
10111	31.46	269.	48556.7
127 14	24.65	215	53060.7
11412	30.75	286.	53056.99
14515	27.32	262.	55257.56
14215	29.48	278.	52355 40
18817	28.14	303.	591 54.5
18115	27.57	317.	65256.69
18215	29.42	346.	64855 10
17316	29.66	820.	58654.3
161 16	25.39	241.	54757.6
17518	24.45		54557.19
17016	24.12	245.	CO159.19
20118	20.26	217.	65260.97
	19.72		75060.88
26221	22.04	265.	67556 16
25421	25.77	302.	61652.50

have been only two years in the twenty-eight when the proof American marriages was less than in 1882; the proportion n marriages was larger than it has been since 1875; the on of mixed marriages, though slightly less than in 1881, was an in any year previous to that. The whole number of marven in this table is 24,287; while in Table VIII., the total ame period of years is 24,293; this discrepancy is owing to hat in some of the early years of the records it was necesmit some marriages from each of these tables on account of statistics. See Report for 1879, page 22. per cent., where the groom was a native of the United States; and 117, or 39.00 per cent., in which the bride was born here. These 137 American grooms who married foreign wives were 18.20 per cent. of the whole number of grooms born in the United States, which was 753. Of the 172 grooms born in Ireland, only 48, or 27.90 per cent., married natives of other countries. Of the 119 grooms born in England, 60, or 50.42 per cent.; of the 31 born in Germany, 17, or 54.84 per cent.; of the 54 born in British America. 26, or 48.15 per cent. Of brides there were 733 who were natives of the United States; of this number, 117, or 15.96 per cent., married foreigners. Of the 199 brides born in Ireland, 75, or 37.69 per cent., married natives of some other country than their own; of the 106 born in England or Scotland, 47, or 44.34 per cent.

The 33 grooms who were natives of "other countries" than those named in the table, were born as follows:—in Sweden, 11; in Italy, 8; in France, 5; in Russia, 2; in the West Indies, 2; in Switzerland, Denmark, Norway, Finland and China, one each.

There were 29 brides who were of "other countries," and were born as follows:—in Sweden, 10; in Italy, 8; in Norway and France, 3 each; in Russia, Switzerland, West Indies and Africa. one each; and one was born "at sea."

In the following table are given the number and percentage of American, foreign and mixed marriages, in each of the twenty-eight years from 1855-1882, inclusive. By mixed marriages are meant those in which one of the party was of American, and one of foreign nativity:—

MARRIAGES: NATIVITY.

	Ame	rican.	F	oreign.	Mixed.
Year.	Number.	Per Cent.	Number.	Per Cent.	Number. Per Cent
1865	308	48.73	265	41.93	59 9.8
18 56	352	. 54.74	212	82.97	7912.2
1857	384	62.54	175	28.50	55 8.9
1858	283	50.54,	214	38.21	6311.2
1859	333	54.50	224	36.66	54 8.8
1860	336	54.19	225	86.29	59 9.5
1861	304	55.58	182	33.27	61
1862	288	52.65	179	82.72	8014.6
1863	342	56.07	192	81.47	7612.4
1864	427	57.78	211	28.55	10113.6
1865	417	58.82	202	28.49	9012.6
1866	471	58.00	244	30.05	9711.9
1867	485	56.73	269	31.46	10111.8
1868	530	60.78	215	24.66	127 14.5
18 69	530	56.99	286	80.75	114 12.2
870	552	57.56	262	27.32	14515.1
1871	523	55 46	278	29.48	14215.0
872	591	54.52	805	28.14	18817.8
873	652	56.69	317	27.57	18115.7
874	648	55 10	846	29.42	18215.4
875		54.31			17316.0
876	547	57.64	241	25.39	161 16.9
877	545	57.19		24.45	
878	€01	59.15	245	24.12	17016.7
879	653	60.97	217	20.26	20118.7
880 ,	750	60.88	243	19.72	23919.4
881	675	56 16	265	22.04	26221.8
382	616	52.56	302	25.77	25421.6

There have been only two years in the twenty-eight when the proportion of American marriages was less than in 1882; the proportion of foreign marriages was larger than it has been since 1875; the proportion of mixed marriages, though slightly less than in 1881, was larger than in any year previous to that. The whole number of marriages given in this table is 24,287; while in Table VIII., the total for the same period of years is 24,293; this discrepancy is owing to the fact that in some of the early years of the records it was necessary to omit some marriages from each of these tables on account of imperfect statistics. See Report for 1879, page 22.

The following figures give the same facts for each of the twenty-seven years, 1856-1882 inclusive, and for the whole period:—

	Males.	Females.
Years.	No. Aggregate age. Average age.	No. Aggregate age. Average u
1856	639 18,104 years28.33 years	637 15,559 years24.42 year
1857	607 16,753 years27.59 years	606 14.427 years23.84 year
1858	560 15,850 years28 29 years	560 13,702 years 24.47 year
1859	611 16,989 years27.81 years	611 14,616 years23.92 year
1860	620 17,135 years27.60 years	620 14.667 years23.66 year
1861	547 15,448 years28.24 years	547 13,316 years
1862	547 15.773 years28.83 years	547 13,565 years24.80 year
1863	610 17,208 years28.21 years	610 14,792 years24 24 year
1864	739 21,361 years28.90 years	739 18,075 years24.45 year
1865	709 20,539 years28.96 years	709 17,605 years24.83 year
1866	812 22.801 years28.08 years	812 20,036 years24.67 yea
1867	855 24 443 years28.58 years	855 21,078 years24.65 yea
1868	872 24.546 years28.15 years	872 21.440 years24 59 yea
1869	930 26,548 years28.55 years	930 23,135 years24.88 yes
1870	959 27,343 years28.51 years	959 23,773 years24.79 yes
1871	943 26,743 years28.37 years	943 23.290 years24.70 yea
1872	1.084 31,110 years28.70 years	1,064 26,835 years24.76 yea
1873	1,150 32,879 years 28.59 years	1,150 28,701 years24.96 yea
1874	1,176 33,927 years28.85 years	1,176 29,510 years25.09 yea
1875	1,079 30,848 years28.59 years	1,079 27,090 years25.11 yea
1876	949 27,496 years28.97 years	949 23,640 years24.91 yea
1877	953 27,904 years29.28 years	953 24,092 years25.28 yes
1878	1,016 29.460 years28.99 years	1,016 25,796 years25 40 year
1879	1,071 31,016 years28.96 years	1,071 27,103 years25.32 yea
1880	1.232 36,127 years29.32 years	1,232 31,015 years25.18 yes
1881	1.202 35,283 years29.35 years	1,202 30.801 years25.62 yea
1882	1,172 31,068 years29.07 years	1,172 29,743 years23.38 year

The number of males and females does not agree in this table because in 1856 the ages of 17 males and 19 females, and in 1857, the ages of 7 males and 8 females, were not given.

Number of times married. Table XI. shows how many of the marriages in 1882 were the first, second or third marriages of the parties:—

GROOMS.	BRIDES. N	Whole		
Number of the Marriage.	First.	Second.	Third.	Grooms.
First	881	78	1	960
Second	123	67	3	193
Third	8	7	3	18
Fourth	1		• • • • • • • • • • • • • • • • • • • •	1
Whole number of Brides	1,013	152	7	1,17

TABLE XI. MARRIAGES, 1882. Number of times married.

The 881 marriages which were the first of both parties, were 75.17 per cent. of the whole number. The whole number of grooms married for the first time, was 960 or 81.91 per cent. of all; of brides, 1,013 or 86.43 per cent. of all. In 67 cases, or 5.72 per cent. of all, it was the second marriage of both parties.

There were, in 1882, 13 marriages in which both parties had been divorced; 20 more in which the man only had been divorced; and 32 others where the woman only had been; making 65 marriages where one or both parties had been divorced; this was 5.55 per cent. of the whole number of marriages, or about one in 18.

Denominational. The 1,172 marriages of 1882, were solemnized by the clergymen of different denominations or by civil authority as follows:—

Roman Catholic438	Latter Day Saints 3
Calvinistic Baptist144	Hebrew 1
Free Baptist 40	Bethel Pastor 5
Methodist179	Lutheran 3
Episcop alian136	New Jerusalem 5
Congregationalist 97	Free Religious 8
Unitarian	Advent 7
Universalist 36	African Union 6
Presbyterian 13	Denomination not specified 6
Christian 9	Judge of Supreme Court 2
Seaman's Mission 8	
Evangelical Advent 4	Total1,173

Nativity and Residence. There were in 1882, 96 marriages, in which both parties were natives of Providence, this was 8.19 per cent. of all the marriages during the year; there were also 115 other marriages, or 9.81 per cent. of all, in which the groom only was born here; and 175 or 14.93 per cent., in which the bride only was born here; this gives a total of 482 persons married, who were natives of the city; this is 20.56 per cent. of the 2,344 persons married in the city during the year. This percentage varies but little from year to year; in 1881 it was 21.76 per cent.; in 1880 it was 22.20 per cent.; in 1879 it was 22.60 per cent.; in 1878, 21.06 per cent.

There were 153 marriages in 1882, or 13.06 per cent. of all, in which neither party was a resident of the city; of these 153 marriages there were 96, or 62.74 per cent. where both parties were not only not residents of the city, but came from outside of the state; and in 74 cases they both came from Massachusetts; in 11 others one of the parties was from that state.

These numbers are all much smaller than the corresponding figures on these points for the past two years, as may be seen from the following table:—

Both parties non-resident.	Per cent.	Both from outside. the state.	Both from Massachusetts	One from
1880 206	16.72	182	104	15
1881 195	16.22	180	102	13
1682 153	13.06	96		11

This decrease is probably the effect of the new marriage law, which went into operation on the first of July, 1881, and one object of which was to accomplish just this result; namely, to prevent the marriage here of runaway parties from neighboring towns and states; especially of minors. That this is the cause of the decrease is still further apparent from the fact, that, for 1881 and 1882, the statistics were made up for the first and last half of the year separately; and in 1881, the figures for the second six months after the new law was in force, were very much less than for the first six months, while in 1882, when the law operated during the whole year, there was very

little difference between the first and last half of the year. Thus in 1881, the number of marriages of non-residents was 195, of which 121 were in the first six months, and 74 in the last six; there were 130 marriages where both were from outside the state; 82 in the first six months, and 48 in the last six; 102 where both parties came from Massachusetts; 67 in the first six months and 35 in the last six.

In 1882, the 153 marriages of non-residents were divided in the first and last half of the year much more evenly; 65 in the first half, and 88 in the last half; of the 96 where both parties were from outside the state; 40 were in the first half, and 56 in the last half; of the 74, where both came from Massachusetts, 35 were in the first half, and 39 in the last half.

The whole number of marriage certificates issued during the year, was 1,245; in 86 cases the parties went out of the city to be married; on the other hand, there were 58 cases where persons came into Providence to be married, with certificates from some other town in the state; so that the marriage is recorded and counted here, though the "Intention of Marriage" is not. There were 45 certificates given which were either not used, or of which no report has been received.

Marriage and education. The 2,344 persons married in Providence in 1882, were divided as follows as to sex and nativity:—

Born in United StatesMa	.les, 753	.Females,	783	Total,	1,486
Born in Foreign countriesMai	les, 419	.Females,	439	rotal,	858

In these different classes, the number and proportion who signed the marriage certificate with a mark, were as follows:—

1882.	Whole Number Married.	Signed with a Mark.	In Each 100.
Males, born in United States	753	28	3.72
Females, born in United States	733	28	3.14
Total, born in United States	1,486	51	8.48
Males, born in foreign countries	419	66	15.75
Females, born in foreign countries	439	88	18.91
Total, born in foreign countries	858	149	17.87
Totals	2,844	200	8.58

Of the 51 persons born in the United States, who signed the marriage certificate with a mark, there were 15 colored persons (11 males and 4 females); and 29 (13 males and 16 females) who were of foreign parentage; this leaves only 7 persons (4 males and 3 females) of white American parentage, who were unable to sign their names; this was 4 less than in 1881; 3 less males and one less female.

The number and proportion who signed with a mark in 1882 were much less in every class than in 1881; the number born in the United States was 21 less (5 males and 16 females), and the proportion 1.04 less; the number of those born in foreign countries, was 22 less (8 males and 14 females), and the proportion 4.22 less; the whole number in 1882 was 43 less than in 1881 and the proportion 1.58 less.

The number of males who signed with a mark were 13 less than in the previous year, 5 American and 8 foreign; the number of females, 30 less, American 16, and foreign 14.

The statistics on this subject for the twenty-three years, 1859-1881, are as follows:—

1869-1881.	Whole Number Married.	Signed with a Mark.	In Each	
Males born in United States	13,237	618	4.6.	
Females born in United States	13,514	895	R.63	
Total born in United States	26,751	1,513	5 60	
Males born in foreign countries	7,429	2,297	30.99	
Females born in foreign countries	7,152	3,822		
Total born in foreign countries	14,581	5,619	38.54	
Totals	41,832	7,182	17.95	

DEATHS.

The number of deaths reported for the year 1882, was 2,242. This was 97 more than in 1881; males, 33 more; females, 64 more; whites, 109 more; colored, 12 less; natives of the United States, 36 less; of foreign countries, 133 more; of American parentage, 34 less; of foreign parentage, 131 more.

The whole number of deaths in twenty-eight years, 1855-1882 inclusive, was 39,957; the average annual death rate, one in 50.86. For 1882 the death rate was one in 51.02 or 19.60 deaths to each thousand of the population.

In the different classes of the population, the ratio was as follows:-

	Estimated		•	Deaths to
	Population,	Deaths,	To population	each 1,000
	1682.	1882.	one death in	of population.
Whole number	114.877	2,242	51.02	19.60
Males	54,601	1,080	50.56	19.78
Females	59,776	1,162	51.44	19.44
White	110,401	2,147	51.42	19.45
Colored	3,976	95	41.85	23.89
Born in United State	s81,666	1,586	51.49	19.42
Born in foreign coun	tries32,711	656	49.86	20 05
American parentage	52,696	965	54.61	18.81
Foreign parentage	61,681	1,277	48.30	20.70

The general statistics in regard to the 2,242 deaths are given in the following table:—



TABLE XII. DEATHS, 1882.

Sex, condition, color, nativity, parentage, locality and season.

1862.	January.	February.	March.	April.	May.	June.	July.	August.	September.	October.	November.	December.	Whole Year
Whole number of Deaths	185	177	166	179	169	165	192	222	167	165	247	215	2,342
Sex.—													
Males Females	78 107	74 108	86 80	85 87	77 92	77 88	95 97	112 110	85 82	92 78	118 129	101 114	1,080 1,163
Condition.—													
Married	62	78	67	75	58	55	41	50	50	60	89	78	758
Single	88	75	81	78	76	78	126	145	98	81	127	108	1,151
Widows	26	22	12	16	25	20	17	21	12	14	19	22	236
Widowers	8	4	6	6	15	10	8	5	4	9	9	12	96
Divorced	1	8	••••	2	••••	2	• • • •	1	3	-1	8	••••	16
Color.—				- 1				- 1					
Whites	179	170	160	164	158	159	188	215	159	157	235	903	2,147
Colored	6	7	6	8	11	6	4	7	8	8	12	19	96
Nativity.—				i				- 1					
United States	127	128	120	109	117	116	145	165	129	115	169	146	1,566
Ireland	84	80	81	49	42	85	85	88	28	33	48	54	457
England and Scotland	15	7	7	11	5	9	8	15	4	13	17	10	121
Germany	8	4	1	2	1	8	1	1	2		1		19
British American	5	8	4	1	8	1	2	1	2	8	7	8	87
Portugal & West'n Islands.		1			1				- 10	1	1	1	5
Other countries	1	2	8			1	1	2	2		4	1,	17
Parentage.—							1					1	
American	86	88	71	63	78	65	76	94	71	75	108	96	965
Irish	61	58	67	77	71	74	81	78	72	60	93	93	885
English and Scotch	19	16	17	20	13	18	28	27	12	- 23	27	16	231
German	6	5	1	23	4	8	2	7	3	3	4	1	12
British American	6	5	6	. 6	4	8	6	9	4	2	6	6	Ø
Portuguese	2	1	1		8		2	5		- 1	1	1	17
Other foreign	5	4	3	- 8	1	2	2	2	5	- 1	8	3	39
Locality, Wards.—												- 1	
I	25	22	18	19	29	27	19	26	14	20	27	31	277
11	6	7	9	6	4	4	4	6	- 6	- 4	8	8	73
III	15	18	17	12	14	10	26	24	16	17	20	27	211
IV	11	14	9	- 9	5	10	to	8	9	10	18	13	196
v	19	20	7	13	17	15		15	7	10	23	17	168
VI	10	9	10	4	8	9	11	10	7	7	8	7	100
VII	18	9	17	20	15	18		16	18	10	26	17	195
VIII	16	19	15	18	18	10	21	17	19	12	24	11	200
IX	27	23	26	23	17	22	26	89	20	25	30	29	307
X	85	84	29	39	88	87	44	47	38	33	48	40	451
Public Institutions	8	7	9	0	9	8	11	14	13	12	20	14	134
Totals	185	177	166	179	169	168	192	222	167	165	247	215	2,24


The largest quarterly total in 1882, was 627 in the fourth quarter, the smallest, 506, in the second. In 1881, the smallest quarterly total was in the fourth quarter. The following figures give the number of deaths in each quarter of the year 1882, and of each of the five years preceding; also the aggregate and average numbers for each quarter, in the period of twenty-seven years, 1855–1881:—

						27 years, 1855-1881.
	1882.	1881.	1880.	1879.	1878.	1877. Aggregate. Av'rge.
January-March	528	. 527	. 560	. 514	541	. 411 9,120 888
April-June	506	. 557	. 583	. 489	423	. 410 8,438 812
July-September	581	. 580	. 525	. 498	520	. 55710,951 406
October-December.	627	. 481	. 402	. 580	505	. 530 9.205 841

The number and percentage for each quarter of the year 1882, and for the twenty-seven years, 1855-1881, were as follows:—

	188	3.	27 years, 1855–1881.		
:	Deaths.	Per cent.	Deaths.	Per cent.	
January-March	528	23.55	9,120	24.18	
April-June	506	22.57	8,438	22.87	
July-September	581	25.91	10,951	29.04	
October-December	627	27.97	9,206	24.41	
Totals	2,242	100.00		100.00	

The following figures give the number and percentage for each quarter of each of the twenty-eight years, 1855-1882 inclusive:—



•	Januar	y-March.	Apri	l-June.	July	-Sept.	OctDec.
	Deaths.	Per cent.	Deaths.	Per cent.	Deaths.	Per cent.	Deaths. Per ct
855	218	21.49	208	20.99	839	84.21	23123.8
856	308	28.92	244	23.91	296	27.80	21790.3
857	214	23.14	178	18.70	286	80.92	25227.2
858	279	27.43	278	27.84	257	25.27	20319 9
859	213	23.69	205	22.80	270	30.04	21123.47
.860	243	24.27	209	20.88	297	29.67	252 25 18
.961	240	22.84	236	22.45	836	31.97	23922.70
862	197	21.55	192	21.01	271	29.65	25427.79
963	287	23.64	248	20.48	870	81.14	801 24.70
964	802	23.58	273	21.31	408	31.46	30323.6
865	200	24.69	240	19.82	372	30.72	30024.7
866	238	23.97	218	21.04	839	32.63	2 4 2 23 34
867	258	26.88	202	21.04	286	29.79	21422.55
868	231	20.81	244	21.98	316	31.17	28926.04
869	831	26.35	250	19.90	863	28.91	31224.84
870	331	26.20	271	21.46	871	29.37	29022.9
871	811	24.80	263	20.98	385	80.70	29523.3
872	830	20.59	361	22.71	518	32.31	39124.39
8 73	427	24.84	407	23.68	504	29.82	38122.10
874	414	22.35	501	25.21	556	27.98	486 24.46
875	488	25.48	401	20.94	531	27.73	49525.s
876	383	20.54	419	22.47	573	80.72	49026.7
877	411	21.21	440	22.70	557	28.74	530 27.3
878	541	27.20	423	21.27	520	26.14	50525.3
879	514	25.37	439	21.67	493	24.33	68028.6
880	569	26.92	533	25.63	525	25.24	40222.21
881	527	24.57	557	25.97	560	27.04	48122.47
882	8.30	09 88	E00	00 87	801	07.01	007 07 0

In the first quarter of 1882, the number of deaths was one more than in the first quarter of 1881, the percentage 1.02 less than in the preceding year. In the second quarter the number and percentage were both considerably less in 1882 than in 1881. In the third quarter, which generally shows the largest number and percentage of the year, the number in 1882 was one more than in 1881, but the percentage 1.13 less. But though this number was the largest quarterly total in 1881, it was not so in 1882; for the epidemic of typhoid fever which prevailed in the last quarter of the year raised the percentage of that quarter to the highest of the year; the number was the largest ever recorded for the fourth quarter, and the percentage larger than in any year except 1879.

Sex and Parentage. The 2,242 decedents of 1882, were divided as follows as to sex and parentage:—

American parentage	465 males	500 femalestotal,	965
Foreign parentage	615 males	662 femalestotal,	1,277
-			
Totals	,080 males1	,162 femalestotal,	2,242

There were 82 more females than males among the decedents; 35 of American and 47 of foreign parentage. There were 312 more decedents of foreign than of American parentage; 150 males and 162 females.

The number of decedents of each sex in the twenty-eight years, 1855-1882 inclusive, was 19,549 males and 20,408 females; total 39,957.

Condition. The number of married persons who died in 1882, was 753; 301 of American and 452 of foreign parentage; it was 84 more than those of the same class in 1881.

The number of single decedents in 1882, 1,151, was 30 less than in 1881; they were 487 of American and 664 of foreign parentage.

The number of decedents who were widows was 226, or 16 more than in the preceding year; they were 116 of American, and 110 of foreign parentage.

The number of widowers who died was 96; 21 more than in the preceding year; 47 of American, and 49 of foreign parentage.

The number of divorced persons who died in 1882, was 16; this was 6 more than in 1881; 14 were of American, and 2 of foreign parentage.

Color. There were 2,147 white decedents in 1882; 874 of American, and 1,273 of foreign parentage; it was 109 more than the number in 1881. The number of colored decedents in 1882, was 95, or 12 less than in the preceding year. Of these 95 decedents, 4 were of foreign parentage. There were 2,692 white children born in 1882; as there were 2,147 white decedents, the proportion of births to deaths in this class was 100 to 79.8. The number of colored children born in 1882, was 96; the number of colored decedents 95; the proportion of births to deaths, 100 to 98.9. In the twenty-eigh

years, 1855–1882, the whole number of colored children born was 2,155; the number of colored decedents in the same time was 2,251; the proportion of births to deaths, 100 to 104.5.

Nativity and Parentage. The decedents in Providence are always registered under both of these classes, and the figures of table XII. show how great is the difference between the two. Thus the number of decedents in 1852, who were natives of the United States, was 1,586 or 70.74 per cent. of all; the number of American parentage was 365, or 43.04 per cent. The number of American nativity was 36 less than in 1881; the percentage, 4.88 less; the number of American parentage was 34 less than in the previous year, and the percentage 3.53 less.

The number of decedents of foreign birth in 1882, was 656, or 29.26 per cent. of all; this was 133 more than in 1881, and 4.87 per cent. more. The number of foreign parentage was 1,277, or 56.96 per cent. of all decedents; it was 131 more than in 1881, and the percentage 8.52 more.

The 17 decedents who were natives of "other countries" than those named in the table, were as follows: 6 were born in Italy; 5 in Sweden; 2 each in Russia and Denmark; and one each in France and Norway.

The 89 decedents of "other foreign" parentage, were as follows: Italian, 13; Swedish, 11; Danish, 4; French, 3; Russian, Austrian and Norwegian, 2 each; Polish and West Indian, 1 each.

Locality. Table XII. gives the number of deaths in each ward and in Public Institutions in each month of the year. The number of deaths in the first ward in 1882, was 5 less than in 1881; in the second ward, 25 less; in the third, 18 more; in the fourth, 4 more: in the fifth, 22 less; in the sixth, 14 less; in the seventh, 10 more; in the eighth, 15 less; in the ninth, 45 more; in the tenth, 87 more: in Public Institutions, 19 more.

The following was the number of decedents, according to parentage, in each ward:

Wards.	American.	Foreign.	Wards.	American.	Foreign.
I	122	155	VI	63	87
II	61	12	VII	83	119
ш	85	128	VIII	119	86
ı ⊽	64	62	IX	157	150
V 70 98			x	97	854
			Public In	stitutions 51	85
Whole	alt-			 965	1 977

The number of deaths in Public Institutions in 1882, and in each of the two years preceding, was as follows:—

•	1892.	1881.	1880
Rhode Island Hospital	76	53	4
Butler Hospital	22	18	•••••
Dexter Asylum	15	21	1
Home for Aged Men	1	0	
Home for Aged Women	5	8	
Children's Home	1	0	
Children's Hospital and Nursery	0	0	• • • • • •
R. C. Orphan Asylum	8	2	
R. C. Convent	1	2	•••••
Beform School	9	1	
Little Sisters' Home for the Poor	6	1	
Friends' School	0	6	
Shelter for Colored Children	0	8	
Hotels	1	4	• • • • • • •
Police Station	1	1	•••••
Total			

Aggregate and Average Age. Table XIII. shows the number of deaths, and the aggregate and average age of the decedents, according to parentage, for each month, and for the whole year 1882:—

Table XIII. Deaths, 1882.

Aggregate and average age, by parentage.

	Amer	ican Par	entage.	Fore	gn Pare	ntage.	Total	America Foreign	an and
MONTHS.	Number of Deaths. Aggregate Agg.		Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.
January	86	8,828	38.69	99	8,235	82.67	186	6,563	25.4
February	88	8,656	42.47	89	2,835	81.85	177	6,491	36.G
March	71	2,562	86.08	95	2,881	80.32	166	5,443	82.7
April	63	2,146	34.06	109	8,768	84.52	179	5,909	84.33
May	73	8,058	41.89	96	8,556	87.04	169	6,614	39.1
June	65	2,237	84.41	100	8,202	82.92	165	5,520	83.5
July	76	1,775	28.85	116	2,678	23.09	192	4,453	23.1
August	94	2,326	24.74	128	2,985	22.93	222	5,261	23.70
September	71	2,171	80.58	96	2,533	26.38	167	4,704	28.17
October	75	2,495	88.29	90	2,808	81.20	165	5,303	32. 10
November	106	8,284	80.41	189	4,514	82.47	247	7,798	81.5
December	95	8,503	87.98	120	3,720	81.06	215	7,223	83.59
Whole Year	965	82,541	83.72	1,277	88,750	80.84	2,242	71,291	81.80

The average age of all decedents in 1882, was 31.80 years; 0.89 years more than in 1881.

The average age of decedents of American parentage, was 33.72 years, or 0.44 years less than in 1881; of decedents of foreign parentage, the average age in 1882, was 30.34 years, which was 2.26 years more than in 1881, and was the highest average age ever recorded for decedents of foreign parentage, as may be seen by reference to table XV.

In comparing 1882 with the previous year we find the following differences in the several months. Among the decedents of American parentage, there were six months of 1882 when the average age was greater than in 1881; namely: January, February, May, July, August and September; in February and May, the average was very

much greater; in the other six months of the year the average age in 1882 was less than in 1881; in June, November and December, very much less.

Among the decedents of foreign parentage, there were eight months in 1882 when the average age was greater than in 1881; in May and June, October and November, considerably greater; in the remaining four months, the average age was less than in the preceding year.

Table XIV. gives the same statistics for the twenty-seven years, 1856-1882:—

TABLE XIV. DEATHS, TWENTY-SEVEN YEARS, 1856-1882.

Recapitulation.	Aggregate	and Avera	rge Age	by	Parentage.

Twenty-Seven	Amer	ican Par	ent ag e.	Fore	gn Pare	ntage.	Total	Americ a n Foreigu.					
YEARS. 1856–18 82 .	Number of Deaths.	Aggregate Agu.	Average Age.	Number of Deaths.	Aggregate Age.	Average Age.	Number of Deuths.	Aggregate Age.	Average Age.				
Јапиагу	1.704	58,907	84.57	1,578	41,484	26.29	3,282	100,391	80.59				
February	1,558	53,424	84.29	1,472	34,953	23.74	3,030	88,376	29.17				
March	1,607	56,796	35.34	1,516	38,645	25.49	3,123	95,431	30.50				
April	1,563	53,525	34 25	1.607	43,883	27.31	3,170	97,408	80.78				
May	1,500	54,426	36.28	1.451	38,960	26.83	2,951	93,380	31.65				
June	1,243	43,248	84.79	1,372	33,147	25.62	2,615	78,395	29.98				
July	1,696	43,350	25.56	2,068	39,080	18.90	3,764	82,430	21.90				
August	1,843	50,348	27.32	2,181	42,073	19.29	4,024	92,421	22.97				
September	1,587	46,454	29.27	1,818	36,789	20.24	3,405	83,243	21.4				
October	1.478	49.248	33.32	1,637	38,887	23.47	3,135	88,135	28.11				
November	1,537	51,239	33 .34	1,642	42,031	25.60	3,179	93,270	29.34				
December	1,663	60,080	86. 08	1,623	43,518	26 .81	3 288	103,596	31.51				
Totals	1:4,981	021,035	32.72	19,985	475,449	23.79	33,966	1,096.484	28.14				

November, 1882, the average age of American decedents was less than that of the foreign class; this is indicated by prefixing the sign—in the column of "difference" to the figures for those months. In the other months of 1882, and in every month in the long period of years, the average age of the American decedents was greater than that of the foreign.

Difference in Years between the Average Age of American and Foreign

Decedents,

		1862.		. 27 years, 1856-1983.							
MONTHS.	Averag	e Age.		Averag							
	American.	Foreign.	Difference.	American.	Foreign.	Difference					
January	38.69	32.67	6.02	84.57	26.29	8.28					
February	42.47	81.85	10.62	34.29	23.74	10.55					
March	86.08	80.32	5.76	85.84	25.49	9.85					
April	34.06	84.52	-0.46	84.25	27.81	6.94					
Мау	41.89	87.04	4.85	36.28	26.85	9.43					
Jane	34.41	32.92	1.49	84.79	25.62	9.17					
Jaly	23.35	23.69	0.28	25.58	18.90	6 66					
August	24.74	22.98	1.81	27.32	19.29	8.03					
September	30.58	26.38	4.20	29.27	20.24	9.03					
October	33.29	31.20	2.09	83.82	23.47	9.85					
November	80.41	32.47	-2.06	83.84	25.60	7.74					
December 87.98		81.00	6.93	36.08	26 81`	9.27					
Whole Year	83.72	80.31	8 38	32.72	23.79	8.93					

In the month of February the difference between the two classes was greater in the single year than in the long period of years; in all the other months it was greater in the long period of time.

In the year 1882, the largest number of deaths in one month in both classes of decedents, was in November; of Americans, 108; of foreign, 139; the smallest monthly total of American decedents, was 63 in April; of foreign, 89 in February



In the twenty-seven years, 1856–1882, the largest monthly total for both classes was in August; American, 1843; foreign, 2,181; the smallest in June; American, 1,243; foreign, 1,372. The percentages of all the deaths in the different quarters of the year, according to parentage, for the period of twenty-seven years, 1856–1882, were as follows:—

			-
	1st Quarter.	2d Quarter.	3d Quarter. 4th Quarter.
American parentage	25.65	22.68	97.01
Foreign parentage	22.85	22.16	21.50

Table XV. gives the number of decedents, and the aggregate and average age, according to parentage, in each of the twenty-seven years, 1856–1882, inclusive, and for the whole period:—

DEATHS: AGGREGATE AND AVERAGE AGE.

THEADEW YOUR PUBLIC LIBRARY

TABLE XV. DEATHS, 1856-1882.

Recapitulation. Aggregate and average age by parent egeculations.

	Amer	ican Par	entage.	Fore	ign Pare	ntage.	Total American and Foreign.				
YEARS.	Deaths.	Aggregate Years.	Average Years.	Deaths.	Aggregate Years.	Average Yeurs.	Deaths.	Azgregate Years.	Average Years.		
856	541	14,554	26.90	524	8,258	15.75	1,065	22,812	21.4		
857	523	16,314	81.19	402	8,007	19.91	925	24,321	26.		
838	512	15,910	81.07	505	8,478	16.78	1,017	24,388	23.		
839	479	14,993	81.30	420	8,640	20.57	899	23,633	26 .		
860	512	16,660	82.54	489	9,751	19.94	1,001	26,411	26.		
861	538	16,827	30.34	513	9,605	18.72	1,051	25,932	24.		
80 2	460	14,589	81.71	454	10,994	24.21	914	25,583	28.		
3 3	597	19,398	82.49	617	13,757	22 29	1,214	83,155	27.		
864	649	21,732	33.48	632	11,365	17.98	1,281	83,097	25.		
865	631	19,628	31.10	580	11,991	20.67	1,211	31,619	26.		
866	513	18,906	36.85	523	12,762	24.40	1,036	31,668	80.		
367	528	19,258	86.47	432	12,149	2 8.12	960	81,407	82		
968	596	21,779	86.54	514	12,539	24.39	1,110	84,318	30.		
999	627	20,490	32.68	629	14,776	23.49	1,256	85,266	28.		
870	612	21,078	82.83	621	14,393	23.18	1,263	85,471	28.		
871	643	22,725	85.34	611	17,005	27.83	1,254	89,780	81.		
872	832	27,572	83.14	771	19,160	21.88	1,603	46,732	29.		
673	862	28,675	83.27	857	19,061	23.24	1,719	47,736	27.		
874	885	27,588	81.17	1,102	24.802	22.51	1,987	52,390	26.		
875	914	80,419	33.28	1,001	24,302	24.28	1,915	54,721	28.		
876	875	29,025	83.17	990	23,959	24.20	1,865	52,984	28.		
877	908	28,353	31.22	1,030	25,410	24.67	1,938	53,763	27.		
978	889	26,724	30.06	1,100	25,178	22.89	1,989	51,902	26.		
879	941	29,896	81.77	1,085	28,446	26.22	2,026	58,842	28.		
880	920	81,775	84.54	1,160	29,729	25 63	2,080	61,504	29.		
881	999	84,126	34.16	1,148	32,182	28.08	2,145	66.308	30.		
882	965	82,541	88.72	1,277	88,750	80.84	2,242	71,291	81.		
years' total	18,981	621,085	82.72	19,985	475,449	28.79	38,966	1,096,484	28.		

The greatest difference between the average age of the two classes of decedents was 15.50 years in 1864; the least difference, 5.55 years, in 1879. The difference between them in the whole period was 9.33 years.

Table XVI. shows the number and percentage, according to parentage of decedents in different divisions of ages, in 1882, and in the twenty-seven years, 1856-1882 inclusive:—

Table XVI. Deaths, 1882.

Percentage at different ages, according to parentage.

	Amer Paren 180	tage.			enn an eign,	meri- d For- 1882.	27 ye 1856-	
AGES.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Under 1 year	186	19.28	230	18 01	416	18.56	7,540	19.2
1 and under 2	61	6.32	68	5.32	129	5.75	8,312	8.5
3 and under 5	69	7.15	64	5 01	133	5.93	3,560	9.1
Total under 5	816	82.75	863	28.84	678	80.24	14,412	36.
5 and under 10	27	2.80	88	2.98	68	2.90	1,968	5.
0 and under 15	19	1.97	26	2.04	45	2.01	794	2.0
5 and under 20	28	2.90	71	5.56	99	4.42	1,301	3.
0 and under 30	86	8.91	186	14.57	272	12.18	8,988	10.
0 and under 40	92	9.58	167	18.08	259	11.55	8,571	9
0 and under 50	77	7.98	110	8.61	187	8.84	2.971	7.
o and under 60	74	7 67	96	7.52	170	7.56	2 837	7.
0 and under 70	95	9.84	86	6.78	181	8.07	2,810	7.5
0 and under 80	82	8.50	95	7.44	177	7.89	2,650	6.
0 and under 90	55	5.70	88	2.58	88	8.98	1,890	8.
0 and over	14	1.45	7	0.55	21	0.94	284	
Total	965	100.00	1,977	100.00	2,242	100.00	38,966	100.0

The percentage of decedents of American parentage under five years of age, was 1.08 less in 1882 than in 1881; the percentage of decedents between 5 and 10 years of age, was also less than in the preceding year; but from 10 to 50 years of age the percentage was 5.46 larger in 1882 than in 1881. Of the decedents of fifty years and over, the percentage in 1882, was 2.78 less than in 1881; this accords with the fact stated under table XIII., that the average age of American decedents in 1882, was less than in the preceding year. Of the decedents of foreign parentage in 1882, the percentage under five years of age, was 5.61 less than in 1881; the percentage between 5 and 10 years, was 1.12 less than in the preceding year. From 10 to 50 years of age the percentage in 1882 was 6.42 more than in 1881; of fifty years and over the percentage in 1882 was 0.31 more than in the preceding year. The average age of all foreign decedents was greater in 1882 than in 1881.

The following statement gives the percentage, according to parentage, of the decedents under five years old, in each of the twenty-seven years, 1856-1882, inclusive:—

•	American. Under five years.	Foreign. Total Under five years.	American & Foreign. Under five years.
856	•	54.95 per cent	-
1857	32.31 per cent	51.74 per cent	40.75 per cent
		55.44 per cent	
1859	28.60 per cent	48.84 per cent	37.82 per cent
		48.87 per cent	
		53.63 per cent	
		42.07 per cent	
1863	28.47 per cent	46.84 per cent	
		51.11 per cent	
		46.37 per cent	
		40 53 per cent	
		36.35 per cent	
		43.19 per cent	
		43.24 per cent	
		43.67 per cent	
		36.01 per cent	
		40.00 per cent	
		43.05 per cent	
		43.55 per cent	
		40 96 per cent	
		43.13 per cent	
		41.26 per cont	
		42.64 per cent	
		37.24 per cent	
		36.21 per cent	
		33.95 per cent	
	32.75 per cent		

In the forty-three years, 1840-1882 inclusive, there were 52,111 decedents in Providence, of whom 20,147 were less than five years old; the number and percentage in each division of age under five years were as follows:—

Whole number of decedents	8	12,1	11
Under one year	5,095	or	9.78 per cent. of all
Total under five years	20,147	OF	38.66 per cent. of all.

Table XVII. shows the number and percentage of decedents in different divisions of age, according to parentage, in the aggregate for the twenty-seven years, 1856-1882 inclusive:—

TABLE XVII. DEATHS, 1856-1882.

Number and percentage at different ages, according to parentage.

		Parentage 1856-1882.		Parentage, 1866–1882.	I'r MICHINE	1 & Foreig 12, 27 year: 1882.
AGES.	Number of Deaths.	Percentage in each division of age.	Number of Deaths.	Percentage in ench division of age.	Number of Deaths.	Percentage in each division of age.
Under 1 year	8,845	17.62	4,195	20.99	7,540	19.35
1 and under 2	1,280	6.48	2,082	10.42	8,312	8.50
3 and nuder 5	1,465	7.72	2,095	10.48	8,500	9.14
Total under 5	6,040	81.82	8,872	41.89	14,413	86.99
5 and under 10	958	5.05	1,010	5.05	1,968	5.05
10 and under 15	410	2.16	884	1.92	794	2.04
15 and under 20	637	8.86	661	8.33	1,301	3.34
00 and under 80	1,718	9.05	2,270	11.36	8,968	10.23
80 and under 40	1,651	8.70	1,920	9.61	8,571	9.16
10 and under 50	1,436	7.57	1,535	7.68	2,971	7.63
0 and under co	1,540	8.11	1,297	6.49	2,837	7.59
0 and under 70	1,650	8.69	1,160	5.80	2,810	7.21
o and under 80	1,787	9.15	913	4.57	2,650	6.80
30 and under 90	1,019	5.87	8/1	1.81	1,880	3.54
0 and over	185	.97	99	.50	284	.78
Totals	18,981	100.00	19,985	100.00	38,906	100.60

The percentage of American decedents under five years of age in 1882, was less than it had been since 1875; and the percentage of foreign decedents under five years old was much smaller than was ever before recorded, and was 4.41 per cent. less than the same class of American decedents; a thing which has never occurred before in the twenty-seven years in which the records have been kept. may see from table XVII. how much the foreign class exceeds the American as a general rule. In the twenty-seven years, 1856-1882, the number of foreign decedents under five years of age, was 2,332 more than the American, and the percentage 10.07 more. The percentages of American decedents of advanced age are much larger than those of the foreign class; the highest percentage of American decedents, beyond five years, was 9.15 between 70 and 80 years old; while the highest percentage of decedents of foreign parentage was 11.36 between 20 and 30 years old. The number of American decedents living beyond the age of 50 years, was 6,131; the percentage 32.29; the number of foreign, 3,830, and the percentage 19.17. This agrees with the facts shown by tables XIII. and XIV., that the average age of American decedents is greater than that of the foreign; in the twenty-seven years it was 8.93 years more.

The proportion of children of American parentage under 5 years old who died in 1882, (Table XVI.) was 0.93 more than the percentage of the same class for the twenty-seven years, 1856–1882, (Table XVII.); the proportion of American decedents of 60 years and over in 1882, was 1.31 more than in the term of years. Of foreign decedents the proportion under 5 years old in 1882, was 13.55 less than in the period of 27 years; the proportion of those over 60 years of age in 1882, was 4.62 more.

Deaths by age and parentage in each month and quarter of the year. Table XVIII. gives the number of deaths at each age, according to parentage, in each month and quarter of the year, thus showing the comparative mortality in the two classes of parentage. The whole number of deaths was 2,242; 965 of American, and 1,277 of foreign parentage; these occurred in the different quarters as follows:—

	1st Quarter.	2d Quarter.	8d Quarter. 4	th Quarter.
American Parentage	25.89	20.83	21.97	28.61
Foreign Parentage	22.16	23.88	26.63	27.23

Among both classes of decedents the largest percentage was in the fourth quarter of the year, owing to the epidemic of typhoid fever which prevailed in the city at that season.

There were 186 children of American parentage under one year old, who died in 1882; this was 20 less than in 1881, and was 19.27 per cent. of all the decedents of American parentage; of these 186 children, 83, or 43.01 per cent. died in the third quarter of the year; the whole number of American decedents in the third quarter, was 241; the 83 children under one year old made 34.44 per cent. of this number.

Of decedents of foreign parentage, the number under one year, was 230; 21 more than in 1881, and 18.01 per cent. of all foreign decedents. Of these 230 children, 116, or 50.43 per cent. died in the third quarter of the year; and these 116 were 67.65 per cent. of all the foreign decedents in the third quarter.

Causes of death, sex. age, parentage, and percentage from known causes. Table XIX. (pages 52-58) gives the number of deaths in 1882 from each cause, in each division of age; also the total number from each cause, according to sex and parentage, and the percentage from each cause of all from known causes.

Season and disease. Table XX. (pages 59-60) gives the number of deaths from certain prominent causes, in each month and quarter of the year, showing the comparative mortality from different diseases at different seasons.

TABLE XVIII. Deaths by age and parentage in each month and quarter, 1882.

T	INTEINE OF THE				DIV	1810	NAL	PE	RIOI	08 O	F LI	FE.				_	
L	YEAR.	Under 1.	1 to 3.	2 to 5.	5 to 10.	10 to 15.	15 to 20.	20 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.	80 to 90.	90 to 100.	100 & OV'T	All ages.
	January February March	11 10 11	9 3 6	8 7 2	9	8 2 1	1 2 2	2 8 8		4	11 8 4	10 10 6	10	6	3		86 85 71
	lst quarter	8:3	18	17	4	-6	5	18	28	18	23	26	27	16	6	1	245
	April	10 13	6 6 1	6 2 4		₂	8 1 1	7 7 2	6 8 9	4 7 6	6 6 4	6 12 7	4 6 9	4 6 9	2		68 78 66
L'ARENTAGE.	2d quarter	31	13	13	6	8	5	16	23	17	16	25	19	12	3	• •	201
	July August September	25 37 21	8 6 5	12 5 3	 2 1 —	i	1 3 	2 6 7			6 4 4	2 5 9	3 7 9	. 8	2	-: -:	76 94 71
AMERICAN	& quarter	83	19	20	8	2	4	15	17	18	14	16	13	16	2		241
₹	October November December	12 12 16	7 22	5 7 8	4 6 4	1 5 2	2 11 1	4 25 8	9	9 6 9	6 6 9	8 8 12	9 6 9		1		75 108 95
	4th quarter	40	11	20	14	8	14	87	24	21	21	28	21	11	2		278
	Whole year	186	61	69	27	19	28	86	92	77	74	95	82	55		1	965
	January February March	19 10 16	3 3 10	7 6 6	3 3	2 1 2	7 5 3	15 19 9	13 9 17		9 10 8	10 7 2	7		1		99 89 95
	1st quarter	38	16	19	6	5	15	43	39	28	27	19	20	7	1		283
AGE	April May June	14 17 10	1 4 4	5 5 6	3 2	 1 2	5 4 6	15 10 23	23 11 13	16 5 8	10 3 9	9 11 8	7 15 5	6 3	1		109 96 100
PAHENTAGE	2d quarter	4.3	9	16	8	8	15	18	47	29	22	28	27	10	2		305
IGN PA	July August September	45 45 26	6 10 7	5 4	1 5 2	4 2 3	2 4 5	10 16 14	10 9 12	12 7	8 10 5	5 4 5	4 8 6	3 2 3	2		116 128 96
PORRIGA	3d quarter	116	23	10	8	9	11	40	31	26	23	14	18	8	3		340
	October November December	8 12 15	6 7	7 6 6	4 5 7	2 5 2	6 13 11	12 27 16	13 20 17	9 10 8	5 9 10	11 10	10 13 7	2	 1		90 139 120
	4th quarter	35	20	19	16	9	30	55	50	27	24	25	30	8			349
	Whole year	230	68	64	38	26	71	186	167	110	96	86	95	33	7	• •	1,277
				_	-								-				

Ş

독 **경**

Riffo of degins from each cause to whole num-ber from known outseas. .IntoT Poreign. А тогісап han 00 .10vo .08 n 1 0% .08 ot 07 DIVISIONAL PERIODS OF LIFE, .07 01 07 .00 to 00 40 10 20 30 to 40. 20 (0 80° 10 10 20. 10 to 12. .01 of a .6 of 2 Under I Year, burd under 3, Total. Гетвіса. Males. : Drowning..... (various)..... Aneurism of Aorts..... Asphyxia..... Poisoning.... Brain, Diseases of..... Railroad Burns and Scalds..... Ansmis Congettion of..... CAUSES OF DEATH. A poplexy..... Bladder, Diseases of..... Shooting Asthma..... Boncs, Discase of..... Ac.idents,

Disease, sex, age, parentage, and percentage from known causes. DEATHS, 1882. TABLE XIX.

TABLE XIX.—CONTINUED.	DEATHS,	THS,	1004	ı					ļ	١	l	۱	l							
		SEX.				ä	DIVISIONAL PERIODS OF LIFE	DNAL	PE	HOD	0 0	Li	pi				PARENTAGE.	MTAG	1	SOUR: SOUR: ONIOU
CAUSES OF DEATH.	Mules.	Females.	Total.	TashaU year. band	2 to 5.	.et o1 &	10 101	12 to 30.	20 to 80.	.04 o1 08	40 to 50.	.09 01 03	.07 03 08	.03 01 08	— Бия 06	.7970	American.	Foreign.	Total	Ratio of the front enem of the front of the front of the front in the
Bruin, luftammation of	~	ង	\$	=	 #	 -	6	84	*	-	<u> </u>	╎╌	H	<u> :</u> :	<u> :</u>	<u> </u>	12	2	18	8.9
Bronchitta	22	2	8	Ħ	-		<u>:</u>	:	_	- 4	÷	:	8	65	10	_	7	28	8	9.68
Cancer, (various)	6	77	83	$\dot{\vdots}$	÷	<u>:</u>	<u>:</u>	<u>:</u>	:	4	-	*	4	-	÷	•	7	a	23	1.30
of Breast	:	2	2	$\frac{\cdot}{!}$	÷	<u>.</u>	<u>:</u>	Ī	:	A	-	••	- 64	_	÷	:	6	4	ă	3 .
of Stomach	-	-65	80	:	<u>:</u>	<u>:</u>	:	Ī		:	-	60	64	_	-	-:	 0-	10	40	8,
of Uterus		F	Ħ	:	$\frac{\cdot}{\cdot}$	-	<u>:</u>	i	:	-	-6-	:	- <u>:</u> -	<u>:</u>	÷	•	æ	•	Ħ	ą.
Carbancle	Ħ	:	Ä	<u>:</u>	<u>:</u>	÷	<u>:</u>		=	÷	$\frac{\cdot}{\cdot}$	÷	÷	:	<u>:</u>	<u>.</u>	:	_	_	ş
Catarrh		A	F	$\frac{\cdot}{\cdot}$	<u>:</u>	<u>:</u>	<u>.</u>	i	:	- ÷	:	÷	÷	-:	-	<u>:</u>	:	_	_	ğ
Cellulitis Pelvic	:		F	+	<u>:</u>	<u>:</u>	:	_:	÷	÷	\div	-:-	÷	\vdots	÷	<u>:</u>	:	_	-	ş
Cbild-birth	:	-	-	<u>:</u>	<u>:</u>	-	<u>:</u>	:	45	61	\vdots	÷	\div	\div	<u>:</u>	<u>.</u>		6	۲۰	.81
Pucrperal Convulsions.	:	•	64	<u>:</u>	<u>:</u>	÷	ij	:	सं	÷	$\frac{\cdot}{\cdot}$	÷	<u>:</u>	:	÷	÷	. :	-		8
Puerperal Fever	:	20	8	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	_	=	40	-	$\frac{\cdot}{\cdot}$:	<u>:</u>	÷	<u>-</u> :	2-	=	82	16.
Cholera Infantum	8	3	130	801	8	- 65		i	÷	÷	:	÷	÷	<u>:</u>	:	-:	8	8	130	5.8
Morbus	16	60	8	÷	<u>:</u>	<u>:</u>	_		$\frac{\cdot}{\cdot}$:	_	-65	10	_	-	<u>:</u>	64	Ξ	13	2 2
Colle	81	_	60	÷	:	<u>:</u>	:	–	÷	÷	\div		<u>:</u>	:	<u>:</u>	÷	:	66	. 65	.13
••••••••••••••		4	=	<u>.</u>	<u>:</u>	<u>.</u>	<u>:</u>	Ī	÷	÷	:	:	_	- <u>:</u> -	÷	-:	- 61	6 4	-	Ħ
Consumption	108	200	â	7	- 		-	3	116	2	3	8 2	2	-6	61		118	3 2	28	15.73

		SEX.					DIVI	DIVISIONAL PERIODS OF LIFE.	AL.	ER	80	8	L L	١.		1	7	PARENTAGE.	AOR.	OHII R	U MOI
CAUSES OF DEATH.	Males.	Females.	.faioT	Year.	l ander 2.	5 to 8.	.01 O1 8	10 to 12.	12 to 30.	30 to 40.	40 (0 0) 0)	.00 to 60.	.07 01 03	.08 01 07	.08 01 08	79VO	American.	Foreign.	-fatoT	ob to olish odose mori e elodive of	ber from kr
Convulsions	व	32	3	80	100	<u> </u>	╁┿	H	H	<u> -</u>	:-	 i	 :	:	<u> </u>	Ŀ	3	3		12	25.5
Croup	#	13	23	4	6	16	6	÷	\div	$\frac{\cdot}{\cdot}$	<u> </u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>		<u> </u>	7	8	- 68	1.43
Debility	21	4	18	2	÷	÷	$\frac{\cdot}{\cdot}$	÷	÷	:	-:	:	:	<u>:</u>	<u>:</u>		<u>×</u>	_	_	<u>e</u>	٣.
" Birth Premature	80	4	18	2	Ė	÷	÷	$\dot{\div}$	-	<u>:</u>	÷	÷	<u>:</u>	<u>:</u>	<u>:</u>				_	<u> </u>	Ę
Diabeten	4	0		:	i	÷	:	÷	-:	_	_	_	_		-				-	-	8.
Diarrhoa	2	18	**	23	- 6	:	÷	<u>:</u>	$\frac{\cdot}{\cdot}$	-	$\frac{\cdot}{\cdot}$		<u>:</u>			<u>:</u>	=	64	<u>~</u>	22	1.70
" Chronic		61	- 25	_	 -	÷	÷	\div	<u>:</u>	\div	\div	<u>.</u>	<u>:</u>	<u>:</u>	<u>:</u>					-	8 1.
Diphtheria	a	8	\$	65	=	8	4		-	<u> </u>		÷	<u>:</u>	<u>:</u>	<u>:</u>		*	ଅ 	_	<u>86</u>	2.15
Dropsy	:	9	•	Ť	÷	÷	÷	:	-	·		-	-	_	≟		•	Ĭ	_	- 65	2 :
" of Chest	:	-		:	:	:	<u>:</u>	÷	\div	-:	=	÷	<u>:</u>	<u>:</u>	<u>:</u>		_			_	Ş
Dysentery	=	2	2		45	÷	÷	\div	:	_	64	91	65	*			2	=	=	_	ġ
Typhoid	64	÷	01	÷	÷	÷	·	$\stackrel{:}{:}$	•	<u>-</u> :-	÷	÷	<u>:</u>	<u>:</u>	_	<u>:</u>			-	-	8
Krabolism	_	_	01	i	÷	÷	$\frac{\cdot}{\cdot}$	÷	:	<u>:</u>	<u>:</u>	-	<u> </u>	<u>:</u>	<u>:</u>		_			- 64	S.
Emphysema	i	-		÷	:	÷		$\frac{\cdot}{\cdot}$	-	\div	-:	<u> </u>	<u>:</u>	<u>:</u>	<u>:</u>		<u>:</u>			_	ş
Enteritis	81	9	2		_		61	<u> </u>	**	_	-	æ	- 6 1	-			<u> </u>	Ť		-6-	ž
" Gastro	=	-	9	-	•	<i>6</i> 4	=	$\frac{\cdot}{\cdot}$	-;	-		<u>:</u>	ൎ	<u>:</u>	<u>:</u>	<u>:</u>	- -		_	-a-	ĸ
Enllengy	7	•	-	_	_	-	-	-	-												

		SEX.					ă	DIVISIONAL PERIODS OF	IVNC	L PE	RIO	80	F L	Lipe.				<u>~</u>	PARENTAGE	rage.	ire ade
CAUSES OF DEATH.	Males.	Females.	Total.	Under 1	l and er S.	.č oj 2	5 to 10.	10 to 15.	12 to 30.	.08 of 02	.04 01 08	.00 01 04	.00 ot 08	.07 04 08	.08 01 07	.08 of 08	Das 08	оует. Атаетісап.	Foreign.	Total.	Ratio of His of the control of the
Erysipelas	•	٦	*		<u> </u>	<u> :</u>	<u> </u>	<u> </u>	<u> </u>	_	_	"	<u> </u>			:	<u> </u>		04	12	12
Fever, Malarial	_	-84	65			_			<u>:</u>	<u>:</u>	<u>:</u>				<u> </u>	$\stackrel{:}{\div}$	<u>:</u>		- 64	_	**
" Remittent	_		_	<u>:</u>		:		<u>.</u>	<u>:</u>	_		i	:	_:	<u>:</u>	\div	<u>:</u>		=	-	_
" Typhoid	8	7.	140		_		-	8 18	8	8	왏	00			8	ᆖ	<u>:</u>	-	<u></u>	-i -88	140
" Typho-Malarial	_	:		<u>:</u>						<u>:</u>	_	<u>:</u>	<u>.</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>-:</u>	_	
Gall Duct, Inflammation of	:					<u>.</u>					:					÷	<u>:</u>		-:	_	
Gangrene	_	-	64							<u>:</u>	_	_	<u>:</u>	<u>:</u>	_:_	<u>:</u>	<u>.</u>			-	<i>6</i> 4
" Sentle	:	_	_	<u>:</u>		≟	: -: -	<u>:</u>				<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u> -	<u>.</u>	<u>:</u>		-	-	_
Glottis, Spasm of	i	-		_	<u>:</u>	≟			<u>:</u>			<u>:</u>		≟	÷	\div	<u>:</u>		-	-:	_
Heart, Diseases of	-8	F	138	<u>:</u>			•	- 8 -	_	2	7	1	18	ି ଶ		28	:		72.	- 20	28
" Hypertrophy of	•	•	R	≟					<u>:</u>		•				65	<u>:</u>	\div		-	- o -	9
Heat	<u> </u>			<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>				<u>:</u>	_	<u>:</u>	<u>:</u>	<u>.</u>	÷	<u>:</u>	<u>:</u>	.	<u>-</u>	_
Hemorrhage	7	8	•	_	<u>:</u>	<u>:</u>	<u>:</u>			<u>.</u>	<u>:</u>		<u>.</u>		≟	÷	\div		4	- ₀₄ -	9
# from Lungs	69	-	_		<u>:</u>	<u>:</u>	<u>:</u>			_	<u>:</u>		<u>:</u>	<u>:</u>		≟	÷		<i>6</i> 4	_04_	-
	10	64	_		<u>:</u>	<u>:</u>	<u>:</u>	_:_	<u>:</u>	<u>:</u>	<u> </u>		_	_	_		- 69		-	_ ₆₀ _	-1-
Jo Disease of			_			<u>.</u>	_				∔		<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	÷	:	_ <u>.</u>		_
L. ing Cough.	16	2	\$	ង						<u>.</u>			<u>:</u>	<u>:</u>	<u>:</u>	÷	÷		8	8	2
a ocephalus	27	•	33		- 	-85			_;				_ <u>;</u>	_ <u>:</u>	_⋮	_:	_:		-	=	88

edia seni	Total. Ratio of definors around to whole in both in betting to be the manual to be the manual to be the best of t	13.	38.	10.	# ·	•	1.08	<u>s</u>	3.26	6 0.	# T	2	81:	\$	\$	9	1.74	- T		
PARENTAGE	Foreign.	1	#	2	-	•	2	_	=	_	6	Ē		_		•	3	-4	•	•
PAI	American.	1	=	•	:	-	-	:	2	_	_	•	64	:	•	-	2			- 2
	90 and Over.	Ī:	:	:	:	i	:	•	:	:	:	:	:	:	i	-:	:			
	.08 01 08	T:	-	:	:	.:	_	:	:	:	:	:	:		:	:	:			
	.08 o1 o7	1:	:	-	:	•	64	:	*	:	:	æ	_	$\overline{\vdots}$:	:	_			:
<u>#</u>	.07 03 00	Ì	4	90	:	:	-	:	**	:	-	_	~	:	-	:	:		- :	
=	50 to 60.	1:	e 1	8		_		:	4	:	:	•	i	:	:	:	:		-	
DIVISIONAL PERIODS OF LIFE	.08 01 04	1:	•	**	_	:	-		64	_	:	٥	=	_:		-:	:			_
RIOI	30 to 40.		1	20	64	_:	-	_	a			•		:	:	:	:		7	- #
PE	.08 o3 0g	1:	:	•		1	•	:	•	:	:	:	:	:	:	:	•		:	
MAL	15 to 20.		:	:	:	:	:	:	:	_	:	_	:	:	:	_ :				_
181	10 to 15.		:		:	<u>:</u>	:	<u>:</u>	:	:	_:_	:		:	:	:		:	_	:
Ā	.01 of 8	1:				_:_		<u>:</u>	:	<u>:</u>			<u>:</u>	:	<u>:</u>	:	_:		_	:
ŀ	3 to 5.	1:		<u>:</u>	<u>:</u>		:		<u>:</u>	<u>:</u>	_	:	<u>:</u>	_:	<u>:</u>	:	_	*	•	_:
	year. baa I s rebau	L	:			<u>:</u>	_	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>		:	<u>:</u>	•	<u>:</u>	:	_:
_	Under I					-	i	<u>:</u>					_:.	:	_:		2	_:		
	Total	1	19	10			51		ě			<u> </u>	•	_	_		2	_	=	
SEX.	Females.		00	10	-	•	11	-	13	:	1	a			1	•	92	•	•	•
	Males.	1	*	20	60	1	12	-	15	a	64	12	- ON .		i	40	3	:	•	
	CAUSES OF DEATH.	Inanition	Insanity	Intemperance	" Delirium Tremens	Intussusception	Kidneys, Diseases of	" Addison's Disease of	" Bright's Disease of	Knee-Joint, Disease of	Laryngitis	Liver, Diseases of	" Inflammation of	Lungs, Disease of	Lymphadenitis	Malformations, (all kinds)	Maraemone	Measles	Meningitis, Cerebro-Spinal	# Spinsl


	4 2	SEX.				4	NVIS	DIVISIONAL	I P	PERIODS	90	OF L	LIFE.				PA	PARENTAGF	. dop.	+dtg មល មល
OAUSES OF DEATH.	maics.	Females.	.laioT	Under 1 year. I and	under 2.	\$ 10 5.	5 to 10.	15 to 20.	-00 to 30.	30 to 40.	40 10 20T	.09 01 08	.07 01 08	.08 of 07	.08 o1 08	90 and Asyo	American.	Foreign.	Total.	เมตาที่ เพลา
	 	-	İ	-	H	 :	<u> </u>	 	<u> </u>	Ľ	<u> </u>	<u> </u>	:	1		1				
	64	:	- 01	-:	÷	<u>:</u>	÷	÷	-		_	<u>:</u>	:	:	:	:	7	_	64	
Negleot and Starvation	_		A	-:-	$\frac{\cdot}{\cdot}$	$\frac{\cdot}{\cdot}$:	-	<u>:</u>	<u>:</u>	<u>.</u>	<u>:</u>	:	:	_;`	\equiv	-		64	
	=	:	÷	÷	\div	: -	- :-	\div	<u>.</u> .	<u>:</u>	<u>:</u>	_	_:	:	Ī			_		
	81	8	2	÷	$\dot{}$	$\frac{\cdot}{\cdot}$:	<u>:</u>	<u>:</u>	<u>:</u>		<u>:</u>	•	12	7	14	9	8	92	
	13	13	8	÷	-:	<u> </u>	\div	÷	<u>:</u>	<u> </u>	~	***	*	-	91	:	13	~	ม	
Locomotor Ataxla	_	69	*	:	\div	<u>:</u>	:	÷	<u>:</u>	<u>:</u>		~	<u>:</u>	_	i		64		•	
	8	æ	11	=	:	:	:	÷	_	<u> </u>		:_	*	:	:	:	80		17	
Picarisy	-	64	*	÷	$\frac{\cdot}{\cdot}$	•	÷	÷	<u>:</u>		<u>:</u>	_	_	:	_	:	Ø9	_	*	
Pneumonia	8	7	170	91	2	a	64	~	7 10	98	2	8	71	g	80	7	8	107	571	
Congestion of Lungs	ca .	च	9	69	<u>:</u>	÷	$\frac{\cdot}{\cdot}$	<u>:</u>	•		<u>:</u>	<u>:</u>	*	:	:	:	8	w	_	
Prostate, Disease of	*	:	ø	÷	<u>:</u> -	÷	<u>:</u>	÷	<u>:</u>	<u>:</u> -	<u>:</u>	_ <u>:</u> _	:	:	8	:	•			
Purpura Hemorrhagica	:	ø	85	-	<u>-</u> :-	:	:	÷	•			<u>:</u>		:	:	:	G9			· •
	:	H	÷	:		:	- :	<u>:</u>	<u>:</u>	- -	<u>:</u>	. :	:	:	:	:	-			
	_∞	ю	œ.	÷	\vdots	:	\div	:	.	_	<u> </u>	-	-	1	:	:	64	₩.	•	
	_연	9	38	_	64	9		<u>:</u>	:	_:		<u>:</u>	:	•	:	:	7	=	81	-24
	~~	-	*	÷	$\frac{\cdot}{\cdot}$	÷	:		:		_	_ :_	:	•	:	:	:		•	
	÷	:	7	=	$\frac{\cdot}{:}$	÷	÷	÷	<u>:</u>	<u>:</u>	<u>:</u>	:_	:	:	:	:	:			
Spine, Diseases of	•	7	÷	=	_ <u>:</u>	_	_	_			_			:			-	_	_	

TABLE XIX.—CONTINUED.	DEA	DEATHS, 1882.	1887		اڅ	ase	3	'n,	ğ	Pag.	nta	ž	and	2	Cen	tage	Disease, sex, age, parentage, and percentage from	# k	nom	known causes.	3
		SEX.				-	Ĕ	DIVISIONAL PERIODS OF LIFE	IL P	ERIO	8	I I	1				PAI	PARENTAGE.	O.E.	·un	UMO
CAUSES OF DEATH.	Males.	Females.	Total.	Under 1 year.	l and under 2.	'g 03 B	.01 o1 d	10 to 15.	20 to 80.	.01 to 40.	.02 of 0	.08 63 05	.07 03 08	.08 of 07	,08 of 08	1940 90 mpd	.naoriean.	Foreign.	.faioT	LOM BRODE I	engses.
Spine, Pott's Disease of	[18	İ	Ħ	†:	 	╎╌	=	 	Ļ	<u> </u>	1 :		T:	1:	1:	1 84	. "		8
Stomach, Diseases of	4	-05	90		:	- -	<u>:</u>	- <u>:</u>							:	<u> </u>	:	60	~		Ą
Stomach, Inflammation of	4	11	19	-;-	-	:	-	-		- 64	<u> </u>	4		- 84	_	:	*	11	*		۶.
Saloide	8	⊸.	18	-	÷		÷		<u> </u>	_	-	_		~	:	-	•	4	×		ş
By neope	-	:	_	Ť	:	_	÷	:	÷	<u>:</u>	<u>:</u>	<u>:</u>			:	:	=	:	_		કં
Syphilis	~~	:	64	÷	\div	-:	$\frac{\cdot}{\cdot}$	-	-:	:	_	:			:	:	64		•		8
" Congenital	- 64	6	-	4	-	-4-	$\frac{1}{1}$	$\frac{\cdot}{\cdot}$		<u>:</u>					-	-	8				₹.
Tabes Mesenterica		**	4		ei	:	=	- :-	-	\div			_:	:		i	-	•	Ĭ		=
Teething	2	•	18	2	-¢-	-:	$\frac{\cdot}{\cdot}$	- <u>:</u> -	<u>:</u>	<u>:</u>		<u>:</u>			:	:	a	66	×		٤
Tetanus and Tris. Nascentium		_	æ	÷	÷	_ <u>:</u>	-	- :	$\stackrel{:}{-}$	_ <u>:</u>			<u>:</u>		:	:	94		•		8
Tuberculosis	-4-	- 63	51	- 84	=	$\frac{\cdot}{\cdot}$	- :		_	60	_	_		•		-:	•	•	×	•	Ę
Tumor	- eo	60	_ •	:			-		÷	—.		_	:	_		:	65	60	Ĭ		5
Ucer		:		÷	- :	:	÷	<u>:</u>	-	_ <u>:</u>	<u>:</u>	_;	:		_	-	-				કું
Uremia	_	:	_	<u>:</u>	:	$\frac{\cdot}{\cdot}$	÷	-	:			:			:	-	:				3
Uterus, Disease of		•	- 🕶		- :	$\frac{\cdot}{\cdot}$		-	$-\frac{\mathbf{i}}{\mathbf{i}}$	_:			:			_	_				=
Unknown	**	•	a	-		-:	-61	-	-:			_:			:						2
Budden		:		÷	:	- <u>-</u> i-	<u>-</u> :	-	÷	<u>-</u> :-	<u>:</u>	_:_		<u>:</u>		-					:
Totale	1				il	1	1	듺		H	1				1	+		T			1

	l		ŀ														
CAUSES OF DEATH.	January.	Pebruary.	March.	First Quarter.	Apdl.	71ay.	, .enut	Second Quarter.	July.	August.	September.	Third Quarter.	.табо1эО	Хочетрег.	ресешрет.	Гопті) Фантет.	Whole Year.
	20	9	8	54	•	1	۽	2	l S		1	22	<u> </u>	·	8	8	F
A poplexy	1-	1-	*	Ä	65	•	-	7	9	*	91	æ	Ξ	13	13	87	28
:	۵	10	-	5	-6	-	80	8	4	-	4	15	-	*	2	19	55
	-	F-	-	19	*	8	•	16	80	64	••	80	64	80	1-	17	8
:	63	8	60	12	8	83	•	15	Oł.	93	•	2	40	*	8	15	55
:	•	+	:	=	4	_	Ø,	7	:	-	i		64	64	4	80	25
<u>:</u>	<u>:</u>	÷	:	Ä		_	*	10	7	23	18	116	1-	Ä	• •	30	130
:	:	:	÷	:		:	:	:	9	4	A	흑	i	:	_	_	13
:	ಹ	8	8	10.	3	8	8	å	23	 	19	5	ន	75	31	8	821
	-	65	04	=	63	8 0		2	ń	40	40	13	60	14	64	10	20
:	-0	-	*	9	4	~	:	10	_		:	_	40	80	*	16	25
:	-	_	_	*	:	:	æ	69	6	*	60	8	ø	-oi-		40	7
:	-	60	*	÷	80		64	•	•	_	_	1-	_	-	2	13	\$
<u>:</u>	<u> </u>	,	:	8	•	•	:	<u> </u>	H	æ	4	=	10	<u>-</u>	:	Φ.	g
•	*	8	64	13	69	•	4	13	4	-	z	10	-	8	64	•	9
:	A	_	**	8	4	-	- 24	1-	:		:	i	i		_	-	#
	•	Ø1	85	a	H	•	-0	-6:T	Ī	4	.	8	65	2	88	108	140
	-	:	i	i		:	i	:	:	:	61	61	_	81		8	•
:	2	*	2	52	2	7	18	;	-	1	80	9	22	16	==	5	143
- :	9 0	•	•	13	63	6	*	13	6	8	61	9.	-	69	69	40	\$
:	04	8	4	•	-84	æ	-84	-	_		-	•	-6	_	69	0	*

유 로!

Whole Year. Fourth Quarter December Холешр**ег.** October. Season and Disease. Third Quarter. Septemb**or.** .tsnyuA July. Second Quarter, DEATES, 1882. .ount Any. A prill. TABLE XX.—CONTINUED. First Quarter. yluı.cp. Kebrum⊅. Januar.y. Old Age..... Paralysis..... Stomach, Discases of...... Liver, Diseases of...... Rhcumatism..... Intemperance..... Insunity Preumonia and Congestion of Peritonitis..... Kidneys, Diseases of...... 8. arjatina...... CAUSES OF DEATH. Sulcido.... Меавјев..... Meningitis, Cerebro-Spinal Tuberculosis..... Marasmus..... Cuknown Tect bing.....



NOTES ON THE CAUSES OF DEATH,

DURING THE YEAR 1882.

The following pages comment on the figures in Tables XIX. and XX.; additional statistics are also given here, which could not be presented in those tables; and compared with the statistics of previous years.

The percentage of deaths is always reckoned on the number from known causes.

The whole number of deaths in Providence in 1882 was 2,242, of which there were 10, or 0.45 per cent., for which the cause was not stated; the number from known causes, and on which the percentages are reckoned, was 2,232. The number of which the cause was not given was 5 less than in 1881, and the percentage, 0.25 less.

During the twenty-seven years, 1856–1882, the whole number of deaths was 3×,966; from unknown causes, 575, or 1.48 per cent.; from known causes, 38,391. In the forty-three years, 1840–1882 inclusive, there were 52,111 deaths; 1,577, or 3.03 per cent. from unknown causes; and 50,534 from known causes.

Table XX. shows the prevalence of different diseases at different seasons. Of diarrhoal diseases there is always a much larger number in the third quarter of the year than at any other season; while the diseases of the respiratory system prevail much less in this, part of the year, and are found largely in the first and fourth quarters. This table shows the extent to which typhoid fever prevailed in the last quarter of 1882; 75.71 per cent. of all the deaths from this disease having been in the last three months of the year.

Abscesses. There were 7 deaths from abscesses in 1882, which was 0.31 per cent. of all from known causes. The number was the

same as in 1881; but the per cent., 0.02 less. Of two the location was not specified; the others were as follows; two of the pelvis, and one each of the jaw, neck and spine. The statistics as to sex and parentage are:—

Males, 0; females, 7; total, 7; or 0.81 per cent. of all. American parentage, 2; foreign parentage 5.

For the twenty-seven years, 1856-1882, the figures are:-

Number of deaths from all known causes, 38,391.

From abscesses, 108; or 0.28 per cent. of all.

Males, 51; females, 57.

American parentage, 49; foreign parentage, 59.

Accidents. The number of deaths from accidents in 1882, was larger than ever before; it was 77, or 3.44 per cent. of all; it was 25 more than in 18~1 and 1.00 per cent. more. There were 9 deaths from burns or scalds; 10 from drowning; 22 from falls; in 1881 there were only 8 deaths caused by falls; and in that year all the decedents were over forty years of age, while in 1882 they ranged from two years old to 70 and over; among these accidents the following may be specified; two fell down stairs and broke the neck; one fell on the ice and broke the thigh; one other fractured the femur and two the spine, it is not stated exactly how; one each fell from a coalpocket, a ladder, a staging, the bowsprit of a vessel, a low-gear, and in an elevator well; four fell from a window or some part of a building, besides the four who jumped from the burning building on Calender street, and died either instantly, or from injuries so received.

There were 3 deaths by poisoning; one was from morphine taken habitually; one from an overdose of morphine; and one from an overdose of gin taken for medicine; there were 6 deaths by railroad accidents; and one from wounds by a toy pistol. There were 25 accidents included under the heading various, which were as follows: fracture of the hip, or thigh (probably by a fall), 5; fracture of tibia, 1; injury to spine, injury of urethra, one each; injury at birth, three; thrown from a carriage or cart, 2; run over by cart, 2; caught in elevator-well, one; in shaft, one; in machinery (not specified), one; hit on the head by the iron door of a cupola, by a lever, by a pickaxe, and by a falling coal-pocket, one each; head crushed by a safe, one; kicked by a horse, one; swallowing a safety-pin, one.

The number of each sex and parentage was as follows:-

Males, 54; females, 23; total, 77; or 3.44 per cent. of all. American parentage, 29; foreign parentage, 48.

For the twenty-seven years, 1856-1882:—

Males, 830; females, 285; total, 1,115; or 2 90 per cent. of all.

American parentage, 430; foreign parentage, 685.

Aneurism. There were two deaths from this cause in 1882; one American and one foreign male died of aneurism of the aorta. The statistics for this cause for the twenty-seven years, 1856-1882, are:—

Males, 24; females, 5; total, 29; or 0.08 per cent of all.

American parentage, 11; foreign parentage, 18.

Apoplexy. The number of deaths from this cause has increased very much in the last few years; the number in 1882 was 89, the largest number ever reported, and 3.98 per cent. of all deaths from known causes; the number was 15 and the per cent. 0.51 more than in 1881. The particulars of sex and parentage are as follows:—

Males, 49: females, 40; total, 89; or 3.98 per cent. of all. American parentage, 55; foreign parentage, 34.

For the twenty-seven years, 1856-1882:-

Males, 441; femules, 446; total, 887; or 2.31 per cent. of all.

American parentage, 613; foreign parentage, 274.

Further details are given under the head of "Paralysis."

Brain, Diseases of. The number of deaths reported in 1882 from different diseases of the brain, was 75, or 3.36 per cent. of all; 13 less deaths and 0.78 per cent. less than in 1881. Of these deaths, 11 were specified as "congestion" and 46 as "inflammation of the brain."

The number of each sex and parentage :-

Males, 34; females, 41; total, 75; or 3.36 per cent. of all. American parentage, 38; foreign parentage, 37. For the twenty-seven years, 1856-1882, the statistics are:-

Males, 792; females, 691; total, 1,483; or 3.86 per cent. of all.

American parentage, 772; foreign parentage, 711.

Bronchitis. This was reported as the cause of 60 deaths in 1882; 7 more than in the preceding year. Thirty-four of the decedents were less than five years old; and 23 were over sixty. The particulars of sex and parentage were:—

Males, 21; females, 39; total, 60; or 2.68 per cent. of all.
American parentage, 24; foreign parentage, 36.

The statistics for the twenty-seven years, 1856-1882, are as follows:—

Males, 255; females, 317; total, 572; or 1.49 per cent. of all.
American parentage, 255; foreign parentage, 317.

Cancers. There were 52 deaths caused by cancers in 1882, which was 2.60 per cent. of all; it was 13 less deaths and the rate 0.45 per cent. less than in the preceding year. Of these 52 cases of cancer, 10 were "cancer of the breast;" the same number as in 1881; 8 "cancer of the stomach;" 3 less than in 1881; and 11 "cancer of the uterus;" 5 less than in the preceding year. There were 23 under the head of various; of these, three were not specified and the others were as follows: 5 of the liver; 3 of the face; and one each of the lip and neck, larynx, tongue, abdomen, omentum, bowels, rectum, spleen, ovary, vagina, thyroid gland and inguinal glands.

The following are the numbers for each sex and parentage:-

Males, 14; females, 38; total, 52; or 2.60 per cent. of all.

American parentage, 31; foreign parentage, 21.

For the twenty-seven years, 1856-1882, the numbers are:-

Males, 257; females, 689; total, 946; or 2.46 per cent. of all.

American parentage, 593; foreign parentage, 853.

()f the 946 decedents, 72.83 per cent. were females; and 62.68 per cent. were of American parentage.

Child-birth. The number of deaths from child-birth in 1882 was 27, including 2 from puerperal convulsions, and 18 from puerperal fever. The whole number was 2 less than in 1881, and the per cent. 0.15 less. The 27 deaths were 1.21 per cent. of all deaths from known causes, and 0.98 per cent. of the 2,759 mothers bearing children during the year; the decedents were 8 of American, and 19 of foreign parentage. In the twenty-seven years, 1856–1882, there have been 461 deaths from child-birth; 1.20 per cent. of all deaths, and 0.86 per cent. of the 53,981 mothers bearing children during that time; 193 of the decedents were of American, and 268 of foreign parentage.

Cholera Infantum. There were 130 deaths from this cause in 1882; 28 more than in 1881. There have been but three years, when the number of deaths from this disease was greater than in 1882, but several years when the rate was higher. The particulars of sex and parentage are as follows:—

Males, 66; females, 64; total, 130; or 5.82 per cent. of all.

American parentage, 50; foreign parentage, 80.

The following are the statistics of sex and parentage for each of the twenty-seven years, 1856-1882, and for the whole period:—

	8	EX.	Parknta	GE.	Whole
	Mules.	Females.	American.	Foreign.	Number. Percentage.
1856	. 85	23	16	41	575.54
1857	. 27	23	26	24	505.56
1858	. 27	22	18	86	494.92
1859	. 14	19	18	20	83
1860	. 85	83	26	42	687.09
1861	. 29	82	25	36	616.02
1879	. 25	19	17	27	444.98
1863	. 86	80	19	47	66 5.56
1864	. 29	29	18	40	584.63
1965	. 20	80	13	87	504.21
1866	. 80	17	15	83	474.44
1867	. 24	25	23	26	495.19
868	. 88	82	23	48	70 6.44
1869	. 85	80	27	88	655.24
870	. 48	45	87	58	98
1871	. 80	82	26	36	63
1872	. 72	79	58	93	1519.56
1878	. 42	50	89	58	925.41
1874	. 72	62	44	90	134 6 85
1875	. 62	74	50	86	1367.18
1876	. 66		48	79	122
877	. 70	52	48	79	123 6.31
1878	. 25	85	26	44	70
879	. 80	29	28	81	59
880	. 42	58	41	59	1004.84
1881	. 58	44	85	67	1034.79
1889	. 68	64	50	80	1305.82

Of the 2,140 decedents, 1,847 or 62.94 per cent. were of foreign parentage.

Cholera Morbus. The number of deaths from this cause in 1882, was larger than usual; it was 13, which was 9 more than in 1881; only in the year 1866, has the number been larger than this; in that year it was 17.

The statistics for 1882, are:-

Males, 5; females, 8; total, 13; or 0.58 per cent. of all. American parentage, 2; foreign parentage, 11. In the twenty-seven years, 1856–1882, the whole number of deaths from this cause was 133, or 0.35 per cent. of all.

Consumption. The number of deaths from consumption in 1882, was 351, which was 7 more than in 1881; the rate in 1882 was 15.72, which was 0.43 less than in the preceding year. The particulars of sex and parentage are as follows:—

American parentagemales,	57females,	56total, 118
Poreign parentagemales,	109temales,	129total, 238
•		
Totalsmales,	166females,	185total, 851

For the twenty-seven years, 1856-1882, the statistics are:-

Whole number of deaths from known causes, 38,391; from consumption, 6,385; or 16.63 per cent. of all. Annual average, 236.

american parentagemales, 1,275females, 1,629total, Foreign parentagemales, 1,638females. 1,848total	
otal	6,885
Males, 45.62 per centfemales, 54.38 per cent.	

American parentage, 45.48 per cent......foreign parentage, 54.52 per cent.

The population of the city being estimated at 114,377, the deathrate from consumption in 1882 was one in 326 of the inhabitants, or 3.07 in each thousand persons; in 1881 the rate was one in 312, or 3.20 in each 1,000. The average annual rate for the fifteen years, 1840–1854, was one in 209.9; for the twenty-eight years, 1855–1882, it was one in 307.6.

We see from the figures given above that more than fifty per cent.

If the decedents from consumption are of foreign parentage. The ollowing figures show the percentage of each class of decedents in ach of the last nine years:—

TE NEW YORK PUBLIC LIBRARY.

ASTOR, LENOX AND TILDEN FOUNDATIONS.

	Deaths from	A	D	Tamilan	Dan
	Consumption.	American.	Per cent.	Foreign.	rer cent
1874	270	106	89.26	164	60.74
1875	299	114	88.18	185	61.87
1876	284	190	42.25	164	57.75
1877	291	116	44.23	178	55 78
1878	805	107	85.08	198	64.95
1879	293	116	89.59	177	60.41
1880	823	111	84.47	¥11	65.43
1881	344	116	33.79	228	66.20
1863	851	118	\$3.19	238	67.8
	 2,763				

The following is a statement of the number and percentage, according to parentage, of decedents from consumption, in periods of five years each from 1856 to 1880 inclusive, and for the years 1881 and 1882:—

Am	erican Par	entage.	Foreign Parentage.	Deaths from
;	Nupiber.	Per cont.	Number. Per ce	ent. Consumption.
1856 to 1860	. 525	56.45	40543	55 980
1861 to 1865	. 493	50.77	48048	.23 975
1866 to 1870	. 532	50.57	520	.48 1,052
1871 to 1875	. 553	44.78	689	. 22 1,535
1876 to 1880	. 570	88.05	92861	.95
1881	. 116	83.79	\$28 60	.28 844
1882	. 113	82.19	238 67	.81 351
27 years' total	.294	45.48	8,48154	526,386

In the report for 1881 a full comparison was made of the number of decedents in each class of parentage and the proportion of each to the population of the corresponding class. It showed that the proportion of decedents to the population of American parentage had decreased steadily during the twenty-six years since 1856; the rate in 1856 being one death in 268.2 of the population; and in 1880 one in 435.2. The ratio among the foreign population has changed but little during the same period; in 1856 it was one in 262.6; in 1880 one in 268.0.

For fuller particulars on this subject, see "City Registrar's Report, 1881," pages 63, 64 and 65.

The following table gives the most important particulars in regard to the decedents from consumption in each of the twenty-seven years, 1856-1882 inclusive:—

DEATHS FROM CONSUMPTION IN PROVIDENCE, TWENTY-SEVEN YEARS, 1856-1882.

Sex; parentage; season; age; whole number, and percentage from known causes.

882. Total	851 6,385 5.72 16.63	166 2,913	113 2,r04 238 3,481	
881.188	154	157	116	-2228888888
880.18	322 8	23	===	8222228228
1879. 188	62 15	152	912	SHEEFFEEFE
8.18	305	170 1	198	22222222222
7. 1878.	1 200	167	92	22422222222
3. 1877.	5.3415.	119	150	SERESERIE
.1876.	1 3		7,10	
1875.	299	157	114	2228222222
1874.	970 13 76	95.7	108	8888888888
1873.	231	131	11	218355555
1872.	241	124	12	222222222
1871.	194	25	88	1222222222
1870.	9.08	187	122	SESSITESESS
1569.1	210	93	88 51	121111111111111
868	9.71	98	100	222222222
1.198	189	8.8	8.5	2222222222
866.10	108.	92	88	22122222222
1865. 18	195	ZE	88	8128882778824
198	197	85	107	283128282283E
863. 18	208	22	901	ESESSIES SESSE
1802.18	1 22	8.8	8.8	=========
-	184	28	8.5	22682778327
860, 1861	712	818	88	ESEE#558138
	12.25	23	28	5625655552
8. 1859.	858	13	22	22212210×45
.1858.	102 16.20	9.0	88	
1857	62	-		00747-0-488
1856	1.5	105	10.	E - 2 21 - 30 30-2
CONSUMPTION.	Whole number 178 Percentage 17.31	Srx: Mules Femuke	PARENTAGE: American Foreign	SEASON: January February March April April June June Septiember Septiember

DEATHS FROM CONSUMPTION IN PROVIDENCE, TWENTY-SEVEN YEARS, 1856-1882.—CONTHUED.

causes.
known
e from ka
and percentage
and
age; whole number,
whole
age;
season;
Sex; parentage;
Sex;

Total	266.2	8,9,9,8, 81,93,4,	25 25 25 25 25 25 25 25 25 25 25 25 25 2
	8328	75 E 8	<u> </u>
1881	8588	ន៍ដូន	44881222528508
.088	2322	115 117	444-885-3868su
1879. 11	2832	5. 25 Z.	
1878. 15	222	282	_4_4@\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
	3335	258	4-84-0854684-
76. 1877	8288	8 63	44-89882±885¤
1875. 1876.	2222	<u> 288</u>	# - # # # # # # # # # # # # # # # # # #
187		- 28 8 2 - 28 8 2	Bond-Pirth
3.1874.	22.23	282	
1873.	5555	858	
1879.	8528	583	
		888	######################################
1870	2882		
	8248	6:5	977-658-9864
	2222	58%	F : 8 4 4 4 4 5 5 5 5 5 5 5
1867.	3222	582	49.9-84666
886	3843	582	20040505050
1864. 1865. 1866.	5328	<u> </u>	Bundan Na Callina
- :	2224	538	**************************************
	&848	759	
1862.	8825	323	<u> </u>
—1 1981 —11	2423	288	4-0-050-3450
1800.18	8228	250	504 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
- 35 - 35	2448	858	원전보원소년중 <u>간원</u> 교통원
1638. 1838.	2222	848	
	\$585 \$585	236	#\$\$\$\$\$\$
<mark></mark>	- 3428 -	222	9 :-845°5333ee-
8	 	: 4:	
CONSUMPTION. 1856.18	First quarter Becond quarter Thirl quarter	First 4 months Second 4 months. Third 4 months	Moder 1 year
COM	First (Second	First & Secon Third	AGE Und Und Und Und Und Und Und Und Und Und

Convulsions. There were 52 deaths from convulsions in 1882; 15 more than in 1881. The particulars as to sex and parentage are:—

Males, 24; females, 28; total, 52; or 2.32 per cent. of all. American parentage, 23; foreign parentage, 29.

For the twenty-seven years, 1856-1882:-

Males, 539; females, 447; total, 986; or 2.56 per cent. of all.

American parentage, 412; foreign parentage, 574.

Croup. The number of deaths from croup in 1882 was 32, which was 17 less than in the preceding year; the rate 1.43 or 0.87 less than in 1881. The deaths in the different quarters of the year were as follows:—

1st quarter10	3d quarter 1
2d quarter 5	4th quarter 16

There were 4 decedents under one year old; 6 between one and two years; 16 between two and five; and 6 between five and ten years. The number of each sex and parentage was:—

Males, 14; females, 18; total, 32; or 1.43 per cent. of all.

American parentage, 14; foreign parentage, 18.

For the twenty-seven years, 1856-1882, the numbers are:—

Whole number of deaths, 848; annual average, 31.

Males, 423; females, 425; total, 848; or 2.21 per cent. of all.

American parentage, 345; foreign parentage, 503.

The following table gives full particulars for each of the twenty-seven years, as to sex, age, parentage, season, locality, color and rate:—

DEATHS FROM CROUP IN PROVIDENCE, TWENTY-SEVEN YEARS, 1856-1882.

71. [872, 1873, 1874, 1875, 1876, 1877, 1878, 1879 1880, 188	83 27 21 29 56 n5 40 38 43 29 49 58 1.31 1.28 1.48 2.93 8.48 2.07 1.93 2.13 1.41 2.80	18 13 6 10 81 82 18 18 27 8 24 17 14 15 10 25 83 22 20 16 21 25	20 6 6 15 19 27 14 18 22 7 17 17 18 21 15 14 35 36 20 21 22 32	0000000 00000 00000000 00000 00000000 00000 00000000
1869, 1870, 1871.	1.63 2.16 2.8	10 9 11 9	13 15 2	0)- 04
. 1867. 1868.	7 2.01 1.01	3 10 6	10 10 4	
4.1865.1866.	45 29 23	25 11 11 26 18 13	19 9 11 26 20 12	**************************************
. 1862, 1863, 1864	24 31 33 3. 27 3.51 2.78 3.	11 20 21 12 20 21	9 15 16 15 16 22	10000000
1860, 1861	1.88	היו	- +3	9-94-H H 4 9
57. 1858. 1859.	22 17 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	10 19 18 18 15 6	18 12 12 10 15 12	*************
1856 1857.	Der 252 3.	16	908	#40H-01-01 , Store
CROUP	Whole number	Ex: Males	American	January Jenuary Merch March May June Auril May June Cotober October December

Debility. The number of deaths for which this cause was assigned in 1882, was 28, of which 12 were from premature birth; the number was 6 less than in the preceding year. The statistics of sex and parentage are:—

Males, 20; females, 8; total, 28; or 1.26 per cent. of all. ... American parentage, 17; foreign parentage, 11.

For the twenty-seven years, 1856-1882.

Whole number of deaths, 662; annual average, 25.

Males. 353; females, 309; total, 662; or 1.72 per cent. of all.

Diarrhoa and Dysentery. The number of deaths from these causes in 1882 was 64; 7 more than in 1881; the rate, 2.86 or 0.18 per cent. more than in the preceding year. The number from each disease was:—

	Mules.	Females.	American.	Foreign.	Total.	Per ct.
Acute Diarrhœa	20	18	17	21	38	1.70
Chronic Diarrhoa	1	2			8	0.18
Dysentery	11	10		11	21	0.94
Typhoid Dysentery.	2	• • • • • • • • • • • • • • • • • • • •		2	2	0.09
Total	34	30	80	84	64	2.86

The deaths from other diseases of this same class were as follows:-

	Males.	Females.	American.	Foreign	Total.	Per ct.
Cholera Infantum	68	64	50	80	130	5 82
Cholera Morbus	5	8	2,	11	13	0.58
Enteritis	12	18	16	14	30	1.84
Enteritis Gastro	14	5	12	7	19	85
Total	 97	95	80	112	192	8.59

The total number of deaths from all these causes during the year, was as follows:—

Males, 131; females, 125; total, 256; or 11.45 per cent. of all.

American parentage, 110; foreign parentage, 146.

The whole number is 74 more than in 1881, and the rate 2.90 more, the number of males, 33 more; of females, 41 more; of American parentage, 36 more; of foreign parentage, 38 more.

The statistics of these diseases for the twenty-seven years, 1856-1882, are as follows:—

Number of deaths, 4,218; annual average, 156.

Males, 2,109; females, 2,109; total, 4,218; or 10.99 per cent. of all.

American parentage, 1,719; foreign parentage, 2,499.

The proportions between the two classes of decedents in 1882 were 132.7 decedents of foreign parentage to each 100 of American parentage; or 57.03 of foreign parentage and 42.97 of American parentage in each 100. Of decedents from all causes in 1882 there were 132.3 of foreign to each 100 of American parentage; or 56.96 of foreign, and 43.04 of American parentage in each one hundred.

The following table shows the number of deaths from each of these causes, with the number from the whole class of causes, and their percentage of all from known causes, reported in Providence during the period from 1840 to 1854 inclusive; during each year, from 1855 to 1882 inclusive, and in the aggregate during the whole period of 43 years, 1840 to 1882 inclusive:—

						Per cent.
C	holera Ch	olera				of all from
In	fantum. M	orbus. Di	arrhœa. Dys	entery. Enteri	is. To	otal. knowu cause
iO-1854	560	68	849	796102	1	.87016.
1855	61	4	37			14014.
1856	87	5	29	35 6.		13211.
1857	50	2	38	29 5		12413.4
1858	49	2	27	12 8		98 9.0
1859	83	1	28	18 4		79 8.
1860	68	2	21	18 9		11811.
1861	61	5	22	32 6		12611.
1862	44	1	31	22 9		10711.
1863	66	5	35	77 9		19215.
1864	58	4	45	40 13		16012
1865	50	7	34	62 4		15712
1866	47	17	28	44 9		14518
1867	49	2	15	43 10		11912
1808	70	3	24	15 8	• • • • • • • • • • • • • • • • • • • •	120 10.
1869	65	6	24	23 6	,	124 9
1870	93	3	22	16 15	•••••	14911
1871	62	6	24	16 18		12610
1872	151	10	52	85 14		26216
1878	92	4	87	7 25		163 9
1874	134	3	50	17 25		22911
1875	136	6	86	14 31	•••••	22311
1876	122	4	40	22 18		20611
1677	122	6	84	21 21	•••••	20110
1878	70	3	2 8	11 18	• • • • • • • • • • • • • • • • • • • •	130 6
1879	59	5	81	9 16	•••••	120 5
1880	100	4	30	12 19		165 7.
1881	102	4	43	14 19	•••••	182 8
1682	130	13	41	23 49		25611

Deaths from Asiatic cholera are not included in these statistics; ey have been as follows: in 1849, 158; in 1854, 159; in 1866, 14; tal, 331.

Diphtheria. There were but 48 deaths from this cause in 1882; smaller number than in any year since 1875; it was 68 less than 1881, and was 2.15 per cent. of all deaths from known causes. There were some deaths in each month and they were distributed in different seasons as follows:—

1st quarter14	8d quarter 7
2d quarter 6	4th quarter21

Of the 48 decedents, 45, or 93.75 per cent., were less than ten year old. The number of each sex and parentage was as follows:—

Males, 20; females, 28; total, 48; or 2.15 per cent. of all. American parentage, 26; foreign parentage, 22.

During the four years' epidemic of diphtheria, 1876–1879, inclusive, the decedents of foreign parentage were greatly in excess of those of American parentage, the proportion being 56.39 of foreign, and 43.61 of American parentage in each 100; previous to the epidemic the reverse of this was true, the figures being 58.68 of American, and 41.32 of foreign parentage in each 100 decedents. Since the epidemic the American class is again in excess; in 1880, 57.38 per cent. of American, and 42.62 of foreign parentage in each 100 decedents; in 18×1; 50.86 of American, and 49.14 of foreign parentage in each 100; in 1882, 54.17 of American, and 45.83 of foreign parentage.

The first cases of diphtheria in Providence were reported in 1858: for the twenty-five years, 1858-1882, inclusive, the statistics are as follows:—

Deaths from known causes in 25 years, 36,463.

From diphtheria, 1,318; annual average, 53.

Males, 660; females, 658; total, 1,318; or 3.61 per cent. of all.

American parentage, 647; foreign parentage, 671.

In the following table, the statistics are given as to sex, parentage, season and locality, for each of the twenty-five years, 1858–1882, and for the whole period:—

DEATHS FROM DIPHTHERIA IN PROVIDENCE, TWENTY-FIVE YEARS, 1858-1882.

Врентивкіл. 18	1898.1	828	.0981	1861.	1859, 1860, 1861, 1862,	1863.	1864	1865.1	1896, 1867, 1868, 1869, 1870, 1871, 1872, 1873, 1874, 1875, 1876, 1877, 1878, 1879.	867.1	868.	1860	.0281	1871.	1879.	1873.	1874	875.	1876.	1877.	878		.088	1881	885	1880, 1881, 1882, Total
Whole number	0.50	1.06	2.60	311	1.00	8.54	3.27	1.35	1 08	0.53	0.46	1 05	0.50	28	122	1.35	1.02	0.74	5.16	5.27	246	5.30	9.82 9.82	116	2.13	3,61
Nules		06-03	##	22	13.4	85	85	24	6.0	00 04		00.45	10.40	51.00	517	14	0.5	0 30	33	148	136	46	8, 8,	510	88	888
ARENTAGE: American	44	19	16	13	10	80,0	23	10	60		.0	91-	4.0	41-	20	55.80	11.9	10	53	139	105	22	56.55	59	83	11.0
Jumary Jumary Jumary March March May Jume July Aug September September Nov mber		94			00		00 40 1-00 40 04 01 70 04 70 40	- 101 - 1-01-400H	H 1-00 1-11010	***************************************	liiilii ilii	20 1 1 1 1 motern	1 1 1 1 1 1 mm ot on 1 20		60 412 03 W. 12 120 04 . 4	2044 4 HL 14	99- 9509-		· +440000001277777	######################################	4528=5 ₂ 22838	Stranger & X 22	F-240400000000	PERSTONE STATE	F848199141F3	82222822222
Wand II	1171171111	-=- ;***	8889984	0-0:004:::	H-04000	84446 A C S		H4000040	H :0101000H	7	111111111111111111111111111111111111111	24444 4	85	4 : 403	0 - 01 - 4 01 01 4 0		- !!		unnannaction	8228228E	8-25552228	745480-1058	000 :	52-1-8-2-55 5-1-5-5-5-5-5-5-5-5-5-5-5-5-5-5-5-5-	0200HHHH40	25 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5

Dropsy. This was returned as the cause of 6 deaths in 1882; all the decedents were females; 2 of American, and 4 of foreign parentage; total 6, or 0.27 per cent of all.

Erysipelas. The number of deaths from this cause in 1882 was 14; 3 less than in 1881 and the per cent. 0.18 less.

Males, 4; females, 10; total, 14; or 0.62 per cent. of all. American parentage, 2; foreign parentage, 12.

For the twenty-seven years, 1856-1882, the statistics are:-

Males, 98; females, 112; total, 210; or 0 55 per cent. of all.

American parentage, 116; foreign parentage, 94.

Fever, Typhoid. The number of deaths from this cause in 1882 was much larger than ever before recorded in Providence; the following figures show the number of each sex and parentage:—

Males, 66; females, 74; total, 140; or 6 27 per cent. of all. American parentage, 54; foreign parentage, 86.

The same statistics for the twenty-seven years, 1856-1882, are:-.

Whole number of deaths, 1,253; annual average, 46.

Males, 637; females, 616; total, 1,253; or 3 21 per cent. of all.

American parentage, 675; foreign parentage, 578.

There have been a few deaths reported simply as fever; some as "bilious fever," and about 30 from typhus fever. There was also in 1869 one death from "intermittent," and one from "malarial" fever; in 1880 there was one death from malarial fever; these are all included in the foregoing figures for the twenty-six years; we have not, however, included the three deaths from malarial fever which occurred in 1881, and five from malarial and remittent fever in 1882, as they indicate what appears to be a new development of the disease in this city, and it seems desirable to keep a separate account of them. The figures for the twenty-six years are found in the following table, which gives all the details of sex, parentage color, season, age, locality and percentage from known causes, for each of the twenty-seven years, 1856–1882, and for the whole period:—

1. 1882. Total	38 110 1233 .78 6.27 8.21	18 66 637 20 74 616	17 54 675 21 86 578	3 62	400120 + 100	11 12 174
880. 1881	2.57 1.73	10.00	28		01-02-40-00 - 00	2118
819. 18	98	13	14	-		899
1878. 1879. 1	1.35	28	23	63	94 : :0101 4 4 00 00 00 00	6100
1877.	# 21 01	88	81 54	CH	0 :00 : H4F-00000	2012
1876.	1.73	119	12	64	0100 :	10 20 20
1875.	3.73	88	40	01	81111 :: 1 95 SE	1000
1874.	3.06	82.24	30	7	e : eee : ce : L	0010
1873.	8.58	22.53	85 83	9	40000 :4046270	ost
1873.	4.31	5%	20.50	100	9 :000000000000000000000000000000000000	200
1871	23	88	8,8	60	<i><u>PAGIMAGIAT-BQWB</u></i>	3-1
1870.	4.16	25.09	85	69	9 444494055	995
1869.	85.35	81 23	83 53	i		200
898	24 21	27	16	:	0 :01-1:1	-0-0
1867.1	3.49	203	10.53	c,		441
1866.	4.54	96	17	1-	H4H9H99900000000	000
1865.	6.90	32.55	64	*	1812 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	2000
1861.	3.85	19	125		9-9-83-83-83 9-9-83-83	91-1
1863.	4.0	208	33	2	004 :01:01-F-400	2002
1862.	25.00	21	14	•	au : 0 : 1 3 1 1 10 1 - 4	02 21 0
1861.	10 01	14	94	ON	9 :430 :44044	0041
1860.	20.00	100	13	24	004 : 104-04 : 22-04	10 00 4
1859.	87.	18	200	:	**************************************	10
1858.	9.4	15	15	i	- 4-40101 00 00 50	21-1
1857.	8.54	19	16	i		*40
1856.	30	123	15	63	01201-0120 0120-0	1-00
TYPHOID FEVER, 1856, 18	Whole number	SEX: Males	PARENTAGE: American Foreign	Colored	SEASON: Junnary March April April May June June Suptimber Octuber Octuber	First quarter

DEATHS FROM TYPHOID FEVER IN PROVIDENCE, TWENTY-SEVEN YEARS, 1856-1882. --CONTINUED.

Sex; parentage; color; season; age; locality; whole number, and percentage from known causes.

Total	20 mg	* 8.48.8.9.8.4.8.4.9.4.4.	25. 25. 25. 25. 25. 25. 25. 25. 25. 25.
1882.	922		20222
1881.	222		4 5488
880.1	552	H : 000000-0040	00 00 00 01 0 04
1879.1	222	: :cu4=cu4a	- 34 - : 00 to
	4808	: :000005-2	9 mmm*
1877, 1878.	P22		10 m 10 m 10 m
1876.18	101-21		2146453
1875.18	2008	.чаю a 5 8 12 4 44 4 : : :	nasaga
1874.18	303	HHH00 0 4 20 0 0 1 1 1 1 1	4-52220
1873. 18	1228		reares
1872. 18	1.23	- : ac = c = c = c = c = c = c = c = c = c	F-#200E
1871.18	21:3	H : 01000 4 E F DOH : 1	0,012422
1870.18	3×2	won-gar-	10 01 00 01 0 00
869.18	1554		80-0 M288
1808, 18	200	04	1-00-00+
867.18	120	H 04-550 1- 1111	646666-
1866, 18	® 21 8	4004000004	B-0-034
1865. 18	*48	- : ::::::::::::::::::::::::::::::::::	80 K 0 H 0 H
1864.18	222		Suuudir
1863. 18	222		4640000
1862.18	000		H : 00045
1801.18	G-E		*******
1860.18	2000	4- :	B-5844
1859.18	202		41-0-00
1858.18	2000		4 9224
1807.18	401		P-01-4-00-4
856.18	888	H :04 : DE040H : : :	0000000
	1:6:		111111
D FEVER.	irst4 months hird 4 months	1 year under	Ë
TYPHOID	First4 months	AGE: Under 1 and 2 2 1 1 2 2 1 1 2 2 1 1 3 0 1 1 3 0 1 1 3 0 1 1 3 0 1 1 3 0 1 1 3 0 1 1 3 0 1 1	Ward

From the table on the preceding pages, we may see the differences in the different classes of decedents from typhoid fever.

Sex. The difference between the number of males and females who die from this cause is comparatively slight. In the twenty-seven years, 1856–1882 inclusive, the number of males was 637 and the number of females 616; a proportion of 103.4 males to each 100 females; or 50 84 males, and 49.16 females in each 100 decedents.

Parentage. The difference between the two classes of parentage is much greater. There were in the twenty-seven years, 675 decedents of American, and 578 of foreign parentage from typhoid fever; a proportion of 116.8 of American to 100 of foreign parentage; or 53.87 American and 46.13 foreign in each 100 decedents.* There have been seven years out of the twenty-seven, when the number of foreign decedents exceeded that of American, but in 1882 the number of the foreign class was so much greater than that of the American, that the difference between the two classes is much less than it has been previous to this date.

Season. It is well known that this disease prevails to a much greater extent in the last part of the year than at any other time; the following figures show the percentage for each quarter for the period of twenty-seven years:—

	Number.	Per cent.	Numb	er. Per cent.
lst quarter	170	18.56	8d quarter 31	825.38
2d quarter	174	13.89	4th quarter59	147.16

There is very little difference between the first and second quarters; both together are but little larger than the number in the third quarter, while the number in the fourth quarter is far larger than in either of the others. Dividing the year in three sections, the proportions are as follows:—

^{*}In the report for 1831, by mistake, these proportions were given in the reverse order, making the number of foreign decedents to appear greater than that of the American.

	Number.	Per cent.
First four months		17.29
Second four months	278	22.19
Third four months	758	60.49

The decedents from typhoid fever are of all ages from the oldest to the youngest, but by far the larger number are between 15 and 40 years of age; the following table shows the number and percentage in each division of age, for 1882, and for the twenty-seven years, 1856-1882:—

Periods.	Under 1 year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and	under 15.	15 and under 20	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	70 and - under 80.	S0 and under 90.	So and	Total
1882. Number.	er.	1	3	3		16	28	50	22	8	5	3	1			341
Per cent 1856-1882.	····	0.71	2.15	2 15	11	43	20.00	85.71	15.71	5.71	3.57	2.15	0.71			100.00
Number. Per cent.	8	13	1 90	82		93	201 16.04	356 28.41	174 13.89	103	88	51		7	1	1,95

In the year 1882, of the 140 decedents, 100 were between 15 and 40 years of age.

The number of deaths in each year, and the aggregate and average age of the decedents were as follows:

ASTOR, LENOX AND TILDEN FOUNDATIONS

Decedents from typhoid fever; aggregate and average age.

:	ber of	Aggre	gate A	ge.	Ave	erage A	ge.
YEARS.	Number Deaths.	Years.	Mos.	Days.	Years.	Mos.	Days.
1856. 1857. 1858. 1859. 1859. 1860. 1861. 1861. 1862. 1863. 1864. 1865. 1866. 1866. 1866. 1866. 1866. 1866. 1866. 1866. 1866. 1866. 1866. 1866. 1866.	80 80 94 87 20 28 21 47 42 83 46 83 24 85 55 55	788 829 1,259 706 722 540 1,456 1,260 2,962 1,420 654 692 1,214 1,421 1,610	5 7 6 8 9 11 4 2 4 7 4 1	8 18 18 19 7 11 28 97 14 17 25 8 13	26 26 34 83 85 81 25 80 82 27 80 19 28 84 27	8 10 3 4 9 11 8 7 10 8 10 8	11 13 29 28 25 25 15 28 9 11 28 28
511	68 61 60 71 83 43 47 40 53 86	1,010 2,161 1,672 1.558 1.988 922 1,243 1,151 1,206 1.505 1,156 3,598	3 6 11 5 8 11 9 9	19 19 24 19 17 28 27 20 4 13	31 27 25 27 27 28 24 80 25 30	9 5 10 8 11 11 6 2 4 5	19 1 24 19 16 5 2 1 24 2
27 years	1,258	85,825	5	4	28	7	3

Previous to the year 1882, the greatest number of deaths from typhoid fever, in any one year, in Providence, was 82 in 1865, and the annual average number was 42, and the percentage of the whole number of deaths was 3.08 per cent. In the year 1882, there was a large and remarkable increase, the disease reaching an epidemic condition, in October and November. There were 140 deaths from typhoid fever, during the year, and the percentage was 6.27 Though the number of deaths from the disease was so much greater than in any previous year, the percentage of all the deaths, was less than in 1865.

The outbreak of fever, in 1882, was so very remarkable that it seems proper that a slight sketch of it should be given here, though the full account of it will be given hereafter, in a special report.

In our past experience, there has been little dread of typhoid fever in Providence, and it has always been considered, more especially, a disease of the country than of the city. The greatest number of deaths from typhoid fever in Providence, in any single month, in the 26 years previous to 1882, was 25 in November, 1875. There has generally been an increase of the disease in the last three or four months of the year, so that this increase has come to be expected; but this has not always been the case. In 1881, there were only 14 deaths from fever in the last four months of the year, and in 1879 only 12 deaths in the same period.

There were no special indications of the fever in 1882, up to the last of October. There were only 42 deaths in all the year to the first of November; there were 9 in September, and only 8 in October. In fact, at the middle of the month of October, there seemed to be less typhoid fever in the city than was usual at that season. Only 13 cases were then, or afterwards, reported as having commenced from the first to the twentieth of October inclusive.

After the twentieth of October, the number of cases increased rapidly, and the disease speedily became epidemic. During the ten days, including the last five days of October and the first five days of November, 212 cases were reported, the greatest daily number being 45 on the first day of November. From that date, the disease declined slowly in November, much more rapidly in December, and almost entirely disappeared in January, 1883.

It seems, then, that the epidemic reached its height in ten or twelve days after its beginning, and that, after a certain date, (October 20th,) at which there was scarcely any typhoid fever in the city, within 15 days, 248 cases were reported. These cases were in almost every portion of the city, and without any apparent tendency to concentration in any particular locality. This fact seems to me important, with reference to the popular theory that there must be a direct connection of every case of typhoid fever with some preceding case.

Though the disease had almost disappeared in January and February, 1883, another epidemic of less severity and extent began about the 16th of March. There were 61 cases reported in the twelve days from March 16th to March 27th inclusive. After the latter date the disease rapidly declined.

As already stated, more full details of the history, and more pecially of the causes of the disease, will be given hereafter in special report upon the epidemic.

Heart Diseases of. There were 142 deaths from diseases of the heart in 1882, including 10 from "hypertrophy." There were also 12 specified as "fatty degeneration" of the heart; 12 "endocarditis;" 3 "pericarditis;" 3 "angina pectoris;" and 1 "dilatation." The number in 1881 was 121 and the per cent. 5.68. The number of each sex and parentage in 1882 was:—

Males, 59; females, 83; total, 142; or 6.36 per cent. of all.

American parentage, 78; foreign parentage, 64.

In the twenty-seven years, 1856-1882, the number of deaths from diseases of the heart was 1,845, or 4.81 per cent. of all; annual average, 68. The particulars for each year and for the whole period are given in the following table:—

	8	EX.	Parknt	AGE.	Whole
	Males.	Females.	American.	Foreign.	Number. Percentage
1856	. 8	18	16	5	212.0
1857	. 18	18	22	9	81
1 85 8	. 18	23	28	8	86,
1859	. 13	11	17	7	24
860	. 15	16	21	10	818.9
861	. 23	18	29	12	414.0
942	. 26	23	83	16	495.5
863	. 23	19	30	13	42 8.5
864	. 26	82	44	14	584.6
965	. 18	29	83	14	47
966	. 19	21	27	18	408.9
967	. 26	23	85	14	495.1
168	. 25	27	84	18	52 4.7
369	. 88	29	85	27	625.0
370	. 28	81	83	26	594.7
71	. 40	37	46	31	77
378	. 43	49	63	29	92
73	. 46	54	59	41	1005.6
74	. 46	58	62	42	1045.5
75	. 43	45	47	41	884.6
76	. 41	····· 47····	51	87	884.7
77	47	48	53	40	934.8
78	. 29	44	45	88	834.9
779	. 57	54	63	48	1115.0
80	. 55	49	59	45	104
81	. 57	64	65	56	121
82	. 59	83	78	64	142

Of the 1,845 decedents, 1,128 or 61.14 per cent. were of American parentage.

Hemorrhage. The number of deaths caused by hemorrhage 1882 was 10. Four were from hemorrhage of the lungs, 2 fro "hemorrhage of the bowels," and 4 were not specified.

Males, 7; females, 8; total, 10; or 0.45 per cent. of all.

American parentage, 6; foreign parentage, 4.

Hernia was given as the cause of 7 deaths in 1882; this was more than in 1881 and 0.17 per cent. more.

Males, 5; females, 2; total, 7; or 0.81 per cent. of all. American parentage, 4; foreign parentage, 3.

Hooping Cough. This was the cause of 40 deaths in 1882; it same number as in the preceding year; though the rate was slight less. There were 23 decedents under one year old; 9 between or and two years; 7 between two and five; and 1 between five and to The statistics of sex and parentage are:—

Males, 16; females, 24; total, 40; or 1.79 per cent. of all. American parentage, 20; foreign parentage, 20.

For the twenty-seven years, 1856-1882:-

Whole number, 477; annual average, 18.

Males, 206; females, 271; total, 477; or 1.24 per cent. of all.

American parentage, 209; foreign parentage, 268.

Hydrocephalus. The number of deaths reported from this case in 1882 was 28; 4 less than in 1881; all but two of the deceder were less than ten years old.

Males, 22; females, 6; total, 28; or 1.25 per cent. of all. American parentage, 17; foreign parentage, 11.

For the twenty-seven years, 1856-1882, the statistics are:-

Whole number, 822; annual average, 30.

Males, 452; females, 370; total, 822; or 2.14 per cent. of all.

American parentage, 422; foreign parentage, 400.

Insanity. There were 12 deaths for which this was assigned as the cause in 1882; 4 less than in 1881.

Males, 4; females, 8; total, 12; or 0.54 per cent. of all. American parentage, 10; foreign parentage, 2.

For the twenty-seven years, 1856-1882:-

Whole number, 284; annual average, 10.

Males, 181; females, 153; total, 284; or 0.74 per cent. of all.

American parentage, 204; foreign parentage, 80.

Of the 284 decedents, 46.13 per cent. were males and 53.87 per cent. females; 71.83 per cent. of American, and 28.17 per cent of foreign parentage.

A large proportion of these deaths occur at the Butler Hospital for the insane, which is within the city limits, but where, of course, the patients are gathered from various places. All but one of the twelve deaths in 1882 occurred at this hospital.

Intemperance. The number of deaths ascribed to this cause in 1882 was 18, including 3 from delirium tremens; it was 4 more than in 1881.

Males, 8; females, 10; total, 18; or 0.80 per cent. of all.

American parentage, 5; foreign parentage, 18.

For the twenty-six years, 1857-1882, the statistics are:—

Whole number from known causes, 37,363.

From intemperance, 270; annual average, 10.

Males, 187; females, 83; total, 270; or 0.72 per cent. of all.

American parentage, 75; foreign parentage, 195.

Of the 270 decedents, 69.26 per cent. were males; and 72.22 per cent. were of foreign parentage.

Kidneys, Diseases of. The number of deaths from diseases of the kidneys was 52, including 28 of "Bright's Disease" and one of "Addison's Disease;" the number was 4 more than in 1881. The number of each sex and parentage was as follows:—

Males, 27; females, 25; total, 52; or 2.32 per cent. of all. American parentage, 21; foreign parentage, 31.

For the twenty-seven years, 1856-1882:-

Whole number, 626; annual average, 23.

Males, 359; females, 267; total, 626; or 1.63 per cent. of all.

American parentage, 340; foreign parentage, 286.

Liver, Diseases of. There were 24 deaths in 1882 from diseases of the liver; this was one more than in the preceding year.

Males, 14; females, 10; total, 24; or 1.07 per cent. of all. American parentage, 11; foreign parentage, 13.

For the twenty-seven years, 1856-1882, the figures are:-

Whole number, 427; annual average, 16.

Males, 221; females, 206; total, 427; or 1.11 per cent. of all.

American parentage, 204; foreign parentage, 223.

Malformations. The deaths reported from malformation in 1882 numbered 9, which was 6 less than in 1881. In one case the special malformation was not stated; in two cases it was of the heart, and in one each, it was of the head, brain, spine, liver, anus and intestines. The number of each sex and parentage:—

Males, 3; females, 6; total, 9; or 0.40 per cent. of all.

American parentage, 7; foreign parentage, 2.

For the twenty-seven years, 1856-1882:—

Males, 136; females, 84; total, 220; or 0.57 per cent. of all.

American parentage, 137; foreign parentage, 83.

Of the 220 decedents, 62.27 per cent. were of foreign parentage.

Marasmus. This was reported as the cause of 39 deaths in 1882; 5 more than in 1881; 33 of the decedents, or 84.62 per cent., were less than a year old.

Males, 24; females, 15; total, 39; or 1.74 per cent. of all. American parentage, 15; foreign parentage, 24. For the twenty-seven years, 1856-1882, the statistics are:-

Whole number, 726; annual average, 26.

Males, 392; females, 334; total, 726; or 1.89 per cent. of all.

American parentage, 361; foreign parentage, 365.

Measles. There were only 4 deaths from this cause in 1882; in 1881 there were 25. It has always been very irregular in its appearance in Providence. All the decedents in 1882 were females of foreign parentage, and the 4 deaths were 0.18 per cent. of all. The number of deaths from measles in each year since 1856 has been as follows:—

1856 1	186316	187016	1877 2
1857 2	1864 7	1871 1	187850
185840	1865 7	1872 7	1879
1859 0	1866 9	187328	1880 8
1860 0	1867 0	1874 4	188195
1861 6	1868 4	1875 0	1882 4
1862 1	1869 6	1876 0	
			Total, 27 yrs239

The statistics for the twenty-seven years, 1856-1882, are:-

Whole number, 239; annual average, 9.

Males, 111; females, 128; total, 239; or 0.62 per cent. of all.

American parentage, 70; foreign parentage, 169.

Of the 239 decedents, 70.71 per cent. were of foreign parentage.

Meningitis, Cerebro-Spinal. The number of deaths from this cause in 1882, was 14; 3 more than 1881.

Males, 6; females, 8; total, 14; or 0.62 per cent. of all.

American parentage, 8; foreign parentage 6.

The first cases in Providence were reported in 1864; the following table shows the number in each year since then:—

1864 2	1871 12	1878
1865 2	187210	1879 7
1866 1	187339	18804
1867 0	187410	1881ll
1868 1	1875 6	1882
1869 0	1876 4	
1870 0	1877 7	Total, 19 yearsls7

Whole number from known causes in nineteen years, 31,354. From cerebro-spinal meningitis, 137.

Males, 73; females, 64; total, 137; or 0.44 per cent. of all.

American parentage, 64; foreign parentage, 73.

The number and percentage in each division of age, and according to sex and parentage, are given in the following table, for the period of nineteen years, 1864-1882 inclusive:—

AGES.	Under 1 Year.	1 and under 2,	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50	50 and under 60	60 and under 70.	70 and under 80.	Males.	Females.	American.	Foreign.	Potal
No.	п	9	50	30	14	16	16	18	00	01		-	73	75	3	18	1000
PER CENT.	8.03	4.38	14.59	21.90	10.93	11.68	10.95	9.49	5.84	1.46	.73	. 57.	53.28	46.73.	46.79	53.28	000 000

Of the 137 decedents, 81, or 59.12 per cent. were less than 15 years old.

Murder. There were two deaths in 1882 under this head; 1 less than in the preceding year; both the decedents were males; one of American and one of foreign parentage.

· Old Age. The number of deaths attributed to this cause in 1881 was 79; 6 more than in 1880 or 1881; fourteen of the decedents were 90 years of age; the same number as in the two preceding years: one in 1882 was 102 years, 8 months, and 15 days old.

The particulars of sex and parentage are :-

Males, 29; females, 50; total, 79; or 3.54 per cent. of all. American parentage, 40; foreign parentage, 89.

For the twenty-seven years, 1856-1882:—

Whole number, 1,4°9; annual average, 55.

Males, 503; females, 976; total, 1,479; or 3.85 per cent of all.

American parentage, 890; foreign parentage, 589.

Of the 1,479 decedents, 65.99 per cent. were females, and 60 17 er cent. were of American parentage.

The following figures will show how small a proportion of the eccedents of advanced age are reported under this head. In the year 882 the number of decedents over 70 years of age was 286; 151 of merican and 135 of foreign parentage; this was 12.81 per cent. of Il decedents from known causes during the year.

In the twenty-seven years, 1856-1882, the number of decedents ver 70 years old, was 4.314; 2,941 of American, and 1,373 of forign parentage; 11.24 per cent. of all deaths from known causes.

Paralysis. There were 28 deaths reported in 1882, from paralyis, including 3 from Locomotor Ataxia.

Males, 13; females, 15; total, 28; or 1.25 per cent. of all.

American parentage, 15; foreign parentage, 18.

For the twenty-seven years, 1856-1832, the figures are:—

Whole number, 493; annual average, 18.

Males, 252; females, 241; total, 493; or 1.28 per cent. of all.

American parentage, 354; foreign parentage, 139.

The statistics for apoplexy and paralysis have generally been given gether in this connection. The following table gives the number of zaths from each, in each of the twenty-seven years, 1856-1882:—

1856 21 2 23 1870 28 24 1857 18 9 27 1871 33 28 1858 15 3 18 1872 31 20 1859 16 6 22 1873 36 21 1860 17 11 28 1874 36 23 1861 14 11 23 1875 41 35 1862 10 5 24 1876 44 24 1863 20 8 28 1877 50 24 1864 13 16 29 1878 42 24 1865 20 18 38 1879 56 30 1866 21 8 29 1880 48 30 1867 31 18 49 1881 74 27 1868 22 24 46	ars. Apople:	ty. Paralysis. Tota	l. Years.	Apoplexy.	Paralysis.	Total.
1858 15 3 18 1872 31 20 1859 16 6 22 1873 36 21 1860 17 11 28 1874 36 23 1861 14 11 23 1875 41 35 1862 10 5 24 1876 44 24 1863 20 8 28 1877 50 24 1864 13 16 29 1878 42 24 1865 20 18 38 1879 56 30 1866 21 8 29 1880 48 30 1867 31 18 49 1881 74 27 1868 22 24 46 1882 89 28	iG21	223	1870	28	24	54
1859 .16 .6 .22 1873 .36 .21 1860 .17 .11 .28 1874 .36 .23 1861 .14 .11 .23 1875 .41 .35 1862 .19 .5 .24 1876 .44 .24 1863 .20 .8 .28 1877 .50 .24 1864 .13 .16 .29 1878 .42 .24 1865 .20 .18 .38 1879 .56 .30 1866 .21 .8 .29 1880 .48 .30 1867 .31 .18 .49 1881 .74 .27 1868 .22 .24 .46 1882 .89 .28	5718	927	1871	33	28	6 1
1860 17 11 28 1874 36 23 1861 14 11 23 1875 41 35 1862 19 5 24 1876 44 24 1863 20 8 28 1877 50 24 1864 13 16 29 1878 42 24 1865 20 18 38 1879 56 30 1866 21 8 29 1880 48 30 1867 31 18 49 1881 74 27 1868 22 24 46 1882 89 28	3815	18	1872	31	20	51
1861. 14. 11. 23 1875. 41. 35. 1862. 19. 5. 24 1876. 44. 24. 1863. 20. 8. 28 1877. 50. 24. 1864. 13. 16. 29 1878. 42. 24. 1865. 20. 18. 38 1879. 56. 30. 1866. 21. 8. 29 1880. 48. 30. 1867. 31. 18. 49 1881. 74. 27. 1868. 22. 24. 46 1882. 89. 28.	5916	622	1873	36	21	57
1862 19 5 24 1876 44 24 1863 20 8 28 1877 50 24 1864 13 16 29 1878 42 24 1865 20 18 38 1879 56 30 1866 21 8 29 1880 48 30 1867 31 18 49 1881 74 27 1868 22 24 46 1882 89 28	3017	1128	1874	36	23	s
1863 20 8 28 1877 50 .24 1864 .13 .16 29 1878 .42 .24 1865 .20 .18 .38 1879 .56 .30 1866 .21 8 .29 1880 .48 .30 1867 .31 .18 .49 1881 .74 .27 1868 .22 .24 .46 1882 .89 .28	3114.	23	1875		35	۲
1864 .13 .16 .29 1878 .42 .24 1865 .20 .18 .38 1879 .56 .30 1866 .21 .8 .29 1880 .48 .30 1867 .31 .18 .49 1881 .74 .27 1868 .22 .24 .46 1882 .89 .28	219	524	1876	44	24	
1865	3320	828	1877	50	24	74
1866. .21 8. .29 1880. .48 .30 1867. .31 .18 .49 1881. .74 .27 1868. .22 .24 .46 1882. .89 .28	3418	1629	1878	42	24	6
1867	3520	1838	1879		30	M
1868	3621	829	1880		30	79
•	3731	1849	1881	74	27	101
•	3822	2446	1882	89	28	117
			•			
Totals, 27 years	• • • • • • • • • • • • • • • • • • • •			897	498	. 134

The statistics of sex and parentage for the two causes for the twenty-seven years, are:—

Whole number, 1,380; annual average, 51.

Males, 693; females, 687; total, 1,380; or 3.59 per cent. of all.

American parentage, 967; foreign parentage, 413.

The decedents were 70.07 per cent. of American and 29.93 per cent. of foreign parentage.

Peritonitis. There were 17 deaths from this cause in 1882; 3 km than in the year previous.

Males, 9; females, 8; total, 17; or 0.76 per cent. of all. American parentage, 8; foreign parentage, 9.

The figures are as follows for the twenty-seven years, 1856-1862:-

Whole number, 266; annual average, 10.

Males, 117; females, 149; total, 266; or 0.69 per cent. of all.

American parentage, 144; foreign parentage, 122.

Pneumonia. There were 170 deaths from pneumonia; 10 more than in 1881; and 6 from congestion of the lungs, which was 8 less than in the preceding year. They were divided as follows in the different seasons:—

1st quarter 56	3d quarter16
2d quarter61	4th quarter43

Of the 176 decedents, 39, or 22.15 per cent. were less than ten years old; and 63, or 35.79 per cent. were over fifty years old.

Males, 98; females, 78; total, 176; or 7.88 per cent. of all.

American parentage, 66; foreign parentage, 110.


For the twenty-seven years, 1856-1882, the statistics are:-


Whole number, 2,859; annual average, 105.

Males, 1,469; females, 1,390; total, 2,859; or 7. 45 per cent. of all.

American parentage, 1,407; foreign parentage, 1,452.

The following table gives the statistics of the deaths from pneumonia for each of the twenty-seven years, 1856–1882, showing the number and percentage, sex, parentage, season and age:—





PREUMONIA AND CONGESTION OF LUNGS IN PROVIDENCE, TWENTY-SEVEN YEARS, 1856-1882.

causes.
known
from
and percentage
and
age; whole number,
whole
age;
season;
Sex; parentage;
Sex;

Total	2 850 7.45	1,478	1.407	855 855 855 851 157 168 168 890 890
	15.35	8 &	8 3	Ex-20-0420
	12.2	88	28	285935 969556
<u>8</u>	25. 9. 28. 38.	88	85	F-28.22 - 48.45-6
879	25.55	£ 8	5.2	8817
	₹ <u>9</u> .	13	55	72772 0-1-12 X 73
37.	8.00	<u> </u>	420	<u> </u>
1876. 1877.	£ 2.	22	5 3	ENTERS OF CITED
1873.	19.01	88	2 2	<u> </u>
18. 18.		88	35	<u> </u>
	NH	6.0	23	
	===	28	£ 3	557-2520 :4635
1871. 1872.	£ 10	73	33	Ö2 ∃@€2000.
1950. 1870.	12.25	43	27	150 - 2 - 10 - 10 - 10 - 10 - 10 - 10 - 10
	22 5	9.0	44	* - 3 x 0 - + 0 4 0 6 0
- 1.567 - 1.683.	25.73.	23	4 4	<u> </u>
1367	25.	8 \$_	- ನನ	10 - 2 C 4 4 2 2 4 4 6
98	2 3 <u>5</u>	\$\$	45	—————————————————————————————————————
- 1365	ा सु	88	43	ELESSON SEE
1464	ا بي الله الله الله الله الله الله الله الل	222	83	
- 1463. 1464 	2,8	28	4%	
1863.	E 3	8.33	22.28	74104-120
-	12.77	- ₹%	88	
	8.5	28	នន	
	£ 4.	28	19	
57. 1838. 1839.	2 6	a	28	
<u>55</u>	5 E	22	88	Marker & 600 rend
25 - 25 	17.5	48	28 S 	
PRETMONIA AND CONG. OF LUNGS	Whole numb r	SEX : Mules Femules	Parkitage: American Foreign	SEARON: January Feiruary March April Juni July Auguste

PREUNONIA AND CONGESTION OF LUNGS IN PROVIDENCE, TWENTY-SEVEN YEARS, 1856-1882.—CONTINUED.

Sex; parentage; season; age; whole number, and percentage from known causes.

PNEUMONIA AND COMG. OF LUNGS.	1856.	1857.	1838.	1859	1860.	1861.	1862.	1863.	1884.	1865.	1806		88	1809	1870.	1871.	1873.	1873.	1874.	1875.	1876.	1.178	878.	879.	1880.	28	1883	Total
AGE: Under I year	158	25.00	1	11/00		64	1	1	1 52 4		51.00	120	1 64 12	82		81-		52	200	48	40	25	82	183	1	1		1
	24	-	= 00	=	440		10 m	30 mg	1-40	+2 0+	- :0	540	E- 00 -	100	700	5-1-	01 00	201	1-100	Zu.	14	201	201-	515-				
15 20	0110		:	•		: :	: :		10 m	::		9~4			11				10 CD 610	134	- 21	.00	***	. 09 00				
100	000	00 t- 1				i-t-	500	21-0	-40	00 F= 5		10 10 1	21.	00	91915	X ICC		00 15 1	00 22 3	22:	19.	1-00	-20	612				
1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	-010	0 04 00						000	0101			0 07.10	7=7	225		215		21-0	. z :	383	182	100	280	222				
80 90	1	11			:	:	11	10 00	*	1		:	90		:	*		: 1	: **	00 01	11	44	*	4	*-	G -		1:0
First quarter Second quarter	281 x	250	88×	27.00	227	224	21223	85 54 °C	522	2500	25 2 × 3	Ego;	222	E 22	21 21 1-1	2112	48.	28 28	30	872	2231	8312	3821	注照二	27.7	645	25.5	-
Fourth quarter	14		2 2	2 0		2 5		1 5	6	3 6	2 4	2 3	3 5	7 6	1 3	2 6	3 3		3 2	3 50	2 5	77	20 5	8 8	5 6	9 3	43	
Second 4 months.	200		10	120	165	18	19	18	16	=	18	0	8	19	2 24	2 5	32	58	5	4	37	16	38	5 63	8:3	200	25	604

Itheumatism. The number of deaths from rheumatism, in 1882, was 8; 4 less than in 1881.

Males, 8; females, 5; total, 8; or 0.36 per cent. of all. American parentage, 2; foreign parentage, 6.

Scarlatina. There were but 18 deaths from this cause in 1882; this was less than in any year since 1877; indeed there have been but five years in the twenty-eight years, from 1855–1882, when the number was less than in 1882. Twelve of the deaths, or 66.67 per cent., were in the first half of the year. The decedents were one under one year old; 2 between one and two years; 10 between two and five; and 5 between five and ten.

Males, 12; females, 6; total, 18; or 0,81 per cent. of all.

American parentage, 7; forcign parentage, 11.

For the twenty-seven years, 1856-1882, the general statistics are:-

Whole number, 2,065; annual average, 76.

Males, 1,017; females, 1,048; total, 2,065; or 5.38 per cent. of all.

American parentage, 846; foreign parentage, 1,219.

The following table shows the particulars of sex, parentage, season and locality for each of the twenty-eight years, 1855-1882:—

SCARLATINA.	1855.	1855, 1836.	1857.	1858.	1859.	1860.	1881.	1862	1863.	1961	1865.	1866.	1867.	1868.	1869	1870, 1871	1871.	.1879.	1873.	874.	1875. 1	1876.	177.	1878.	1879.	1880.1	881.	883.	Total
Whole number		4.37 14.00	33.35	1.83	5,18	1.17	98.76	1.58	87.3	1.2	9.08	0.39	1.08	4.60	138	2.80	1.69	1.20	132	268	93	1.87	13	2.89	252	17.	19	18	5,30
Males	218	55	51	288	9191	101	113	5 30	13	13.88	53		99 1-	83	EE	15	15 30	81	25	128	53	16	400	88	135	193	18	51.0	1,088
PARENTAGE: American Foreign	19	70	81	85	87	111	114	60 11	16	82	69	24 14	1-00	978	25.88	55.53	25	81	827	191	50	22	10 00	17	101	38	728	-=	25.5
SEASON: January February March March April May June June September September September September September September September September		228245400-20	00 H410 4014H01 :H01			**************************************	* :**********************************	- : : : : : : : : : : : : : : : : : : :	HIGH SON HILLS	H48048888	888888894 : al	-117171111111	-01::-		812184415148	No.04 : 1 : 1 : 10	@F4=01-		880000000000000000000000000000000000000	82222455558	885858848868	400 40 : : : : : : : : : : : : : : : : :		ผลผายเฉรียงเรียง	E8880044008821	528888888 EXE	·0 × 41-× 44	0100 H-0001 :01 :: H00	25 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
LOCALITY: Mard II III V V V V V V V V V V V V V V V	2-2	8752828	E001-40-	0041-888	9-138608	1 11	+01- :00001 : :	0000000000	∞e∞#####	10110084	**************	[["]["]]]	401		208065000	H - HH401-01-	- 01	7 ::: 4 :u40	803000128	######################################	200-02450	4 ::		2-2 : Surati	828110181	11-51-12182	00 400004	B-H : H : 18H	25 25 25 25 25 25 25 25 25 25 25 25 25 2

Scrofula. The number of deaths from this cause in 1882 was 3; 1 less than in 1881, and 0.13 per cent. of all from known causes.

Males, 2; females, 1; total, 3; or 0.13 per cent. of all. American parentage, 0; foreign parentage, 3.

Skin, Disease of. There was but one death reported under this head in 1882; the decedent was a foreign male.

Small-Pox. No death from this disease occurred in Providence in 1882; from the table given below it will be seen that there had been none since 1875.

The statistics for the twenty-seven years, 1856-1882, are:-

Males, 58; females, 21; total, 79; or 0.21 per cent. of all. American parentage, 48; foreign parentage, 31.

The particulars for each year are as follows:-

SMALL-Pox.

	8	EX.	PARE	NTAGR.	Whole	
	Males.	Females.	American.	Foreign.	Number.	Percentag
856	7		5	4	9	
857	•••••					
			5			
			8			
			•••••			
			1			
			•••••			
			8 8			
			8			
			1			
	.		8			
78	12	9	8	6	14	
74		••••	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • • • • • • • • • • • •	• • • • • • • • • •
75	1		• • • • • • • • • • • • • • • • • • • •		1	0.1
78			• • • • • • • • • • • • • • • • • • • •			
77			· · · · · · · · · · · · · · · · · · ·			
8			•••••		• • • • • • • • • • • • • • • • • • • •	
79		• • • • • • • • • • • • • • • • • • • •	· · · · · · · · · · · · · · · · · · ·			
		•••••				
			• • • • • • • • • • • • • • • • • • • •			•••••
52		• • • • • • • • • • • • • • • • • • • •				•••••

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Spine, Diseases of. The number of deaths in 1882 from diseases of the spine was 9; 3 more than in 1881.

Males, 4; females, 5; total, 9; or 0.40 per cent. of all. American parentage, 3; foreign parentage, 6.

Stomach, Diseases of. There were 24 deaths in 1882 from diseases of the stomach; 16 of them were specified as "inflammation."

Males, 10; females, 14; total, 24; or 1.08 per cent. of all.

American parentage, 2; foreign parentage, 22.

Suicide. The number of suicides in 1882 was 12; 2 less than in 1881. They were specified as follows:—Shooting, 3; arsenic and laudanum, 2 each; opium, morphine and Paris green, one each; one person jumped from a window, and one cut his throat.

Males, 8; females, 4; total, 12; or 0.54 per cent. of all.

American parentage, 8; foreign parentage, 4.

The statistics for the twenty-seven years, 1856-1882, are:—

Males, 123; females, 50; total, 173; or 0.45 per cent. of all. American parentage, 97; foreign parentage, 76.

Teething. The number of deaths reported from this cause in 1882 was 16; 7 more than in 1881.

Males, 10; females, 6; total, 16; or 0.72 per cent. of all.

American parentage, 8; foreign parentage, 8.

Tuberculosis was stated as the cause of 12 deaths in 1882; 13 less than in 1881.

Males, 7; females, 5; total, 12; or 0.54 per cent. of all. American parentage, 6; foreign parentage, 6. The various diseases which properly come under this head have never been reckoned in one class in these reports; if they were so classified the percentage of such a class would exceed that of any other. The heading of "tuberculosis" was not used separately until 1867; since then the deaths have been as follows:—

Whole number from known causes in sixteen years, 27,099.

From tuberculosis, 218; annual average, 14.

Males, 98; females, 120; total, 218; or 0.80 per cent. of all.

American parentage, 142; foreign parentage, 76.

Tumors. There were 6 deaths from tumors in 1882; the same number as in the two preceding years. Two were "ovarian" tumors; 1 abdominal; 1 of the spleen, and 1 of the nose; one was not specified.

Males, 3; females, 3; total, 6; or 0.27 per cent. of all. American parentage, 3; foreign parentage, 3.

Unknown. There were 10 deaths in 1882, for which the cause was not stated, it was 0.45 per cent. of all the deaths during the year. The number was 5 less, and the per cent. 0.25 less than in 1881.

Males, 4; females, 6; total, 10; or 0.45 per cent. of all. American parentage, 2; foreign parentage, 8.

The statistics on this subject for the twenty-seven years, 1856-1882, are:—

Whole number, 575; annual average, 21.

Males, 298; females, 277; total, 575; or 1.48 per cent. of all

American parentage, 196; foreign parentage, 379.

Of the 575 decedents, 65.91 per cent. were of foreign parentage.

STILL-BORN.

The following statistics are given in regard to the still-born children in Providence in 1882.

Number. There were 162 still-born children in 1882; this was only one more than in the year previous; this number was 5.81 per cent. of the 2,788 children born during the year, or one in 17.21.

Sex. The number of males was 96; 5 less than in 1881; the number of females, 66, or 6 more than in the preceding year.

Color. White, 158; colored, 4.

Locality. First ward, 21; second, 4; third, 17; fourth, 13; fifth, 14; sixth, 10; seventh, 12; eighth, 14; ninth, 21; tenth, 36.

Season. January, 14; February, 19; March, 9; April, 8; May, 13; June, 15; July, 15; August, 10; September, 12; October, 13; November, 19; December, 15.

The number and percentage in each quarter of the year were :-

	Number.	Per cent.	Number.	Per cent.
lst quarter	42	25.93	8d quarter37	22.84
2d quarter	86	22.22	4th quarter47	29.01

Parentage. The following table shows the parentage of the 162 still born children in 1882; the first letter indicates the birth-place of the father; the second that of the mother:—

A. and A63	R. and R 1	A. and E
I. and I37	I. and A 7	I. and E
E. and E 8	E. and A 2	A. and 81
8. and 8 2	S. and A 1	A. and B. A
G. and G 3	G. and A 2	I. and B. A1
B. A. and B. A 7	B. A. and A 2	E. and B. A
P. and P 1	A. and I 8	G. and Sw
It. and It 2	E. and I 1	_
N. and N 1	8. and I 1	Total
	l	1

A.—American; I.—Irish; E.—English; G.—German; B. A.—British American P.—Portuguese; It.—Italian; S.—Scotch; N.—Norwegian; R.—Russian; 8%—Swedish.

The 162 still-born children were 5.81 per cent. of all the children born in the city during the year. Of the 2,788 children born, 1,424, or 51 08 per cent. had American mothers; 742, or 26.61 per cent. had Irish mothers; and 622, or 22.31 per cent. had mothers of other foreign nations.

Of the 162 still-born children, 77, or 47.53 per cent. had American mothers; 47, or 29.01 per cent. had Irish mothers; and 38, or 23 46 per cent. had mothers of other foreign nations.

The proportion of still-born to the whole number of children borne by mothers of each class, was as follows: of the 1,424 children of American mothers, 77, or 5.41 per cent. were still-born; that is one in 18.49; of the 742 children of Irish mothers, 47, or 6.33 per cent. or one in 15.79 were still-born; of the 622 children of other foreign mothers, 38 were still-born, or 6.11 per cent., or one in 16 37.

The whole number of children borne by American mothers in 1882 was slightly less than in 1881; but the proportion of these to all the children born was greater than in 1881; of still-born children of American mothers, the number and proportion were less in 1882 than in the preceding year.

Of children of Irish mothers in 1882, the number and proportion were less than in 1881; of still-born children of Irish mothers, the number and proportion were also less than in the year previous.

Of children of other foreign mothers the number and proportion were greater in 1882 than in 1881; also the number and proportion of still-born children.

For the twenty-seven years, 1856-1882, the statistics are: Whole number of children born, 54,640. Number of still-born, 3,171, or 5.80 per cent. of all; or one in 17.23. Of these 3,171 still-born children, 2,973 were white, and 198 colored; 1,838, or 57.96 per cent. were males; and 1,333, or 42.04 per cent. females.

Reckoned according to the fathers' birth-places, 1,387, or 43.74 per cent. were American; 1,332, or 42.01 per cent. were Irish; and 445, or 14.03 per cent. were children of other foreign fathers; of 7 the parentage was not stated.

The following statement shows the whole number of children borne by mothers of different nativities, during the twenty-seven years, 1856-1882, inclusive; also the number of still-born children to mothers of each class; and the percentage of the latter to the whole number of children:—

	Whole number	Still-born	Percentage	\ Or 1
	of children.	children.	of still-born.	in
American mothers	25,887	1,384	5.85	18.70
Irish mothers	20,595	1,407	6.83	14.64
English, Scotch or Welsh mo	others 4,187	204	4 93	20.28
German mothers	1,096	83	8.01	33.21
British American mothers	1,921	85	4.42	22.60
Other foreign mothers	1,004	58	5.78.,	17.81
			5.80	

THE NEW YORK PUBLIC LIBRARY.

ASTOR, LENOX AND TILDEN FOUNDATIONS

CENSUS OF PROVIDENCE, 1880.

Statement showing the nativity, parentage, sex, color, and total population of the city of Providence by wards, according to the United States census of 1880:—

	NATI	VE HTE.	FORE	IGN HTE.	Core	RED.	Total	Total	Total
WARDS.	Males.	F'mls.	Males.	F'mls.	M'les.	Fm's.	Males.	F'mls.	Popula tion.
L	4.501	4,039	1,736		228	258	6,465		11,71
I	1,598	1.812	248	806	230	308	2,076		5 00
III	2,997	3 087	1,089	1.290	115	142	4,201	4,519	8.73
V	2.202	2,353		851	70	76	2,921	3,280	6.29
V	2,912	3.073	1,251	1,423	49	3.5	4,212	4,53)	8,75
VI	2,256	2,531	342	536	219	306	2.817	3,373	6.19
VII	3.169	3,579	1,015	1,240	180	199	4,370	5,036	9,46
VIII	4.615	5,112	1,199	1,568	135	192	5,149	6,867	12.5
X	5,508	5,684	1,458	1.728	420	458	7.386	7,870	15,53
X	5,717	5,576	3,666	3,831	7	13	9,390	9.42	18,81
City	35,475	37,764	12,653	15,319	1,659	1.987	49,787	35,070	114.8

WARDS.	American.	Foreign.	Am. father and foreign mother.		Total. Population
I	5,517	7,311	403	482	13,713
II	3,383	1,429	77	113	5 992
III	3,544	4.598	270	308	8,730
IV	3,171	2.661	118	251	6,201 8 743
V	3.397	4.873	202	271	8 743
V1	4,244	1,676	95	175	6,136
VII	4,524	4.289	228	365	9 406
VIII	7.002	5,138	294	382	12,516
IX	7.601	6,664	464	527	15,256
X	3,303	14,420	474	613	18.810
City	45,686	53,059	2,625	3.487	104,857

The percentage of each class, for the whole city, is as follows: American parentage, 43.57; foreign parentage, 50.60; American father and foreign mother, 2.50; foreign father and American mother, 3.33.

CENSUS OF PROVIDENCE, 1883.

Dwelling houses, tenements, families, sex, and total population by wards.

WARDS.	Dwelling. houses.	Unoccupied houses,	Tenements.	Unoccupied Tenements.	Families.	Males.	Females.	Total Pepulation.
I	1,929	23	3,480	104	3.376	7,337	7,861	15,59
H	828	18	1.072	33	1,039	2,389	3,214	0.60
HI	1,109	8	2,182	47	2,135	4.478	4,987	2,88
IV	930	12	1.658	39	1,619	3,460	3,528	6,277
V	1.084	13	2,214	35	2.179	4.702	4 88	9,555
VI	930	5	1,732	26	1,706	2,957	3,009	6,5W
VII	1.312	5	2,393	35 26 21 73	2,372	4.793	5,385	10,128
VIII	1,797	19 59	3,332	73	3,259	6.563	7.696	24,28
1X	2.421	59	3,999	182	3,817	8,112	8,655	16,75
X	2,691	64	4,689	214	4,475	10,950	11.261	2120
Whole city	15,031	227	26,751	774	25,977	55,736	61,019	116,78

•		
		•
	. ,	
	•	

CENSUS OF PROVIDENCE, 1880.

Statement showing the nativity, parentage, sex, color, and total population of the city of Providence by wards, according to the United States census of 1880:—

WARDS.	NATIVE WHITE.		FOREIGN WHITE.		COLORED.		Total	Total	Total	
	Males.	F'mis.	Males.	F'mls.	M'les.	Fm's.	Males.	F'mis.	Popula tion.	
I	4.501		1,786			258	6,465			
<u> </u>	1,598		248		280	808	2,076			
[[] [V	2,947		1,089		115		4,201	4,519		
	2.202 2.912		649 1.251	851 1.428	70 49		2,921 4,212		8.7 K	
V VI			842		219		2.817		6.19	
V 11	3.169		1.015		184	199	4.870			
VIII	4.615		1,199		135		5.149		12.410	
X	5,508		1,458		420	458	7.886			
X	8.717		8.666		7	18	9,390		18,510	
24		- 0,0.0	0,000	0,001						
City	85,475	87.764	12,658	15,819	1,659	1.9%7	49.787	55,070	1(14.85)	

WARDS.	American.	Foreign.	Am. father and foreign mother.	father and	Total. Population
	5.517	7,811	403	482	13,713
I		1.429	77	113	5 002
[[4.598	270	308	8,720
V		2,661	118	251	6,201
· · · · · · · · · · · · · · · · · · ·		4,873	202	271	8 743
·I		1,676	95	175	6.190
11	4,524	4.289	228	865	9 406
'iii		5,188	294	882	12,616
X		6,664	464	527	15,256
£		14,420	474	613	18.810
City	45,686	58,059	2,625	3,487	104,857

The percentage of each class, for the whole city, is as follows: American parentage, 43.57; foreign parentage, 50.60; American father and foreign mother, 2.50; foreign father and American mother, 3.33.

CENSUS OF PROVIDENCE, 1883.

Dwelling houses, tenements, families, sex, and total population by wards.

WARDS.	Dwelling. houses.	Unoccupied houses.	Tenements.	Unoccupied Tenements.	Families.	Males.	Females.	Total Population.
1	1,929	23	8,480	104	8.876	7,837	7,861	15,1%
11	128	18	1,072	83	1,089	2,389	8,214	5.663
HI	1,109		2,182	47	2,135	4.478	4,967	9,160
IV	930	12	1.658	39	1,619	8,460	8.528	6,550
V	1.084	18	2,214	85	2 170	4,702	4 Rt 8	9,545
VI	980	5	1,782	85 26	1.708	2,957	8.609	665
VII	1.812	6	2,898	21	2,872	4.793	5,885	10,178
VIII	1,797	19	8.832	21 78	8,259	6.563	7.026	14,159
IX	2.421	59	8,999	182	8,817	8,112	8,655	16,767
x	2,691	64	4,689	214	4,475	10,950	11.261	22,211
Whole city	15,081	227	26,751	774	25,977	55,736	61,019	116,755



	•	,		
			•	
		•		



The second

TWENTY-NINTH ANNUAL REPO.

UPON THE

BIRTHS, MARRIAGES, AND I

IN THE

CITY OF PROVIDENCE:

FOR THE YEAR

1883.

By EDWIN M. SNOW, M. D.,

CITY REGISTRAR.



PROVIDENCE:

PROVIDENCE PRESS COMPANY, PRINTERS TO THE CI1884.



THE CITY OF PROVIDENCE.

JOINT RESOLUTION OF THE CITY COUNCIL.

[Approved April 24, 1884.]

RESOLVED, That the twenty-ninth annual report of the City Registr for the year ending December 31, 1883, be received, and that six hand copies thereof, and of the communication accompanying the same, printed in pamphlet form for the use of the City Council, under direct of the joint committee on printing.

A true copy: witness,

HENRY V. A. JOSLIN, CITY CLERK.

INTRODUCTION.

CITY REGISTRAR'S OFFICE, PROVIDENCE, April 21, 1884.

To the Honorable the City Council:

The annual report upon the births, marriages, and deaths in the city of Providence, is herewith presented. The present report contains the statistics upon this subject, for the year 1883, and is the twenty-ninth annual report that I have presented.

During the year 1888, there were 2,916 births reported in Providence, or 128 more than in 1882; the number of persons married was 2,854, or 10 more than in the year preceding; and the number of deaths was 2,328, or 86 more.

The population of the city, by the census of 1880, was 104,857, and by the city census of January, 1883, it was 116,755. The same rate of increase would make the population 119,405 in the middle of the year 1883. The proportions of births, marriages, and deaths to population, in 1883, were, therefore, as follows:

Births, 2,916, or one in 40.95; or 24.42 in each 1,000.

Persons married, 2,354, or one in 50.72; or 19.71 in each 1,000.

Deaths, 2,828, or one in 51.29; or 19.50 in each 1,000.

The present report contains the usual tables of previous reports, and various additional tables, illustrating the subjects given. Some of the tables have been condensed, so that the report is shorter than those for the last few years.

EDWIN M. SNOW, M. D.,

City Registrar.

CONTENTS.

Вівтня	PAGE.
BIRTHS	i
Births in connection with season, and sex	2-4
Number and proportion of the sexes, 30 years (Table II)	3-4
Color:-Number and proportion of the sexes, 28 years	Š
Color:—Number and proportion of the sexes, 25 years. Color:—Births, marriages and deaths in 28 years. Parentage of children born in each ward, (Table III.) Number and percentage of American and of foreign children, 22 years. Parentage:—Mixture of different nations, (Table IV.) Number of 1st, 2d, etc., children in 1833. and 1855-1883, (Table V.) Average number of children to each mother, 29 years, (Table VI.). Number of the child, with age of the mother, (Table VII.). Plurality-births, statistics for 1833. Plurality-births, 20 years' summary.	5
Number and nercentage of American and of foreign children Street	
Parentage:—Mixture of different nations, (Table IV.)	. '3
Number of 1st, 2d, etc., children in 1883. and 1855-1882, (Table V.)	10
Average number of children to each mother, 29 years, (Table VI.)	18 18
Plurality births, statistics for 1883	14-15
Plurality-births, 29 years' summary	15-17
MARRIAGES. Marriages at different seasons, 29 years, (Table VIII.). Birthplaces of partics married in 1883, (Table IX.). Bride and groom born in same country, etc. Proportions of American, foreign, and mixed marriages, 29 years Marriages of colored persons.	18-3
Marriages at different seasons, 29 years, (Table VIII.)	18-11
Birthplaces of parties married in 1883, (Table IX.)	. 💥
Proportions of American foreign and mixed marriages 90 years	. 20-2.
Marriages of colored persons	
Age of persons married in 1888, (Table X.)	. :
Average age of persons married, 28 years. (Table)	. 2
Marriages of divorced persons	. 9
Denominational	. 25-3
Nativity and residence	. 25-1
Bride and groom born in same country, etc. Proportions of American, foreign, and mixed marriages, 29 years Marriages of colored persons Age of persons married in 1883, (Table X.). Average age of persons married, 28 years. (Table). Number of times married, (Table XI.). Marriages of divorced persons Denominational Nativity and residence. Marriage and education; certificate signed with a mark	, 24-2
Deaths to population in 1883; proportion in each class	•
(Table XII.)	. 3
Deaths by season; numbers and proportions; recapitulation	. N-3
Sex, condition, color, nativity, parentage, and locality of decedents	
Aggregate and average age by parentage, 1883, (Table XIII.)	. ~ i
Aggregate and average age by parentage, 28 years, (Tables XIV. and	d
XV.). Difference between average age of American and foreign decedents (Table).	. <i>3₁-</i> 9 L
(Table)	ີ 3
Percentages at different ages by parentage, 1883 (Table XVI.)	. !
Percentages under five years of age by parentage, 25 years (Table)	: :
Deaths by age and parentage in each month and quarter, 1883, (Table	ė
(Table). Percentages at different ages by parentage, 1883 (Table XVI.) Percentages under five years of age by parentage, 28 years (Table) Percentages at different ages by parentage, 29 years (Table) Deaths by age and parentage in each month and quarter, 1883, (Table XVII.) Causes of death, sex, age, parentage and percentages, (Table XIX.) Causes of death in connection with season, (Table XX.)	. 454
Causes of death in connection with season. (Table XX.)	51-5
Notes on the Causes of Death in 1883	.as-m 5-3
Consumption; sex and parentage	
Consumption; change in rate of mortality	
Cronn: stutistics for 1883 and 28 years (Table)	. N-6
Diarrhoa and dysentery; recapitulation for 44 years	. 64
Diphtheria; statistics for 1883, and 26 years. (Table)	. 6-7
Comparison of Diphtheria, Scarlatina and Croup	. (j)
Heart, diseases of: decedents by sex and parentage, 28 years, (Table).	. 80-6
Meningitis. Cerebro spinal; deaths at each age. 20 years	8
Programming Say Parentage Season Age	. (U-1) (L-10)
Scarlatina; statistics for 1883 and 29 years, (Table)	. 94
Small-pox; recapitulatory table, 28 years	. 🏋
Unknown causes	آليون آ
Still-born children, 1856–1883.	. H
Cholera Infantum; decedents by sex and parentage, 28 years, (Table). Consumption; sex and parentage. Consumption; change in rate of mortality. Consumption in Providence, 28 years, (Table). Croup; statistics for 1883, and 28 years, (Table). Diarrhoa and dysentery; recapitulation for 44 years. Diphtheria; statistics for 1883. and 26 years, (Table). Comparison of Diphtheria, Scarlatina and Croup. Fever, Typhoid, 1883, and 28 years, (Table). Heart, diseases of; decedents by sex and parentage, 28 years, (Table). Meningitis. Cerebro-spinal; deaths at each age, 20 years. Pneumonia; statistics for 1883 and 28 years, (Table). Pneumonia, Sex, Parentage, Season, Age. Scarlatina; statistics for 1883 and 29 years, (Table). Small-pox; recapitulatory table, 28 years. Unknown causes. Still-born children, 1856-1883.	
Births. Marriages and Deaths: numbers and proportions. 29 years	S.
(Table XXI.)	10
(Table XXI.). Births and dethis by parentage; excess of births, 28 years, (Table XXII.)	le 16
Census of Providence, 1880, sex, nativity, parentage, and color	
XXII.). Census of Providence. 1880, sex. nativity, parentage, and color Census of Providence, 1883, houses, tenements, families, sex and to	tal 🤅

CITY REGISTRAR'S REPORT.

1883.

BIRTHS.

The number of births reported in Providence for the year 1883, was 2,916.

The first table given below shows the number in each ward, and for the whole city, in each month and quarter of the year. Also for the city, the number of children of each sex, and the number and sex of the colored children born in each month and quarter of the year:—

TABLE I. BIRTHS, 1883. Wards, Months, Sex and Color.

MONTHS.					WARDS.										Colored children in- cluded in the preceding.		
MONTHS.	ı.	11.	111.	ıv.	v.	VI.	VII.	vm	ıx	x.	Males.	Females.	Total.	м.	F.	Total.	
January February March	24 37 28	5 7 2	18 20 22	6 16 11		9 5 8	16 21 29	21 20 25	34 24 46	42	98 102 143	107 105 119	200 207 262	8 6	4 4 3	19	
1st Quarter	89	14	60	33	60	22	66	66	104	155	338	331	669	17	11	28	
April May June	32 24 36	9 1 7	20 18 26	16 8 15	19 19 34	10 7 10	14 27 13	18 28 32	32 39 53	58 63 56	128 117 131	100 117 151	228 234 282	5 7 5	5 4 9	10 11 14	
2d Quarter	92	17	64	89	72	27	54	78	124	177	376	368	744	17	18	35	
July August September	34 31 35	6 9 7	. 25 38 26	12 7	26 15 14	6 8	14 22 21	34 27 32	36 38 37	65 67 68	136 121 143	114 144 112	250 265 255	6 4	6 5 5	19	
3d Quarter	100	22	89	23	55	20	57	93	111	200	400	370	770	14	16	30	
October November December	37 50 33	5 11 8	29 23 21	11 11 12	20 17 13	14 11 5	22 21 17	28 34 31	42 42 30	55 52 48	123 128 112	140 124 106	2/3 252 218	6 3 4	9 2	10 5 6	
4th Quarter	100	24	73	34	50	30	60	93	114	155	363	370	733	13	8	21	
Whole Year	381	77	286	129	237	119	237	330	453	687	1,477	1,439	2,916	61	53	114	

The number of births in 1883, was 128 more than in 1882, and is considerably larger than ever before recorded in Providence. The differences in the several wards, as compared with the preceding year, are as follows: in the first ward, 10 more; second and third, each 20 more; in the fifth, 26 more; in the eighth, 34 more; in the tenth, 86 more; a total increase of 196; the remaining wards lost in numbers, in 1883, as follows: fourth ward, 10 less; sixth and seventh, each 19 less; ninth, 20 less; a combined loss of 68.

Ratio. A census of the city was taken in January, 1883, which showed the population to be 116,755. But allowing a rate of increase between January and July, equal to what it had been since the last preceding census, (June, 1880), we have as the population of the city in the middle of the year 1883, 119,405; and on this basis, all the rates in the present report are reckoned. Thus we have as the ratio of births in 1883, one in 40.95 of the population, or 24.42 births to each thousand inhabitants.

Season. In January, February, April and December, 1883, the number of births was less than in 1882, in all the remaining months the number was larger than in 1882. In the first and last quarters of the year 1883, the number was less than in the corresponding seasons of 1882; in the second and third quarters it was greater. In the first half of 1883, the number of births was 1,413, or 25 more than in 1882; in the last half of the year it was 1,503, or 103 more than in 1882.

The largest monthly total in 1883, was 282 in June; the smallest, 200 in January.

The following statement shows the aggregate number of births in each quarter of the year for the twenty-seven years, 1856 to 1882; the number in each quarter of 1883; and the same for the twenty-eight years, 1856 to 1883.

j	Jan.	April.	First	July.	Oct.	Second	Whole
. м	arch.	June.	6 mos.	Sept.	Dec.	6 mos.	Year.
Twenty-seven years	3,279	.13,611	26,890	18,568	.14,182	27,750	54,649
1883	669	. 744	1,413	770	. 733	1,503	2.916
-							
Twenty-eight years1	3,948	.14,855	28,308	14,338	.14,915	29,253	57.554

For the twenty-nine years, 1855 to 1883, the aggregate monthly

totals of births are as follows; the daily average for each month and for the whole period is also given:—

January.	February.	March.	April.	May.	June.
4,619	4,521	5,165	4,609	4,879	5.230
5.14	5.51	5.74	5.30	5.82	6.01
July.	August.	Sept.	Oct.	Nov.	Dec
4,859	4,976	4,918	5,154	5.015	5.211
5.40	5.53	5.65	5.73	5.76	5.80
	4,619 5.14 July. 4,859	4,6194,521 5.145.51 July. August4,8594,978	5.14 5.51 5.74 July. August. Sept4,8594,9764,918		4,6194,5215,1654,6094,879 5.145.515.745.305.82

For a number of years these figures showed the largest aggregate number of births to be in December; the figures for 1883 show the largest aggregate number in June; the smallest in February. The daily averages hold the same relative positions that they have done before; the highest in June and the lowest in January. In the months of March, June, September, October, November and December, the daily average was above the general average for the period; in the remaining six months it was below the general average.

Proportions of the Sexes. There was less difference than usual in 1883, between the number of males and females born; there were 1,477 male, and 1,439 female children; 38 more males than females; a proportion of 102.6 boys to 100 girls; or 50.65 boys, and 49.35 girls in each 100 children born. In the months of January, February, June, August and October, there were more female than male children born; in the remaining seven months the number of males was larger. The number and proportion of each sex in each quarter of the year were as follows:—

Males. Fema	les.	Males. Fema	les.	Males. Females.
JanMarch338331,	or	102.1 to 100,	or	50.52 and 49.48 in each 100 born.
April-June376368,	or	102.2 to 100,	or	50.54 and 49.46 in each 100 born.
July-Sept400370,	or	108.1 to 100,	or	51.95 and 48.05 in each 100 born.
OctDec363 370,	or	98.1 to 100,	or	49.52 and 50.48 in each 100 born.
Whole year.1,4771,439,	or	102.6 to 100,	or	50 65 and 49.85 in each 100 born.

The first and second quarters are nearly alike; while in the third quarter the excess of males is much larger, and in the fourth much less than in the first half of the year. The same statistics for the twenty-eight years, 1856–1883, the whole number of children born during the period being 57,556, are as follows:—

	Males.	Female	s. Males	. Fema	les. I	Males. F	emales.		
JanMarch	7,217	6,731,	or 107.2	to 100,	or 5	1.74 and	48.26 in	each	100 born.
April-June	7,379	6,976,	or 105.8	to 100,	or 5	1.40 and	48.60 in	each	100 born.
July-Sept	7,309	7,029,	or 103.9	to 100,	or 5	0.97 and	49.08 in	each	100 born.
OctDec	7,685	7,230,	or 106.2	to 100,	or 5	1.52 and	48.48 in	each	100 born.
					_				
28 years	. 29,590	27,966,	or 105.8	to 100,	or 5	1.41 and	48.59 in	each	100 born.

In the thirty years, 1854 to 1883 inclusive, there were 60,525 children born in Providence; 31,108 males, and 29,417 females. The following table shows the proportions of the sexes, in periods of five years:—

TABLE II. BIRTHS, 1854 to 1883. Proportions of the Sexes.

	Males.	Female	8.	Males	. F	emal	le s .	Male	в.	Fema	les			
1854-1858	4,171	3,885,	or	107.4	to	100,	or	51.77	and	48.23	in	each	100	born.
1859-1863	4,050	3,752,	or	107.9	to	100,	or	51.91	and	48.09	in	each	100	born.
1864-1868	3,977	3,787,	or	105.0	to	100,	or	51.23	and	48.78	in	each	100	born.
1869-1873	5,056	4,916,	or	102.2	to	100,	or	50.55	and	49.45	in	each	100	born.
1874-1878	6,817	6,428,	or	106.0	to	100,	or	51.47	and	48.53	in	each	100	born.
1879-1883	7,037	6,619,	or	106.3	to	100,	or	51.58	and	49.47	in	each	100	born.
					_									
30 years	31,108	29,417,	01	105.7	to	100,	or	51.40	and	48.60	in	each	100	bors.

These statisties will be found for each year separately, in the report for 1882, page 5.

Color. The number of colored children born in Providence in 1883, was 114, or 18 more than in 1882. If we assume that the colored inhabitants bear the same proportion to the whole population as in the census of 1880, it gives as the colored population in 1883, 4,151. From these figures we find the ratio of births in this class to be one in 36.41, or 27.46 in each thousand of the colored population. The ratio of all births during the year to the total population was one in 40.95, or 24.42 to each thousand persons. The white population of the city was 115,254; the number of white children born, 2,802, the ratio one in 41.13, or 24.31 to each thousand persons.

The number of colored children was as follows in each ward: first ward, 13; second, 14; third, 4; fourth, 5; sixth, 13; seventh, 15; eighth, 7; ninth, 43; there were no colored children born in the fifth or tenth wards. In the different months the numbers were: January.

7, February 12, March 9, April 10, May 11, June 14, July 12, August 9, September 9, October 10, November 5, December 6. The proportion of the sexes in this class of children were as follows: 61 males and 53 females; or 115.1 males to 100 females; or 53.51 males and 46.49 females in each 100 born. In the twenty-eight years, 1856 to 1883, the proportions have been as follows; the statistics are given by periods of five years from 1856 to 1880; in the report for 1882, page 6, they are given for each year separately.

	Total.	Males.	Females.	Males.	Female	s. Males.	Females.		
1856-1860	222	119	108, o	r 115.5	to 100, e	or 53.60 a	nd 46.40 in	each	100 born.
1861-1865	228	122	106, o	r 115.1	to 100,	or 53.51 a	nd 46.49 in	each	100 born.
1866-1870	421	233	188, 0	r 123.9	to 100,	or 55.34 a	nd 44.66 in	each	100 born.
1871-1875	512	267	215. o	r 108.9	to 190.	or 52.15 a	nd 47.85 in	each	100 born.
1876-1890			•		-				
1881									
1882					•				
1863			•		•				
28 vears.	2 225	1.179	1.046, 0	r 112.7	to 100.	or 52.99 a	nd 47.01 in	each	100 born.

The number of births, marriages and deaths among the colored people of Providence during this same period of twenty-eight years, was as follows:—

Twenty-seven years, 1856-1882	2,111 births	1,042 marriages	2,197 deaths.
1888	114 births	54 marriages	120 deaths.
• -			
Twenty-eight years, 1856-1883	2,225 births	1,006 marriages	2,317 deaths.

The proportion of deaths to births in 1883, was 105.2 to 100; for the whole period it was 104.1 deaths to 100 births.

The births, marriages and deaths among the colored population are given below, in periods of five years from 1856 to 1880, and for each of the last three years, 1881, 1882, and 1883:—

	Births.	Marriages.	Deaths
1856-1860		75	
1861-1865	228		291
1866-1870			86
1871-1875		257	
1876-1880		238	
1881		61	107
18/2	96	53	9:
1883	114	54	120
8 years	2,225		

The figures for each year separately, may be found on page 7, of the report for 1882.

PARENTAGE.

The following tables in relation to the parentage of the children born in the city, contain many interesting facts, showing the great and constantly increasing variety of nations represented in our population.

Table III. shows how many children of different classes of parentage were born in each ward; also the number and percentage of each nationality in the whole city. Where the parents were of different foreign nations, the parentage is given according to the birth-place of the father:—

TABLE III. BIRTHS, 1883. Wards and Parentage.

20020000					WA	RDS.					City.	th 100
PARENTAGE.	I.	11.	ш.	ıv.	y.	vi.	VII.	viii	IX.	x.	Whole	In en
American	145	51	101	58	82	63	110	151	246	179	1,186	40.5
Irish	83	7	81	24	58	8	58	54	81	165	619	21.2
English, Scotch and Welsh	20	3	12	8	14		8	42	25	122	260	9.0
German	7	3	1	1	14	1	5	3	15	28	78	2.6
British American	12	1	4		11	1	9	21	8	22	89	1.0
Portuguese			22					1			23	.7
Other Foreign	12		1	16	7	4	10	2	7	6	65	2.0
Total Foreign	143	14	121	49	104	14	90	123	136	343	1,137	38.99
American father and For- eign mother	45	10	26	11	19	9	18	26	44	63	271	9.30
Foreign father and American mother	48	2	38	11	32	13	19	30	27	102	322	11.04
Ward Totals	381	77	286	129	237	99	237	330	453	687	2,916	100.00

This table has for a number of years shown that the foreign population is largely to be found in the Third, Fifth and Tenth wards; these wards always showing a large excess of children of foreign parentage; in some years the children of Irish parentage alone.

exceeding the number of American parentage; but that is not the case in any ward in 1883. Sometimes the First ward has also shown an excess of the foreign element; in 1883 it was not so, but the number of children of American parentage was only two more than of the foreign class. The "other foreign" not specified in the table were as follows: In the First ward, 12 of Russian and Polish parentage; Third, 1 of Swedish parentage; Fourth, 9 Italian, 6 Swedish, 1 Norwegian; Fifth, 3 Swedish, 2 French, 1 Dutch, 1 Italian; Sixth, 2 Swedish, 1 Russian, 1 Italian; Seventh, 10 Italian; Eighth, 1 Russian, 1 Swedish; Ninth, 5 Swedish, 2 French; Tenth, 3 Swedish, 2 Italian, 1 Austrian. In the whole city there were 23 of Italian parentage; 21 of Swedish; 14 Russian; 4 French; 1 Norwegian; 1 Dutch; 1 Austrian.

The following is a statement of the number and percentage of children of American, foreign and mixed parentage, born since 1856. The statistics are given for periods of five years, from 1856 to 1880 inclusive, and for each of the three years, 1881, 1882, and 1883; also for the whole period:—

					Am.	father.	For. fath	er.
	Ame	rican.	For	Foreign.		mother.	Am. mother	
Years.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.	Number. P	er ct
856-1860	8,327	39.95	4,424	53.10	292	3.51	285	8.4
861-1865	2,706	37 . 57	8,809 .	52 89	352	4.89	336	4.6
866 -1870	3,702	41.92	4,118	46.64	447	5.06	563	6.8
871-1875	4,792	40.23	5.353	45.61	693	5.90	970	8.2
876-1880	5,178	89 95	5.597	43.22	929	7 17	1,251	9.0
1881	1.141	40.71	1,151	41.06	224	7.99	287	10.2
1882	1,146	41.10	1.140	40.90	224	8.03	278	9.9
1002	1 198	40.67	1,187	38 99	971	9.30	899	11.0

It will be seen that the percentage of children of American parentage has varied very little throughout the entire period of twenty-eight years; while the percentage of foreign parentage has decreased considerably. In both classes of mixed parentage, the percentage has steadily increased. The number of children of American parentage in 1883, was larger than ever before, though there have been some years when the percentage was larger. The number of children of foreign parentage in 1883, was but 3 less than in the year before, but the percentage was 1.91 less, and was the lowest ever re-

corded in this class. In the two classes of mixed parentage the number and percentage were both the largest ever recorded.

These particulars are given for each year, from 1856 to 1882, on page 9 of the report for 1882.

The percentage of children born in Providence of parents of each nation, is as follows, for each of the twenty-eight years, 1856-1883, inclusive:—

			English.	British.	Other.
	American.	Irish.	8. & W.	German. American. Po	rtuguese. for a
	Per cent.	Per cent.	Per cent.	Per cent. Per cent. P	er cent. Perct
1856	42.09	41.79	4.84	2.27	: £
1857	38 51	44.82	5.92	3.73	1.51
1858	41.30	42.98	4.82	2.26	1.25
1859	37.60	45.58	5 34	3.26	151
1860	40.11	40.96	6.55	2.85	144
1861	37.91	42.61	6.03		156
1862	38.52	41.53	6.02	3.07	1.64
1863	35.73	43.08	6.35	2.98	1.99
1864	38.69	41.86	6.55	2.75	1.36
1865	36.67	42.90	5.93	1.85	1.55
1866	40.20	39.52	6.25		
1867	39.72	37.81	5.66	3.02	25
1868	41.75	36.01	6.48	1.881.07	
1869	45.19	34.14	4.01	1.811.32	L#
1870	42 85	84.68	6. 19	1.96 1.91	
1871	44.85	32.25	4.54	1.68 2.09	1.73
1872	41.12	32.46	5.67	2.772.31	21
1878	40.84	30.50	7.19	1.882.49	
1874	38.42	32.63	8.13	2.842.19	اللاا
l 875 .	37.50	31.53	7.69	2 852.67	981.48
1876	38.27	31.05	7.75	2.59 2 89	.1.22155
1877	41.45	28.08	6.96	1 . 96 2 . 56	.1.:01.39
				2.552.66	
				1.982.54	
				1.673.27	
				2.033.00	
				1.974.16	
				2 673.05	

The British American were included in the "other foreign" until 1868; and the Portuguese until 1874.

Table IV. gives the birth-blace of both parents, thus showing more exactly the parentage of the 2,916 children born in 1883:—

TABLE IV. BIRTHS, 1883. Parentage.

	BIRTH-PLACES OF MOTHERS.									
BIRTH-PLACES OF FATHERS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number according to the father's birth- place.		
United States	1,186	138	66	2	58		7	1,457		
Ireland	152	578	84	1	10		1	771		
England, Scotland and Wales	78	89	197	2	24		1	341		
Germany	19	5	4	63	2		4	97		
British America	46	5	7		77			135		
Portugal and Western Islands	5	1				22		28		
Other Countries	22	1	2	2	4		56	87		
Whole number according to the mother's birth place	1,508	762	310	70	175	22	69	2,916		

There were 1,186 children whose parents were both natives of the United States; 40.67 per cent. of all children born; the number was 40 more and the per cent. 0.43 less than in 1882. There were also 271 children whose fathers were American and mothers foreign; this made a total of 1,457 children of American fathers, or 49.97 per cent. of all; this was 87 more children, and 0.84 per cent. more than in 1882. There were 322 children whose mothers were American and fathers foreign; this gave a total of 1,508 children of American mothers, or 51.71 per cent. of all; 84 more children, and 0.64 per cent. more than in the previous year. The children of both classes of mixed parentage numbered 593, or 20.34 per cent. of all; this was 91 more children, and 2.34 per cent. more than in 1882. The number of children in 1883, whose parents were both foreign was 1,137, or 38.99 per cent. of all; this was 3 less children and 1.91 per cent. less than in 1882. Of these 1,137 children, there were 988, or 86.89 per cent., whose parents were natives of the same foreign country; and 149, or 13.11 per cent., whose parents were of different foreign countries.

NUMBER OF THE CHILD.

Table V. shows how many of the 2,916 children born in 1883 were the first, second, third, etc., children of their mothers; also the same facts in regard to the 56,240 children born during the twenty-eight years preceding, 1855–1882, inclusive:—

TABLE V. BIRTHS, 1883. Number of the Child.

NUMB	EB	OF C	HILD.	1883.	28 years, 1855–1882.	NUMBER	OF.	CH	ILD.	1883.	26 years, 1855-1882
First oh	ua	of the	mother	755	13,749	Twelfth chi	ld of	the	moth'r	10	
Second	••	44	"	567	11,427	Thirteenth	14	••	••	1	
Third	(F	**	"	484	8,913	Fourteenth		••	**	5	•
Fourth	"	44	**	326	6,809	Fisteenth	**	44	**	 .	2
Fifth	66	44	4	272	5,022	Sixteenth	••	"	"		1
Sixth	44	66	66	188	3,598	Seventeenth	•	44	••		'
Seventh	• •	**	**	184	2,554	Eighteenth	**	44	44	 	
E ighth	"	"	"	85	1,648	Nineteenth	**	••	"	 .	
Ninth	44	60	44	72	1,054						
Tenth	**		"	45	676	Whole nu	mbe	r o	f chil-		
Eleventh	**	44	44	27	370	dren	• • • • •	••••	•••••	2,916	

The 755 children who were the first children of their mothers, were 25.89 per cent. of the 2,916 children born in 1883; this was 1.18 per cent. more than the proportion of first children in 1882. Of the 56,240 children born in the twenty-eight years, 1855 to 1882, there were 13,749, or 24.44 per cent., who were the first children of their mothers.

Average number of children to each mother. There were 31 twin births in Providence in 1883; this number taken from the 2,916 children born during the year leaves 2.885 as the number of mothers bearing children in 1883. Table VII. shows that these 2,885 mothers had borne in all 10,048 children; an average of 3.48 children to each mother.

The next table shows the average number of children to each mother, in each of the twenty-nine years, 1855-1883, inclusive.

This table shows the number of children the mothers included in the figures had borne, previous to the end of the year 1883; but it does not show the average number of children the same mothers may have in their life-time:—

TABLE VI. BIRTHS, 1855 to 1883.

Average number of children to each mother.

	No. of	No. of	Aggregate.
Years.	Mothers.	Children.	No. of Children. Average.
1855	1,588	1,600	5.214 3.28 children to each mothe
1856	1,658	1,675	5,562 3.35 children to each mothe
1857	1,664	1,688	5,786 8.44 children to each mothe
1858	1,706	1,724	5,685 8.88 children to each mothe
1859	1,570	1,593	5,618 3.58 children to each mothe
1860	1,627	1,648	5,841 3.59 children to each mothe
1861	1.694	1,725	6,017 8.55 children to each mothe
18 02 .	1,512	1,529	5,822 8.52 children to each mothe
1863	1,296	1,807	4,717 8.64 children to each mothe
1864	1,327	1,844	4,887 8.68 children to each mothe
1865	1,279	1,998	4,625 8.60 children to each mothe
1866	1,614	1,632	5,744 8.56 children to each mothe
1967	1,595	1,624	5,497 8.50 children to each mothe
1 86 8	1,841	1,866	6,338 8.44 children to each mothe
1869	1,797	1,819	5,954 3.30 children to each mothe
1870	1.859	1,889	6,145 3.31 children to each mothe
871	1,987	1,960	5,420 3.31 children to each mothe
872	2,174	2,206	7,846 8.88 children to each mothe
878	2,104	2,128	6,781 8.20 children to each mothe
1874	2,788	2,780	8,960 8.27 children to each mothe
1875	2,684	2,664	8,977 8.41 children to each mothe
1876	2,604	2,681	8,914 3.42 children to each mothe
1877	2,516	2,548	8,705 8.46 children to each mothe
1878	2,597	2,627	9.417 8.63 children to each mothe
1879	2,499	2.522	8,942 8.58 children to each mothe
1880	2,600	2,627	9,386 3.61 children to each mothe
881	2,780	2,808	10,111 8.64 children to each mothe
1982	2,759	2,788	9,974 8.62 children to each mothe
1868	2,885	-	10,048 8.48 children to each mothe
M	89.484	50 156	202,833 3.47 children to each mothe

Table VII. gives the number of the child in connection with the age of the mother; shows how many first, second, third, etc., children were borne in 1883 by mothers at the ages given; the number of mothers of each age, and the aggregate and average number of children they had borne:—

TABLE VII. BIRTHS, 1883. Number of child with age of mother.

			N	UMB	ER ()F C	HILE	۰.						each	983.	borne.	children age.
THE SOTHERS IN YEARS.														er of mothers at	Children they bore in 1883.	Total children they had borne	Average number of chito to mothers at each ag
18t	åg.	3d.	4th.	5tb.	etb.	7tb.	Stb.	9th.	10tb.	11th.	12t lt.	13th.	14th.	Number Age.	Childr	Total	Avera to m
3. 1 3. 2 7. 12 8. 92 8. 92 8. 97 9. 143 1. 66 2. 98 3. *77 4. 77 6. 90 9. 22 9. 15 3. 11 4. 88 9. 22 9. 30 9. 30 1. 11 9. 14 9. 14 9. 15 9. 15 9. 11 9. 15	*7		29 89 166 188 25 22 *82 56 15 15 15 15 16 16 18 18 18 18 18 18 18 18 18 18 18 18 18	21 11 233 199 40 21 17 17 17 9 5 4 4 7 7 8 8 8 1 1		2 3 3 9 10 9 15 18 13 *14 16 8 8 8 8 2 2 2 1 1	1 1	11 10 7	10000001540001					1 2 3 3 3 3 4 7 7 1 1 1 2 1 3 1 3 3 3 4 7 7 1 1 1 4 1 1 5 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 2 2 3 8 8 477 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	11 22 344 577 344 577 344 577 345 345 345 345 345 345 345 345 345 345	1.0.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1.1

^{*} One of these mothers bore twins.

The reference marks in this table show the number of the first child in each case of twin births; thus at the age of 26, three mothers bore twins; in two cases they were the second and third children, and in one, the third and fourth; at the age of 34, two mothers bore twins; in one case they were the third and fourth children; and in one, the fifth and sixth.

There were in 1883, 153 mothers of 40 years of age and over, who bore 157 children during the year; they had borne in all 1,155

[†] Two of these mothers bore twins.

children, an average of 7.55 to each mother. The record of the ages of the mothers has been kept since July 1, 1863, a period of twenty years and six months; in this time there have been 2,205 mothers of 40 years and over, and they have borne 2,231 children during the time.

The whole number of children borne by these mothers was 15,988, an average of 7.25 children to each mother.

During this twenty years and six months, since July 1, 1863, the number of children borne by mothers whose age was stated, was 45,225; the number of children to mothers at each age, was as follows:—

Age of mother. C	hildren.	Age of mother.	Children.	Age of mother.	Children
14	. 5	27	2,715	. 40	1,004
15	21	28	8,146	41	804
16	64	29	2.336	49	872
17	193	80	3.561	48	207
18	466	81	1,586	44	124
19	776	32	2,165	45	129
20	1,236	83	1,790	46	32
21	1,459	84	1,532	47	81
22	2.183	35	2,261	48	14
23	2,399	36	1,401	49	8
24	2,680	87	1,166	50	11
25	8.025	38	1,252		
26	2,771	89	805	All known ages.	45,225

These 45,225 children were divided as follows, to mothers of different divisions of age:—

Numb	ber of mothers.	Per cent
Under twenty years	. 1,525	3.8
Twenty, and under twenty-five	. 9,957	22.0
Twenty-five, and under thirty	. 13,993	80.9
Thirty, and under thirty-five	10,634	23.5
Thirth-five, and under forty		
Forty, and under forty-five	. 2,011	4.4
Forty-five and over	. 220	0.49
Total	45,225	100.00

PLURALITY BIRTHS.

There were 31 cases of plurality births in Providence in 1883; this was two more than in the preceding year.

The general statistics of these cases are as follows:-

Cases. 31; children, 62.

Locality. First ward, 2; Second, 0; Third, 6; Fourth, 0; Fifth, 5; Sixth, 0; Seventh, 1; Eighth, 3; Ninth, 5; Tenth, 9.

Season. January, 0; February, 0; March, 4; April, 3; May. 3; June, 8; July, 2; August, 3; September, 4; October, 2: November, 0; December, 2.

Sex. In 8 cases, both children were males; in 12, both were females; and in 11 cases there was one child of each sex; total?7 male and 35 female children.

Color. All the parents were white.

Number of the children and age of the mother. From the reference marks in Table VII. we learn the number of plurality births to mothers of different ages; also the number of the children in each In nine cases they were the first and second children of mothers, two of whom were twenty years old; two were 24; and one each of 22, 23, 27, 28, and 31 years; in five cases they were the second and third children of mothers, two of whom were 26 years old, and one each of 28, 31, and 37 years; in three cases the third and fourth children of mothers 26, 34 and 37 years old; in two cases the fourth and fifth children of mothers 28 and 29 years old; in one case the fifth and sixth children of a mother 34 years old; in four, the sixth and seventh children of mothers 31, 35, 37, and 40 years old; in three, the seventh and eighth children of mothers 29, 37, and 43 years old; in two, the eighth and ninth children of mothers 35 and 38 years old; in one case the ninth and tenth children of a mother 47 years old; and in one, the tenth and eleventh children of a mother 40 years old.

Parentage. The parentage of the 31 cases was as follows:-

Cases.	Parentage.	Cases.	Parentage.
10	American.	1	English.
4	Irish.	1	Scotch.
2	Br. American.	2	American and Irish.
1	German.	4	Irish and American.
1	Irish and Scotch.	1	English and Scotch.
1	Irish and English.	1	English and B. American.
1	American and English	h. 1	B. American and American.

TWENTY-NINE YEARS' SUMMARY OF PLURALITY BIRTHS.

There have been 693 cases of plurality births in Providence in the twenty-nine years, 1855 to 1883; 684 of twins and nine of triplets.

Sex. Of the nine cases of triplets, in one case, all the children were males; in five cases, all were females; and in three cases, there were two male children and one female; making 9 male and 18 female children in the nine cases.

Of the 684 cases of twins there were 223 cases where both children were males; in 233 cases, both were females, and in 228 there was one male and one female child; 674 male, and 694 female children.

In the 693 cases of plurality births there were 1,395 children, of whom 683 were males and 712 females; or 104.2 females to 100 males; or 51.04 girls and 49.96 boys in each 100 children.

Color. In all the cases of triplets, the parents were white; in the 684 cases of twins, there were 633 where the parents were white, and 51 where they were colored.

Parentage. Of the triplets, in one case the parents were both American; in five cases, both were Irish; in one case, both German; in one, the father was English and the mother Irish; and in one, the father Scotch and the mother Irish. There was but one American mother in the nine cases.

The nativity of both parents is given in the following table for the 584 cases of twin births; the first letter indicating the father's birth-place and the second the mother's.

A., American; I., Irish; E., English; S., Scotch; W., Welsh; G., German; B. A., British American; D., Dutch; P., Portuguese; It., Italian; Sw., Swedish; N., Norwegian; F., French.

Parentage. No	Parentage. No	Parentage. No.	Parentage. No.
A. and A230	I. and A20	B. A. and I3	
I. and I251	E. and A14	8w. and I1	E. and S3
E. and E 81	8. and A3	It. and I1	E. and B. A1
S. and S 8	B. A. and A 7	P. and I1	I. and W1
G. and G 18	G. and A 2	A. and E5	A. and B. A
B. A. and B. A 18	Sw. and A 1	I. and E	I. and B. A?
D. and D 1	A. and I17	S. and E1	F. and B. A 1
N. and N 1	E. and [12	G. and E1	8. und B. A
P. and P 9	8. and 1 8	B. A. and E1	A. and 8w1
It. and It 2	G. and I 1	A. and S1	Total684

In these 684 cases there were 276 American mothers, or 40.35 per cent. of all; the number of Irish mothers was 289, or 42.25 per cent.; the mothers of other foreign birth numbered 119, or 17.40 per cent. In the 693 cases of plurality births there have been 277 American mothers; this was about 1.00 per cent. of the 27,816 American mothers who bore children during these twenty-nine years. There were 416 mothers of foreign birth, which was 1.36 per cent. of the 30,638 foreign mothers bearing children in this period of years.

Proportion of plurality births to whole number. The following figures show the number and proportion of plurality births in 1883, and in the twenty-nine years, 1855 to 1883:—

28 years, 1855-188254,907		
29 years, 1855-188857,761	6849	58,45459,156
Proportion of cases of	plurality births :—	
	Twins.	Briplets.
1855-1882		
1888	1 in 98.1	•••••
1855–1883	1 in 85.5	1 in 6.494.8

Proportion of children of plurality births:-

	Twins.	Triplets.
L855-1882	1 in 43.1	1 in 2,082.9
1883	1 in 47.0	•••••
		
1855-1883	1 in 43.2	1 in 2,190.9

Proportion of plurality births by parentage. In the twenty-nine years, 1855 to 1883, the whole number of births was 58,454. The following table shows the number of births of American, foreign, and mixed parentage, and the number and proportion of plurality births in each of these classes:—

Whole No	. Births. Plura	lity Cases.	1 in	Per Ct.
American parents2	8,510	231	101.7	0.98
Foreign parents 2	7,190	886	70.4	1.42
Am. father, foreign mother.	3,448	30	114.9	0.87
Foreign father, Am. mother.	4,306	46	93.6	1.07
_				
Total	3,454	693	84.3	1.19

Of the 693 cases during this period, the 231 of American parentage, were 33.33 per cent.; the 386 of foreign parentage, were 55.70 per cent.; the 30 with American fathers and foreign mothers were 4.33 per cent.; and the 46 with foreign fathers and American mothers were 6.64 per cent.

MARRIAGES.

The number of marriages in Providence in 1883, was 1,177.

Table VIII. gives the number of marriages in each month and quarter of the year 1883, and of each of the six years preceding; also the aggregate and average number for each month and quarter for the twenty-nine years, 1855 to 1883:—

TABLE VIII. MARRIAGES. Season.

MONTHS.	1883.	1882.	1881.	1880.	1879.	1878.	1877.	29 years. 1855-1883.	average.
January	113	105	117	105	102	76	68	2,241	п
February	65	82	112	100	85	73	65	1,970	0
March	66	55	81	53	47	73	38	1.312	43
1st Quarter	214	242	310	258	234	222	171	5,523	190
April	117	109	95	98	83	77	-83	2,052	71
May	97	91	114	109	98	~68	85	2,195	76
June	127	101	131	111	96	104	83	2,338	60
2d Quarter	341	301	340	318	277	249	251	6,575	227
July	83	70	88	97	59	94	85	1,926	
August	72	80	69	75	88	62	76	1,665	57
September	93	111	81	115	109	93	87	2,309	80
3d Quarter	248	261	238	287	253	249	248	5,890	905
October	143	125	134	108	103	99	99	2,631	91
November	133	149	114	148	128	115	102	2,834	36
December	68	94	66	113	76	82	82	2,017	Ø
4th Quarter	344	368	314	869	307	296	283	7,482	230
Yearly totals	1,177	1,172	1,202	1,232	1,071	1,016	953	25,470	87

The number of marriages in 1883 was but five more than in 1882. On page 2, we estimated the population of the city at 119,405; the 2,354 persons married in 1883, give, therefore, a ratio of one in 50.72 of the population, 19.71 persons married in each thousand. The ratio of persons married to births, was 80.73 to 100; of persons married to deaths 101.1 to 100.

Season. These marriages were more evenly divided than usual between the first and last half of the year, there being only 7 more in the last six months than in the first six. The largest quarterly total was 344 in the fourth quarter; the smallest, 244 in the first quarter. The largest number in any one month was 143 in October, and the smallest, 65 in February.

The following figures give the number and percentage in each quarter of the year 1883, and of the twenty-eight years, 1855-1882:—

	18	83.	1855-1882.			
1	Marriages.	Per cent.	Marriages.	Per cent.		
JanMarch	. 244	20.78	5,279	21.78		
April-June	841	28.97	6,234	25.66		
July-Sept	248	91.07	5,649	23.22		
OctDec	844	29.28	7,138	29.89		
Year	1.177	100.00	24,293	100.00		

Table IX. gives the birth-place of both parties in each of the 1,177 marriages during the year 1883:—

Germany...... British America.....

Portugal and Western Islands ...

Other Countries.....

Whole number of brides.....

	BIRTH-PLACES OF BRIDES.							
BIRTH-PLACES OF GROOMS.	United States.	Ireland.	England, Scotland and Wales.	Germany.	British America.	Portugal and Western Islands.	Other Countries.	Whole number of gre
United States	606	44	42	3	36	3	2	736
Ireland	39	135	7		6			15
England, Scotland and Wales	34	17	64		6			121

26

1

11

725

11

18

23 1,1.,

TABLE IX. MARRIAGES, 1883. Nativity of the parties.

Bride and Groom born in the same country. The number of marriages in 1883, in which the bride and groom were natives of the same country was 873, or 74.17 per cent. of all; this was one more marriage, but 0.23 per cent. less than in 1882. In 606 marriages, or 51.49 per cent. of all, both parties were natives of the United States; this was 10 less marriages, and 1.07 less per cent. than in the preceding year. In 135 marriages, or 11.47 per cent., both parties were born in Ireland; 11 more marriages, 0.89 per cent. more than in 1882.

There were 64 marriages where both were born in England or Scotland; 5 more than in 1882; 12 where both were natives of Germany, or 2 less than in the preceding year; and 27 where both were born in British America; 1 less than in 1882.

Bride and Groom born in different countries. In 304 marriages, or 25.83 per cent. of all, the bride and groom were natives of different countries. The number of grooms born in the United States, who married women of foreign nativity was 128, or 42.11 per cent. of the 304. The number of American women who married foreign

men was 119, or 39.14 per cent. of the 304. There were in all 784 grooms born in the United States; of these, the 128 who married foreign women were 17.44 per cent. The whole number of brides born in this country was 725; the 119 who married foreigners were 16.41 per cent. of this number. There were 187 grooms born in Ireland, of whom 52, or 27.81 per cent. married natives of other countries; and 209 brides born in Ireland, of whom 74, or 35.41 per cent., married natives of some other country. Of the 121 grooms born in England or Scotland, 57, or 47.11 per cent., married women of other countries; of the 27 born in Germany, 15, or 55.55 per cent.; of the 57 born in British America, 30, or 52.63 per cent.

The 35 grooms who were natives of "other countries" than those specified in the table were born as follows: in Sweden, 12; in Italy, 7; in Russia, Norway, the West Indies, and on the Atlantic Ocean, 2 each; and 1 each in France, Spain, Denmark, Australia, Sandwich Islands, Calcutta, Island of Corfu, and Island of Guan.

There were 23 brides, natives of "other countries" who were born as follows: in Sweden, 12; in Italy, 6; in Russia, Norway, Bohemia, Switzerland and West Indies, 1 each.

The following table shows the number and percentage of American, foreign and mixed marriages in the twenty-nine years, from 1855 to 1883; the statistics are given for periods of five years from 1855 to 1879; and for each of the four years 1880, 1881, 1882, 1883. The same statistics for each one of the twenty-eight years, 1855-1882, will be found on page 25 of the Report for 1882:—

	Ameri	can.	Fore	ign.	Mixed.	
Years.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent
1855-1859	1,660	54.25	1,090	85.62	310	10.13
1860-1864	1,697	55.40	989	32.29	377	12.31
1865-1869	2,433	58.23	1,216	29.11	529	12.66
870-1874	2.966	55.84	1.508	28.39	838	15.77
875-1879						
	750					
1881	675	56.16	265	23.04	283	21.80
	616					
	606					

The proportion of American marriages during the last four years averages 55.34 per cent.; the foreign 23.71 per cent.; and the mixed 20.95 per cent.

The changes in the proportion of American marriages have been comparatively slight; the proportion of foreign marriages has decreased considerably, and the proportion of mixed marriages (where one of the parties was of American, and the other of foreign birth) has doubled. The whole number of marriages as given in this table is 25,464, while in Table VIII. the total is 25,470; this is owing to imperfect statistics in the early years of the records; see Report for 1879, page 22.

Marriages of colored persons. The number of marriages of colored persons in 1883, was 53; and one in which the man was colored, (Indian), and the woman white; making 54, or 1 more than in 1882. In 1881 and in 1882 there were 5 marriages, where one of the parties was colored, and one white.

The aggregate and average age of the parties in 1883 were:-

54 males......Aggregate age, 1,616 years......Average age, 29.33 year-53 females.....Aggregate age, 1,401 years......Average age, 26.43 year-

The difference in the number of males and females is owing to the fact of the mixed marriages just alluded to. The same difference of course will be observed in the aggregate numbers given below for the period of 28 years, 1856–1883 inclusive. The whole number of marriages of colored persons during that time, either one or both parties being colored, was 1,096, the aggregate and average age of the parties was as follows:—

1,095 males......Agregate age, 32,431 years......Average age, 29.63 years. 1,086 females.....Agregate age, 28,494 years........Average age, 26.24 years.

Ages of persons married. Table X. gives the number and percentage of brides and grooms in each division of ages:—

		AGES OF THE BRIDES.									ber 8.	Jo	
AGES OF THE GROOMS.	Under 20 years.	20 to 25.	25 to 30.	30 to 35.	35 to 40.	40 to 45.	45 to 50.	50 to 55.	55 to 60.	60 to 65.	65 to 70.	Whole number of Grooms.	Percentage Grooms.
Under 20 years	7	4										11	0.93
20 to 25	88	254	53	7	1							403	34.24
25 to 30	29	184	126	15	6	8	1					364	30.93
80 to 35	4	47	67	28	9	1				.,		156	13.25
35 to 40	4	20	21	31	16	3	3					98	8.33
40 to 45	1	5	10	11	13	9	4	1				.54	4.59
45 to 50		3	3	7	15	11	3	1				43	3.65
50 to 55			9	3	5	5	3	4				29	2.46
55 to 60		1			3	3	1		2			10	0.85
60 to 65					1	1			1			3	0.25
65 to 70					2					2		4	0.34
70 to 75								1				1	0.09
75 to 80											1	1	0.09
Whole number of Brides	133	518	289	102	71	36	15	7	3	2	1	1,177	
Percentages of Brides	11.30	44.01	24.55	8.67	6.03	3.06	1.27	0.60	0.25	0.17	0.09		100.00

TABLE X. MARRIAGES, 1883. Ages of the parties.

In 449 marriages, or 38.15 per cent. of all, the bride and groom were in the same division of age.

There were 414 grooms less than 25 years old, or 35.17 per cent. of all; the number of brides less than 25 was 651, or 55.31 per cent. of all.

The number of grooms less than 20 years old was only 11; while the number of brides in the same division of age, was twelve times that number, or 133.

The aggregate and average age of all persons married during the year, were as follows:—

1,177 malesAggregate age, 34,505 yearsAverage a	ge, 29 32 years.
1,177 females Aggregate age, 30,232 years Average a	ge, 25.68 years

The following figures give these statistics for the period of twenty-eight years, 1856-1883; by periods of five years from 1856 to 1880 inclusive; and for each of the three years, 1881, 1882, 1883. They are given for each year separately, in the report for 1882, page 28.

		Males.		Females.				
Years.	No.	Aggregate age.	Average age.	No.	Aggregate age. Average age			
1 856 -1860	.3,037.	84,831 years.	27.93 years	3,034	72,971 years24.05 years			
1861-1865.	3,152	90,329 years	28.66 years	8,152	77,353 years24.54 years			
1 86 6–1870	.4,428.	125,681 years.	28.38 years	4,42H	109,462 years24.72 year			
1871-1875	.5,432.	155,507 years.	28.63 years	5,482.	135,426 years24.93 year			
1876-1880	.5,221.	152,003 years.	29.11 years	5,221.	131,646 years25.21 year			
1881	.1,203.	35,283 years.	29.35 years	1,202	30,801 years25.62 year			
1882	.1,172.	34,068 years.	29.07 years	1,172	29,743 years25.38 year			
1883	.1,177.	34,505 years.	29.32 years	1,177	30,232 years 25.63 year			

The number of males and females does not agree in this table, because in 1856 the ages of 17 males and 19 females, and in 1857, the ages of 7 males and 8 females, were not given.

Number of times married. Table XI. shows how many of the marriages in 1883 were the first, second or third marriages of the parties:—

TABLE XI. MARRIAGES, 1883. Number of times married.

GROOMS.	BRIDES.	Whole number of			
Number of the Marriage.	First.	Second.	Third.	Grooms.	
First	850	75	1	25	
Second	148	69	8	225	
Third	12	10	1	23	
Fourth	3		1	3	
Whole number of Brides	1,012	154	11	1,177	

In 850 cases, or 72.22 per cent. of all, it was the first marriage of both groom and bride; this was 31 less marriages and 2.95 per cent.

less than the first marriages in 1882. The whole number of grooms married for the first time was 926, or 78.67 per cent. of all; of brides, 1,012, or 85.98 per cent.

In 69 cases, or 5.86 per cent., it was the second marriage of both parties.

Marriages of divorced persons. There were, in 1883, 8 marriages in which both parties had been divorced; 42 in which the man only had been divorced; and 45 where the woman only had been divorced; making 95 marriages where one or both parties had been divorced; this was 8.07 per cent. of all marriages during the year, or about one in 12; in 1882 the proportion was 5.55 per cent., or one in 18.

There were 2,354 persons married in Providence in 1883, of whom 103, or 4.37 per cent. of all, had been divorced. Of these 103 divorced persons, 50 were males, and 53 were females.

Of the 95 marriages, in which one or both the parties had been divorced, 2 were remarriages of the same persons.

The 95 marriaged of divorced persons were solemnized by clergymen of different religious denominations as follows:—

Methodist Episcopal30	Episcopal
Congregational 6	Unitarian E
Baptist19	Universalist10
Free Baptist 8	Judges of Supreme Court 3
Seaman's Mission 2	
Advent 1	Free Religious
Latter Day Saints 1	Denomination not known

Denominational. The 1,177 marriages in 1883 were solemnized as follows by clergymen of different denominations or by civil authority:—

Roman Catholic419	Swedish Christian 1
Baptist189	Hebrew 3
Free Baptist 44	Friend's Ceremony
Methodist186	Lutheran
Episcopalian144	Latter Day Sainte
Congregationalist 80	Regenerated Church of Christ 1
Unitarian 34	New Jerusalem Church 4
Universalist 38	Free Religious Society 1
Presbyterian 5	Gospel Free Church 3
United Presbyterian 19	Denomination not specified 6
Seaman's Mission 10	Judge of Supreme Court 10
Evangelical Advent 11	_
Advent Christian 8	Total1.17
	i

Nativity and residence. In 72 marriages, or 6.12 per cent. of all, the bride and groom were both natives of Providence; in 89 cases, the groom only was born here; and in 186, the bride only; this made 419 persons married, who were natives of Providence; this was 17.80 per cent. of the 2,354 persons married during the year.

There were 111 marriages, or 9.43 per cent. of all, where neither party was a resident of the city; of these 111, there were 79, or 71.17 per cent. where both parties came from outside the State; and in 58 cases they both came from Massachusetts; in 8 others, one of the parties was from Massachusetts.

There has been a constant decrease in the number of marriages of non-residents, since the passage of the new law (taking effect July 1st, 1881), requiring all persons married here to obtain a license at the office of the Registrar, previous to the marriage. This may be seen by the following figures:—

5 -41		Both from outside the	Both from One from
Both parties			
non-resident.	Per cent.	State.	Massachusetts.
1860 206	16.72	182	104 15
1881 195	16.22	180	102
1882 153	13.06	96	74
1883 111	9.43	79	58

The number of certificates issued from the Registrar's office during 1883 was 1,288, or 43 more than in 1882; in 96 cases the parties went out of the city to be married; there were 43 cases where persons came to be married in Providence, bringing a certificate from some other town in Rhode Island, where they resided; these marriages are recorded here, although the "Intention of Marriage" is not. There were 58 certificates or 4 50 per cent. of all, that were either not used, or of which no report has been received; it is quite likely that many of these persons also were married in some other town.

Marriage and education. The 2,354 persons married in 1883 were divided as follows, as to sex and nativity:—

Born in United StatesMales	, 784Females	, 725Total, 1	1,459
Born in Foreign countriesMale	s, 443Females	, 452Total,	895

In these different classes, the number and proportion who signed the marriage certificate with a mark were as follows:—

	Whole Number	Signed with	In each
1888.	Married.	a Mark.	100
Males born in United States	734	39	5.3
Females born in United States	725	51	7.00
Total born in United States.	1,459	90	6.1
Males born in foreign countries.	443	61	18.7
Females born in foreign countrie			
Total born in foreign countri	les 895	164	-
Totals			

The whole number of persons born in the United States, who signed the certificate with a mark was 90; 39 males and 51 females; but 26 of these were colored persons (11 males and 15 females); and 46 were children of foreign parents, (19 males and 27 females); so that only 18 persons (9 males and 9 females), of white American parentage signed with a mark; this was 11 more than in 1882, (5 more males and 6 more females).

In every class, except "males born in foreign countries," the number and proportion who signed with a mark in 1883 were larger

than in the year previous. The number born in the United States was 39 more; the proportion 2.74 more. Of the foreign born, the number was 15 more, and the proportion 0.95 more. The whole number was 54 more and the proportion 2.26 more. The increase was largely among persons of American nativity, and much greater among females than males. The whole number of males who signed with a mark in 1883 was 100; 6 more than in 1882; the whole number of females was 154; 48 more than in the preceding year.

The following figures show these facts for the period of twenty-four years, 1859 to 1883:—

	Whole Number	Signed with	In each
1859-1883.	Married.	a Mark.	340
Males born in United States	18,990	646	
Females born in United States		918	64
Total born in United States	28,237	1,564	5.5
Males born in foreign countries.	7,848	2,363	30.11
Females born in foreign countrie	s 7,591	3,405	
Total born in foreign countri	es15,439	5,768	
Totals	43,676	7,332	с.н

DEATHS.

The number of deaths in Providence, during the year 1883, was 2,328; a larger number than ever before reported in the city. The number was 86 more than in 1882, as follows: males, 50 more; females, 36 more; whites, 61 more; colored, 25 more; natives of the United States, 55 more; natives of foreign countries, 31 more; of American parentage, 82 more; of foreign parentage, 4 more.

During the twenty-nine years, from 1855 to 1883 inclusive, the whole number of deaths in Providence, was 42,285; the average annual death rate was one in 50.89 of the population. In 1883, the death rate was one in 51.29, or 19.50 in each 1,000 of the population, estimated at 119,405.

The rate of mortality, in different classes of the population, in 1883, was as follows:—

	Estimated			Deaths to
	Population,	Deaths,	To population	each 1,000 of
	1883.	1883.	one death in	population.
Whole number	119,405	2,328	51.29	19.50
Males	57,001	1,130		19.82
Females	62,404	1.198		19.90
Whites		2,208		19.16
Colored	•	•		
Born in United States				
Born in foreign countr		•		
American parentage	*			
Foreign parentage	•	•		

The general statistics relating to the 2,328 deaths, in Providence, in 1883, are given in Table XII. on the next page:—

TABLE XII. DEATHS, 1883.

Sex, condition, color, nativity, parentage, locality and season.

1883.	January.	February.	March.	April.	May.	Jane.	July.	August.	September.	October.	November.	December.	Whole Year.
Whole number of deaths	218	146	193	227	204	187	250	198	161	195	165	184	2,394
Sex.— Males Females	105 113	75 71	101 92	106 121	85 119	92 95	121 129	92 106	80 81	95 100	84 81	94 90	1,130
Condition.— Married	67 110 32 8 1	54 67 17 5	56 101 24 12	86 109 25 7	77 87 83 7	61 102 20 4	61 164 18 6 1	65 104 19 9	61 82 15 3	70 100 16 8 1	54 78 27 5	67 86 16 14	7,196 1,196 201 58
Color.— White Colored	207 11	140	181 12	213 14	194	175 12	235 15	190	152 9	183 12	163	175	2,208
Nativity.— United States Ireland England and Scotland Germany British America Portugal & West'n Isl'ds Other countries	160 41 8 3 4 1	103 28 7 2 4 1	140 40 7 1 4 1	154 49 14 4 4	126 53 14 2 7 1	146 30 6 2 1	198 37 19 1 6	136 49 10 1 4 1	107 40 9 3 2	143 33 11 3 3 1	108 42 8 1 6	125 43 9 3 2 1	1,641 428 113 26 47 14
Parentage.— American Irish. English and Scotch. German British American Portuguese. Other foreign.	102 75 20 6 6 27	64 52 14 4 5 3	85 84 15 2 4 1	104 85 23 5 5 1 4	90 76 23 5 5 1	95 59 16 4 6 2 5	115 76 34 10 12	89 69 22 1 7 3	69 66 18 4 2	92 65 20 3 8 2	59 75 17 3 11	81 75 16 5 1	1,00° 807 288 59 77 17 65
Locality, Wards.— I II II IV V VI VII VIII IX X Public Institutions.	24 9 29 14 15 11 15 16 29 42	15 4 19 6 12 3 11 14 24 26 12	20 8 24 14 19 13 18 11 24 32	20 8 18 6 26 16 19 26 41 30	27 9 23 13 9 15 20 21 19 35	26 11 18 16 17 7 12 10 28 30 12	22 7 18 20 18 13 13 21 49 57	19 5 9 7 30 10 16 18 30 38 16	10 4 12 7 17 17 12 7 17 12 17 21 39 15	22 5 17 10 18 5 22 20 23 40 13	17 6 14 5 11 10 12 18 14 44 14	25 22 12 12 12 8 17 21 27 33 5	200 年 100 年 100 年 100 日
Totals	218	146	193	227	204	187	250	198	161	195	165	184	2,22

Season. The subject of mortality in its relations to season is of very great interest and importance. In these reports we have, usually, considered the subject only with reference to the four quarters of the year, though the facts are given each year which would enable us to study the mortality by months.

In the year 1883, the largest quarterly mortality was 618 deaths in the second quarter; the smallest, 544 deaths in the fourth quarter. In the previous year, 1882, the largest mortality was in the fourth quarter, and the smallest in the second. The following figures show the number of deaths in each quarter of each of the last six years; also the aggregate and average number of deaths, in each quarter of the year, in the total for twenty-eight years, 1855 to 1882.

	1883.	1882.	1881.	1880.	1879.	1878.	Aggregate. Av'rge
January-March	557	528	. 527	. 560	. 514	541	9,648 841
April-June	618	506	557	533	439	428	8 944 819
							11,532 419
							9,838 351

The number and percentage of deaths in each quarter of the year 1883, and for the twenty-eight years, 1855 to 1882 inclusive, were as follows—

	1	883.	28 years, 1855–188 2.				
	Deaths.	Per cent.	Deaths.	Per cent			
January-March	557	23.92	9,648	24.18			
April-June	618	26.55	8,944	23.80			
July-September	609	2 6.16	11,582	28.80			
October-December	544	28.87	9,833	24.6			

In the long period of twenty-eight years, it may be taken for granted, that the percentages of mortality in the different quarters of the year, are very nearly the standard rule for the city of Providence. At least, the relations of the several quarters of the year to each other, are probably the same as they would be in a much longer period of time.

In the last annual report for 1882, page 36, a table was given, continued from previous reports, showing the number and the percentage of deaths, in each quarter of each of the twenty-eight years, from 1855 to 1882 inclusive. It does not seem to be necessary to print this table in full, from year to year; but rather to begin a new annual statement, and also give some notes upon the general results of the annual statements up to 1882.

First quarter of the year. In 19 of the twenty-eight years, from 1855 to 1882 inclusive, the percentage of deaths in the first quarter, was less than one quarter of the whole; in nine years it was greater. The largest proportion in the first quarter of the year, was 28.92 per cent. in 1856; the smallest proportion, was 20.54 in 1876.

Second quarter. The percentage of deaths in the second quarter of the year, was less than one-fourth of the whole, in 24 of the 28 years, and greater than one-fourth in only 4 years. The largest proportion, in the second quarter of the year, was 27.34 per cent. in 1858; the smallest proportion was 18.70 in 1857.

Third quarter. In only one of the 28 years, 1855 to 1882 inclusive, was the percentage of deaths, in the third quarter of the year less than one-fourth of the whole. The largest proportion, in the third quarter, was 34.21 per cent. in 1855; the smallest proportion was 24.33 in 1879.

Fourth quarter. In 18 of the 28 years, the percentage of deaths in the fourth quarter of the year, was less than one-fourth of all; in 10 years it was greater than one-fourth. The largest proportion, in the fourth quarter, was 28.63 per cent. in 1879; the smallest proportion was 19.96 per cent. in 1858.

The extremly large percentages in any quarter of any year can usually be explained by the character of the mortality in that quarter. Thus, in 1856, there was an epidemic of scarlatina which caused a large mortality in the first quarter of the year; in 1858, the same disease caused an unusual mortality in the second quarter of the year; in 1855, there was a very large proportional mortality from diarrhoad diseases in the third quarter of the year; and in 1879, scarlatina and diphtheria both caused an unusually large mortality in the fourth quarter of the year; scarlatina alone causing 161 deaths in that quarter.

In the whole 28 years, from 1855 to 1882 inclusive, in each 100 deaths in Providence, there were, very nearly, 24 in the first quarter of the year, 22 in the second quarter, 29 in the third, and 25 in the fourth.

The following shows the number and percentage of deaths in Providence, in each quarter of the year, in the aggregate for the 28 years, from 1855 to 1882 inclusive; and also in the year 1883.—

January-March.	April-June.	July-Sept.	OctDec.
Deaths. Per cent.	Deaths. Per ce	nt. Deaths. Per cent.	Deaths. Per ct.
1855-18829,64824.15	.8,94422.38.	11,53228.86	9,63324.61
1883 55723.92	. 618 26.55.	60926.16	54423.37

Sex and Parentage. During the 29 years, 1855 to 1883 inclusive, 42,285 persons died in Providence, of whom 20,679, or 48.90 per cent. were males, and 21,606, or 51.10 per cent. were females.

For many years, in the early history of the present system of registration in Providence, there was, generally, a considerable excess of females among the decedents of American parentage, and an excess of males among the decedents of foreign parentage. During later years, this difference has gradually disappeared, and there is now an excess of females, generally, among the decedents of both American and foreign parentage. In 1883, the decedents were divided, by sex and parentage as follows:—

American parentage		
Totals	 1.198 females	total, 2,328

The decedents of foreign parentage were 234 more than those of American parentage; and the female decedents were 68 more than the male decedents.

Condition. The 2,328 decedents, in 1883, were divided as follows:—

Married	American	parentage	٠.	350 ;	foreign	parentage	٠	429;	total	8	779
Single	**	11		506;	••	••		684;	**	1	,190
Widows	44	"	••	141;	**	44	••	121;	66		262
Widowers	6.	4.	••	44;	44	**		44;	6.		88
Divorced	**	44	••	6;	**	**	••	8;	44	••	9
Totals	**	44	1	.047;	44	"	1	, 2 81 ;	**		2,328

Of the 1,047 decedents of American parentage, 506, or 48.33 per cent. were "single," that is had never been married; and 541, or 51.67 per cent. were married at the time of death, or had been previously. Of the 1,281 decedents of foreign parentage, 684, or

53.39 per cent. were single, and 597, or 46.61 per cent. were, or had been married. This illustrates a difference in the population of American, and of foreign parentage.

In the whole number of decedents the numbers of the several classes, in 1883, as compared with the numbers in 1882, were so follows: Married, 26 more; single, 39 more; widows, 36 more; widowers, 8 less; divorced, 7 less.

Color. There were 2,208 white, and 120 colored decedents in Providence, in 1883. The ratio of deaths to population was: white decedents, 19.16 in each 1,000; colored decedents, 28.91 in each 1.000.

Of the white decedents, 932 were of American, and 1,276 of foreign parentage; of the 120 colored decedents, 115 were of American and 5 of foreign parentage. The white decedents were 61 more, and the colored decedents 25 more, in 1883 than in 1882.

Nativity and Parentage. The record of each decedent in Providence, since 1855, has shown his or her birth-place, and also the birth-places of the decedents' parents; or in other words the nativity, and the parentage of each decedent. The facts obtained from the records in this particular, have been very valuable in many ways. The figures in table XII. show the differences between the nativity and the parentage of the decedents. Thus, there were 1,641 decedents in 1883, who were born in the United States, or 70.49 per cent. of all; but only 1,047, or 44.97 per cent. of all were of American parentage.

There were 687 decedents of foreign birth, or 29.51 per cent. of all; but, including these, there were 1,281 decedents of foreign parestage, or 55.03 per cent. of all.

The decedents of American birth were 55 more in 1883 than in 1882; those of American parentage were 82 more.

The decedents of foreign birth were 31 more in 1883 than in 1882: those of foreign parentage were 4 more.

There were 14 decedents in 1883, who were natives of other for eign countries than those named in table XII. They were as follows: 5 born in Italy; 4 in Sweden; and 1 each in Switzerland, France, Island of Malta, Denmark, and Finland.

There were 45 decedents in 1883; of "other foreign" parentage.

as follows: Italian, 16; Swedish, 8; French, 6; West Indian, 3; Polish and Swiss, and Austrian, 2 each; and 1 each, Russian, African, Danish, Norwegian, Dutch, and Finnish.

Locality. The following table shows the number of deaths, by parentage, in each ward of the city in 1883:—

Wards.	American.	Foreign.	Wards.	American.	Foreign
1	116	131	VII	89	93
H	66	12	VIII	97	110
III	91	132	1X	173	150
IV	64	66	x	98	84
v	91	113	Fub. Inst	itutions. 69	8
V1	93	30			
Whol	e citv				1.28

The whole number of deaths in 1883, in the first ward, was 30 less than in 1882; in the second ward, 5 more; third ward, 12 more; fourth, 4 more; fifth, 36 more; sixth, 23 more; seventh, 13 less; eighth, 13 more; ninth, 22 more; tenth, 5 less; public institutions, 19 more; whole city, 86 more.

The deaths in *Public Institutions*, in each of the last four years, have been as follows:—

	1883.	1882.	1881.	1880
Rhode Island Hospital	79	76	53	40
Butler Hospital	19	22	18	8
Dexter Asylum	22	15	21	18
Home for Aged Men	2	1	0	
Home for Aged Women	5	5	3	7
Children's Home	0	1	0	(
R. C. Orphan Asylum	1	3	2	8
R. C. Convent	3	1	2	
Reform School	0	2	1	(
Little Sisters' Home for the Poor.	7	6	1	(
Friends' School	0	0	6	(
Shelter for colored children	0	0	3	(
Hotels	8	1	4	(
Police Stations				
Railroad Station	1	0	0	(
Small-Pox Hospital	1	0	0	(
8(. Elizabeth Home	1	0	0	(
Cove Promenade	1	0	0	(
Railroad track	1	0	0	(
Fire Engine Station	1	0	0	(
Ray and Rivers	5	0	0	(
				_
Totals	153	134	115	74

In the aggregate mortality for twenty-eight years, the average age of the decedents of American parentage was 8.70 years more than the average of the decedents of foreign parentage.

The highest average age of the decedents of American parentage was 36.56 years, in May; the highest of those of foreign parentage was 27.75 years, in April. The lowest American average was 25.77, in July; the lowest foreign average was 19.12 in the same month.

The following table shows the difference in the average age of the American and foreign decedents, by parentage, in each month of the year 1883, and also in each month for the aggregate mortality in the twenty-eight years, from 1856 to 1883, inclusive:—

Difference in the Average Age of American and Foreign Decedents.

		1883.		28 7	enra, 1856-	1883.
MONTHS.	Averag Yea		Difference.	Averag		Difference
	American.	Foreign.	Years.	American.	Foreign.	Years.
January	88.30	36.88	7.42	84.78	26.60	8.18
February	38.78	32.72	6.06	84.47	24.22	10.25
March	42.71	29.33	13.38	85.71	25.75	9.96
April	38.61	33.48	5.18	34.52	27.75	6.77
May	41.18	87.29	3.89	86.56	27.61	8.95
June	86.99	28.90	8.09	84.95	25.82	9.13
July	28.80	22.47	6.33	25.77	19 12	6.65
August	32.67	31.02	1.65	27.57	19.85	7.73
September	87.45	31.48	5.97	29.61	20.78	8.83
October	32.02	28.64	3.38	88.24	23.77	9.67
November	39.07	32.53	6.54	38.55	26.02	7.58
December	88.06	25.86	2.20	86.18	27.84	8.84
Whole Year	36.74	81.06	5.68	82.93	24.23	8.70

The greatest monthly difference, in the average age of American, and of foreign decedents, by parentage, was 13.38 years in March, 1883; and 10.25 years for February in the aggregate of twenty-eight years, from 1856 to 1883, inclusive.

The least difference was 1.65 years in August 1883; and 6.65 years for July, in the long period.

The next table (XV), shows the aggregate and the average age of the decedents, by parentage, in periods of five years, from 1856 to 1880 inclusive; also in each of the last three years; also in the total for twenty-eight years, from 1856 to 1883 inclusive:—

Table XV. Deaths, 1856-1883.

Recapitulation. Aggregate and average age by parentage.

	Americ	nn l'arei	itage.	Foreig	n Paren	tage.		American Toreign.	and
YEARS.	Deaths.	Aggregate Age. Years.	Average Age. Years.	Deaths.	Aggregate Age Years.	Average Age. Years.	Deaths.	Aggregato Age. Years.	Average Age. Years.
1856-1860	2,567	78,431	30.55	2,340	43,134	18.43	4,907	121,565	24.77
1861-1865	2,875	91,674	31.88	2,796	57,712	20.64	5,671	149.3%6	26.34
1866-1870	2,906	101,511	34.93	2,719	66 619	24.50	5,625	168,1 3 0	29.89
1871-1875	4,136	136,979	33.11	4,342	104,330	24.02	8,478	241,309	28.46
1876-1880	4,533	145,773	32.15	5,365	132,722	24.78	4 63,6	278,495	28.13
1861	999	34.126	34.16	1,146	32,182	28.08	2,145	66,308	30.91
1882	965	32.541	83.72	1,277	38,750	30.34	2,242	71,291	31.80
1883	1,047	38.468	36.74	1,981	89,792	31.06	2.328	78,260	33.62
28 years totals	20,028	659,503	32.93	21,266	515,241	24.23	41.294	1,174,744	28.45

The preceding table differs from table XV. in the report for the year 1882. That table showed the facts for each separate year, from 1856 to 1882 inclusive. In the above table we have divided the facts into periods of five years each, thus showing more distinctly the changes that have taken place in the average age of the decedents according to parentage.

It will be noticed that the average age of the decedents of American parentage has changed but little during the whole period of twenty-eight years. In 1860, it was almost the same as the average for the whole period, and in the five years, from 1861 to 1865, the average was only about one year less than for the whole period.

In only one of the twenty-eight years was the average age of the decedents of American parentage less than thirty years; but in six of the twenty-eight years the average age of the decedents of foreign parentage was less than twenty years, and in no year, until 1882, was it as high as thirty years. (See page 45 of the Report for 1882).

The preceding table shows that the average age of the decedents of foreign parentage has increased with much regularity and almost constantly since 1856. The average age for the first period of five years, was 18.43 years; and for the last period of five years. (1875-1880), the average was 24.73 years. Since 1880, the average has increased still faster.

The reasons for the increase in the average age of the decedents of foreign parentage, are obvious on reflection. In 1856, the foreign population, by parentage, was composed mostly of middle aged, or younger persons, and children. There were, comparatively, few aged persons. Thus at that time, the average age of the living population of foreign parentage was much less than that of the American population, and of course the average age of the decedents was less.

In 1883, the middle aged persons of foreign parentage have become aged, and the average age of the whole population of foreign parentage is much greater than in 1856. The average age of the decedents is, therefore, greater.

The next table (XVI), shows the number and percentage of the decedents, according to parentage, in different divisions of ages. in 1883, and also in the twenty-eight years, from 1856 to 1883, inclusive:—

Table XVI. Deaths, 1883.

Percentages at different ages, according to parentage.

	Amei Parei 18	itage,	Fore Parer 18	itage,	Total. can an eign,		28 yes 1856-1	
AGES.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.
Under 1 year	209	19.96	209	16.31	418	17.96	7.958	19.27
1 and under 2	56	5.35	74	5.78	130	5.58	3,442	8.83
2 and under 5	44	4.20	74	5.78	118	5.07	8,678	8.91
Total under 5	309	29.51	357	27.87	C66	28.61	15,078	36.51
5 and under 10	33	3.15	48	8.75	81	3.48	2,049	4.96
10 and under 15	21	2.01	21	1.64	42	1.80	886	2.03
15 and under 20	24	2.29	58	4.53	82	3.52	1,383	3.35
20 and under 30	82	7.83	179	13.97	261	11.21	4,219	10.29
30 and under 40	80	7.64	145	11.32	225	9.67	3,796	9.19
40 and under 50	94	8.98	118	9.21	212	9.11	3,183	7.71
50 and under 60	96	9.17	138	10.77	234	10.05	3,071	7.44
60 and under 70	95	9.07	98	7.65	193	8.29	8,003	7.27
70 and under 80	127	12.13	80	6.25	207	8.89	2.857	6.99
80 and under 90	72	6.88	36	2.81	108	4.64	1,488	3.60
90 and over	14	1.34	8	.23	17	.73	301	. 73
Total	1,047	100.00	1.281	100.00	2,328	100.00	41,294	100.00

Under one year of age, in 1883, the percentage of decedents of American parentage was considerably greater than the percentage of foreign parentage. Between 10 and 15 years of age, the percentage of American was also greater. With these two exceptions in every division of age up to 60 years, the percentage of American decedents was less, and sometimes very much less, than the percentage of decedents of foreign parentage.

Above 60 years, the percentage of decedents of American parentage, is much greater than the percentage of decedents of foreign parentage; the aggregate of decedents above 60 years of age, being:

American decedents, 29.42 per cent. of all; foreign decedents, 16.94 per cent. of all.

Of the decedents in 1883, of American parentage, 29.51 per cent. and of those of foreign parentage, 27.87 per cent., were under 5 years of age. The percentage of American parentage, under 5 years, was considerably less than in the previous year, and less than the average for 28 years past. The percentage of foreign parentage was less than ever before in any year.

The following shows the percentage, according to parentage, of the decedents under 5 years of age, in Providence, in periods of five years each, from 1856 to 1880; also in each of the three years since 1880; also for the whole period of twenty-eight years, from 1856 to 1883, inclusive:—

	age,	28 yes 56-188	ars,	Foreign 28 ye	1 l'are ars, 1: 1883.	ntage 856-	Parents	and For age. 28 856-1883.	years
PERIODS.	Total Deaths. American Parentage.	Deaths under 5 years. Am. Parentage.	Percentage un- der 5 years. Am. Parentage.	Total Deaths. Foreign Parentage.	Deaths under 5 years. For. Parentage.	Percentage un- der 5 years. For. Parentage.	Total Deaths. American and Foreign.	Deaths under 5 years. American and For.	Percentage un-
1856-1860	2,567	856		2,340	1,218	52.05	4,907	2,074	42.2
1861-1865	2,875	882	30.68	2,796	1,342	48.00	5,671	2,224	39.25
1866-1870	2,906	804	27.67	2,719	1,128	41.49	5,625	1,982	34.55
1871-1875	4,136	1.297	31.36	4,342	1,788	41.18	8,478	3.085	36.39
1876-1880	4,533	1,547	34.13	5,865	2,145	39.98	9,898	3,692	37.30
1881	909	338	33.83	1,146	389	33.95	2,145	727	33.90
1882	965	816	32.75	1,277	362	28.34	2,242	678	30.94
1883	1,047	309	29.51	1,281	357	27.87	2,328	666	28.61
1854-1883	20,028	6,349	81.70	21,266	8,729	41.05	41,294	15,078	36.51

In previous reports, the above table has been given showing the percentages for each year separately, since 1856. A condensation of the facts, as above, seems to illustrate the subject better than by single years.

It will be seen, from the preceding table, that the percentage of decedents of American parentage, under 5 years of age, has not

changed materially since 1856. The percentage in the first quinquennial period, was 33.35; in the last similar period, it was 34.13.

On the other hand, the percentage of decedents of foreign parentage, under 5 years, has steadily decreased in each period from 1856 to the present time. In the first quinquennial period, the percentage was 52.05; in the last, it was 39.98.

In the last three years, the percentage of decedents of both American and foreign parentage, under 5 years, has regularly decreased, until last year, when, in both classes, the percentage was smaller than will probably be maintained in the future. The cause of the very small percentage, under 5 years, in the last two years, was probably, the large number of deaths from typhoid fever, the decedents from this cause being mostly adults, and thus the percentage of adults being increased, the percentage of children must be decreased.

During forty-four years, from 1840 to 1883 inclusive, there were 54,439 decedents in Providence, of whom 20,813, or 38.23 per cent. were under five years of age. The number and percentage in each division of age, under 5 years, were as follows:—

							_
Whole number of decedents	• • • • • • • • • • • • • • • • • • • •	•••	. 54,43	9			
Under one year of age	10,463,	or	19.22	per	cent.	of al	11.
One and under two years	5,225,	or	9.60	per	cent.	of al	ı.
Two and under five years	5,125,	or	9.41	per	cent.	of al	11.
Total under five years	20,813,	or	86.23	per	cent.	of al	11.

The next table, XVII, shows the number and percentage of the decedents according to parentage, in each division of age, in the aggregate for twenty-eight years, from 1856 to 1883, inclusive:—

TABLE XVII, DEATHS, 1856-1883.

Number and percentage at different ages, according to parentage.

	American 28 years,	Parentage 1856–1883.	Foreign 28 years,	Parentage, 1856–1883.	American Parentage 1856-	, 25 years
AGES.	Number of Deaths.	Percentage in cach division of age.	Number of Deaths.	Percentage in each dlytsion of age.	Number of Deaths.	Percentage in cach division of age.
Under 1 year	3,554	17.75	4,404	20.71	7,958	19.27
1 and under 2	1,286	6.42	2.156	10.14	8,442	8.33
2 and under 5	1,509	7.53	2,169	10.20	3.678	8.91
Total under 5	6,319	31.70	8,729	41.05	15,078	36.51
5 and under 10	901	4.95	1,058	4.97	2,049	4.96
10 and under 15	431	2.15	405	1.90	836	2.03
15 and under 20	661	3.30	722	3.40	1.383	3.35
20 and under 30	1,800	8.99	2,449	11.52	4,249	10.29
80 and under 40	1,731	8.64	2,065	9.71	8,796	9.19
40 and under 50	1,530	7.64	1,653	7.77	3,183	7.71
50 and under 60	1,636	8.17	1,435	6.75	3,071	7.44
60 a nd under 70	1,745	8.71	1,258	5.91	3.003	7.27
70 and under 80	1,864	9.31	993	4.67	2,857	6.93
80 and under 90	1,091	5.45	397	1.87	1,488	3.69
90 and over	199	.99	102	.48	301	.73
Totals	20,028	100.00	21,266	100.00	41,294	100.00

In the preceding table, showing the aggregate of decedents, according to parentage, in twenty-eight years past, marked differences will be noticed in the percentages at different ages, between the decedents of American parentage and those of foreign parentage.

The number and the percentage of the decedents of foreign parentage are greater than of those of American parentage, in every division of age up to fifty years, except between 10 and 15 years. The total under 50 years of age shows:—

American parentage	13,493	decedents,	or 67.37 pe	er cent.
Foreign parentage	17,081	decedents,	or 80.32 pe	er cent

Above 50 years, the totals are as follows:-

The explanation of this difference is probably this: that under 50 years of age, the number and percentage of the living population of foreign parentage in Providence, are greater than of American parentage; while above 50 years of age, the opposite is true.

Deaths by age, parentage, and season. Table XVIII, which follows, shows the number of deaths at each age, according to parentage, in each month and quarter of the year. This shows the comparative mortality of the population by parentage, and by season.

The whole number of decedents, during the year 1883, was 2,328, of whom 1,047 were of American, and 1,281 of foreign parentage. The percentages of each class, in the different quarters of the year, were as follows:—

	1st Quarter.	2d Quarter.	3d Quarter.	4th Quarter.
American parentage.	23.97	27 60	26.08	
Foreign parentage				

The decedents under one year of age of American parentage were 209, or 19.96 per cent. of all of American parentage. The decedents of foreign parentage, of the same age, numbered 209; but they were only 16.31 per cent. of all of foreign parentage.

Of the 209 decedents of American parentage, under one year of age, 93, or 44.50 per cent., died in the third quarter of the year. Of the 209 of foreign parentage, under one year of age, 78, or 37.32 per cent. died in the third quarter.

Causes of death. Table XIX, (pages 47-50), shows the number of deaths in Providence, in 1883, from each cause given, and the number from each cause in each division of ages; also the number from each cause, according to sex and parentage; also the percentage, from each cause, of the whole number from known causes.

Season and Disease. Table XX, pages (51-52), shows the number of deaths, from each of the most prominent causes, in each month and quarter of the year 1883.

Table XVIII. Deaths by age and parentage in each month and quarter, 1883.

					Dr	VISIO	ONAL	PE	RIOD	S OF	LI	E.			_	
Dı	YEAR.	Under I.	1 to 2.	2 to 5.	5 to 10.	10 to 15.	15 to 20.	20 to 30.	30 to 40.	40 to 50.	50 to 60.	60 to 70.	70 to 80.		90 and over.	All Ages.
	January February March	18 10 11	5 7	4 3 4	7 2 1	i	2 1 3	7 7 8	7 6 4	10 5 5	8 2 8	12 5 7	9 13 11	8 5 16	4	161 64 85
	1st Quarter	89	16	11	10	3	6	22	17	20	18	24	33	27	5	25
GE.	April May June	8 8 18	8 6 3	3 8 2	4 2 5	i	9	13 9 12	6 11 9	7 8 13	15 8 7	15 4 7	9 15 12	9 5		10 9
PARENTAGE.	2d Quarter	31	12	13	11	8	10	34	26	28	30	26	36	18	8	28
	July August September	45 31 17	9 4 2	5 4	3	200	 1	1 5 4	8 2 5	8 9 5	999	6 5 10	12 10 7	7 7 4		112 85 65
MERICAN	3d Quarter	93	15	9	4	5	3	10	15	22	23	21	29	18	4	277
AM	October November December	21 9 13	94 85 85		5	2	3 1 1	4 5 7	13 5 4	12 7 5	12 4 7	6 9	6 9 14	2 2 5	. 1 i	96 59 83
	4th Quarter	43	13	11	8	5	5	16	.22	24	23	24	29	9	2	234
	Whole Year	209	56	44	33	21	24	82	80	94	96	95	127	72	14	1,007
	January February March	18 14 17	11 4 5	5 6 5	5 2 4	1 2	8 2 6	11 12 17	11 8 20	11 8 8	16 7 10	6 10 7	5 5 4	7 9 3	···i	116 82 108
	1st Quarter	49	20	16	11	5	16	40	39	27	33	23	14	12	1	306
AGE.	April May June	12 13 13	8	4 7 7	4 6	3	11 9 5	22 16 12	17 10 6	10 13 8	19 13 12	4 16 2	9 10 5	4 3 5		123 114 22
PARENTAGE	2d Quarter	38	13	18	14	5	25	50	33	31	44	99	24	12		129
		41 23 14	16 5	6	7 3 3	2 2	3 1 5	17 14 13	6 15 11	14 9 7	8 12 11	6 11 10	7 4 6	4	1	155 109 92
FOREIGN	3d Quarter	78	26	10	13	6	9	44	32	30	31	27	17	6	1	336
	October November December	16	5 5	6	3	2 2 1	5	17 16 12		9 7 14	8 10 12	9 8	5 8 12	04 01 04	···i	105 106 100
	4th Quarter	1	15		10		1 3	45		30	30	26	25	6		310
	Whole Year	209	74	74	48	21	58	179	145	118	138	98	80	36	S	1,28
7	otal Am. & For	418	130	118	81	49	82	261	225	212	234	193	207	108	17	2,32

E

CAUSES OF DEATH.	solal.	emales.)otal.	Tager 1 Year.	bna 2 Toban	.6 05.	10 10.	10 10.10.10.10.10.10.10.10.10.10.10.10.10.1	\$ 10.20. \$	0 to 30.	0 0 0 0 0 0	.09 01 0	.07 o3 G	.08 01 0	.06 01 0	o and o	1 400/2044	Oreign.	- Age distriction	atto of deaths om each cause mun elolw o
	PE	æ	- 1	ام	أ			\div		÷	-+	-	-+	202	08	06	V	H.	T	;; ;;
(various)Burns and Scalds	21.	60 64	g _a	- 01		810	М.	87 :		81 :	- :	80	80 :	<u> </u>				101	8 m	ञ्च
Drowning	40		F 7	:	:	<u> </u>		8	<u>:</u>	:0	C) (-	:	:						
Polsoning	9 04		90	=	<u> </u>		• :		<u>:</u> : :	• ;	<u>:</u>	<u>' :</u>	•	:	:	<u>: :</u>		:		P 80
Railroad	a	:	90	:	<u> </u>	:	<u>:</u>	:	21	4	_	:	\vdots	:	:	:				80
	01 (_	00	:	÷	:	:		:	<u>:</u>	:	<u>:</u>	-	_	≟	<u>:</u>	_			92 (
denitis	01	:	οı ·	- -	÷	:::	÷	:	<u>:</u>	;	=	:	<u>:</u>	:	<u>:</u>	<u>:</u>	:	•		91.
A næmla	- 1	•	7 1	Ť	<u>:</u>		÷	:	_	<u>:</u>	:0	<u>:</u>	:0		<u>:</u>	_			20 00	10
Anomiation	8	2	2				:		_		:	. •	90	• 6		: *	, 2	_		
Asthenia	7	<u>:</u>	-	F		<u>:</u>	:		_	:	-	-		<u>'</u>	:	<u> </u>		<u>:</u>		_
Asthma	87	:	84	:	:	:	:		:	:	:	<u>:</u>	-	:	:	:			_	87
A telectasis Pulmonum	σ,	_	₩.	4	:	<u>:</u>	<u>:</u>	:		:	<u>:</u>	:	÷	<u>:</u>	<u>:</u>	<u>:</u>		04 0	69 /	.
Sindder, Discuses of	÷ -	:	• 0	:	:	<u>:</u> :	<u>:</u>	:		<u>:</u> :	<u>:</u> :	<u>:</u>	-	N		<u>:</u>		0 -		• 0
Bones, Discuses of	- 4		7 [•	•	•	•	:	<u>:</u>	<u>:</u>	:-	<u>:</u>	<u>:</u>		:	<u>:</u>				-
Diseases of	25		47	916	-	4 -	4-	•	<u>:</u> :-	:-	- 0	: 00	- ~						· 6	• [-
Inflammation of	72		2	15	6	=	600	07	a	_	61	_		_	_		64			
	86		22	18	7	9	69	:	:	_	∞	01		<u></u>	~	_	04			_
(various)	9	8:	8;	:	:	<u>:</u>		:	-	_	∞.	۰.	- (<u>.</u>	<u>:</u>	_			<u>ن</u>
Breast	:	***	* 0	:	:	:	<u>:</u>	:	<u>:</u> :	:	٦,	• 6	÷		<u>:</u>	<u>:</u>	_	·		+ 0
Stomach	₹	9.5	91		<u>:</u>	<u>:</u>	<u>:</u>	:	<u>:</u> :	:		70		G =	~ ~	<u>:</u>	_	- 		01
		-		:				<u>:</u> :		:	•	> -	5-	_		<u>:</u>	_	-	3 -	- 0
Carbuncie.	•	- 0	10	<u>:</u>	:	:		<u>:</u>		<u>:</u> :	:-		<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>		- 0	_	10
		• •	10	:	:	<u>:</u>	<u>:</u> :	<u>. </u>	:-	: 0		:	<u>:</u>	<u>:</u>	: :	<u>:</u>	_	:	• 0	• 0
Puerneral Convul-		•	•	:	:	<u>:</u>	:	:	•	•	•	•	<u>:</u>	<u>:</u>	<u>:</u>	<u>:</u>	_	_		•
	:	01	01	:	:	:	:	:	:	_	=	<u>:</u>	<u>:</u>	<u>:</u>				-		69
Child birth, Puerperal Fever	:	36	9	:	:	<u>:</u>	<u>:</u>	:	:	8	9	:	-	:	:	-	_	_	_ 6	8
Cholera Infantum	1		5									_					•			•

Table, XIX.—Continued. Deatils, 1883. Disease, sex, age, parentage, and percentage from known causes.

		SEX.					DIV	(STO)	MY	PER	IOD	40 8	DIVISIONAL PERIODS OF LIFE.	62				PAR	PARENTAGE.	E.
CAUSES OF DEATH.	Males.	Females.	Total.	Under I.	I and I and er 2.	2 10 2	5 to 10.	.61 03 01	15 to 20.	20 10 20.	30 to 40.	40 to 50.	.00 to 60.	.07 01 00	.08 of 07	.00 of 08	90 and Over.	American.	Foreign.	Total.
Oblic	1,00	1000	1 900	:	1 : :0	11:	::0	1::=	: :8	1 5	:-2	1 : : 2	:-2	1:-5	1::	iir		1 101	-616	1 98
Convulsions	30	18	88	40	19	1619	0100	_										138	38	88
Debility Premature Birth	24	20.00	14	120-1	04		1		i				-	00	۳.	i	1	10 00	00 10	18
Diabetes	78	6, 88	9 80	: 55	: 9					-			29 01			-	:-	40	6. 55	59.6
: :	700	20.20	7.3	G9 00	:6		191	·	:-	H	-	01 :	T	٦:	11	-	11	* 64	50 27	20
Dropsy	- 9	103	16	- 00	Ħ	11	- :	Ħ	ii	- :	:-	:-	i	:00	-01	-	11	40	::	16
Emphyseina	-1-	:00	-61	-	:00		Ħ	::		: :	Ì	:-	: 00	:-	7 :		::	:0	-1-	- 57
Golo	-1-	010	200	-0	21 -	:-	:		1	:-	:-	:	i	:01	:	:	1	err-	- 6	20.00
Epilepsy	41	41	00	:	1	-	-	-	:	-	:	24	÷	:	-	_	1	- 01	¥	000
Fever, Typhoid	70	299	128	00 -	:0	. 7	:00	: 25	- 83	:4	** 2	C0 -0	: 30	- 00	7	-	:	0 Z	œ ;:	198
Mularial	7	01	9	-	-	1	:	:	-	-	:	Ì	:	:	01.	-	:	21.	7	9
Gangrene				i		ì			i			1	:-	H	1:	i			1	-
Heart, Diseases of	3-	1.0	169	-	-	69	9	93	0	a	21-	81-	57	50	15	21	1	2-	30 CH	169
leat			- 5							÷	:-	-		1	1		i	. 00	-n	-0:
ornia. from Lunga	**	ex si	en	-				Ħ	H		-:	20-	::	:"		11	::	e 01	- 00	ons
property County	40	- 50	4	1		:	59	•		÷	:	:								•

Table XIX.—Continued. Deaths, 1883. Disease, sex, age, parenlage, and percentage from known causes.

CAUSES OF DEATH.		-									-	DIVISIONAL LERICORS OF LIVE								U
	Males.	Femules.	Total.	Under I Year.	I and under 2.	3 10 5.	5 to 10.	10 to 15.	15 to 20.	30 10 40.	10 10 20"	.00 to 60.	.07 of 00	70 10 80.	.00 01 08	90 and over.	Атметіскп.	Foreign.	Total	Ratio of de- from each co to whole n ber from kn causes.
Insmitton	101	2,2	1-2	9 :	iii						:	1 : 30	1 . 70	1:00	1:-	111	82		1-8	.38
Intemperance Delirium Tremens	t- 04		12 00	1					1		00 01	9 :	:	::		1	1	CD 02		.56
Intussusception	141		01 25		::			:	:-	: *		:	:	. 05	: 01		:	21 21		1.08
Brights Disease of	54	13	25		:	÷	;	-	:	*-	29	8 10		40			18			1.47
Lencocythamia			;	İ	i							11	1	:	11	Н	-			. 7.
" Inflammation of	20 04		2 53							: :		9 -	:	: :	1			201		.13
Lungs, Discases of	92 10	01 2	100	:	:	:	:					:	:	_		_	90 0		200	315
Marasmus	· 25		200	7	-	:00								. 00		: :	55		48	2.07
Measles	9		œ;	:	04 0	10		:		:	:	:	:	:	:	:	=1		30 4	56.
Meningitis, Cerebro-Spinal	20		940		, e	0	9	. :					:				- 00	0 01	250	00.00
Metritis		1	-	i	:	:		:		:	:	:	:		_		:		1	10.
Mumps	1	******	-	-		*****			:	:	:	:			:	:	-	:::	-	8
:	-0	:	-0	İ			:					,	1			-	-0		-0	100
		-															-		•	10.
Old Age	27	25	8					_				:			_	12	44	40	35	3.62
Oplum Habit	- 0.	:	-6	İ	:	1		-	:	:	:	:	:	ť	:	:	-	:	100	10.
Locomotor Ataxia	4	10	4						-			20 01	2 -	9			7	- 00	4	17
Peritonitis	9		01		_	_		4		-					01		13	6	81	.96
Pleurisy	9		1-	****	:	:	:	:					00		:	_	4		-	.30
Pneumonia	100	8	188	98	13	10	9	:	2	3 21		30				•	8	107	188	8.10
	20 1	-	97	0	00	:	:	:		:		**	1	000	091	:	30 4	200	16	69
Prostate, Disease of	•		-	i	:	÷	÷	+		:	÷	:	_	2		:	0.	24		98.

7

Table XIX.—Continued. Deaths, 1883. Disease, sex, age, parentage, and percentage from known causes.

		SEX.					DIV	1810	DIVISIONAL PERIODS OF LIFE.	PER	TODE	OF	LIF	E.				PARENTAGE.	TAGE.		diam
CAUSES OF DEATH.	Males.	Females	.latoT	Under ! Year.	I and I and I	3 to p.	51010.	10 to 12:	.02 01 61	.08 of 02	30 to 40.	40 to 50.	20 to 60.	'04 04 09	70 to 80.	.00 ot 08 bria 00	.1970	American	.IntoT		Matio of di
Pyernia Scarlatina Scarlatina Scarlatina Septiramia Septiramia Septiramia Septiramia Septiral Stomach, Diseases of Stronosh, Diseases of Stronosh, Congenital Congenital Tabes Mescinerical Tectinus and Tris. Nascentium Throat, Disease of Thorroush Throat, Disease of Thorroush Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of Throat, Disease of		01-04-14 615-0 80 15E 1-0-	2.2.24-4.2.3.2.2.4.4.43.22.4.4.				- 20				.00 .00 .00 .00 .00 .00 .00 .00 .00 .00	na		(A) (B) (B) (B) (B) (B) (B) (B) (B) (B) (B	04 .0				900 to : : : : : : : : : : : : : : : : : :		2001-4-2002279491-8-12-92
Totals	1 130	1,198	2.328	18	130	1	ā	10	00	1000	200	010	188	1901	202	108	150	1.047	1.281	100	9,828

Table XX. Deaths, 1883. Season and Disease.

Whole Year.	95 75													
Роцтір Финтег.	18	19	2 10	4				010	14.0	90 0	0.	56	04 SŽ	. 54
December.	13.5					25 ,				10	'	9	16	
November.	144	- X	- 0	CH	31	211	0.54		- 64		-	Œ		i
October.	O t-	2 00	9-	.04	38	00 4	9-		· ·	0+0	•	12	-=	. 04
Third Quarter.	1500	30	21.	8:	- 01	7	0 40	00.0	2 22	100	0 01	30	00	- 6
September.	0. m .		200	*	13	**	0 04	00.0	0 00	7-	1	16	9 24	
'Jsn2n V	000	2 61	es es	64	- 63	7	: 00		, eo	00 4	•	2	=	
July.	1000	E 00	œ-	22	8	9		NG N	2 01	00 0	9 -	6	-1:	
Second Quarter.	225	17	18	9	100	910			17.0	***	- 00	40	: 8	120
Jane.	20:	==	0.4	10	98	40	* :	***	· X	•		00	=	10
May.	917	0 0	00 4	-	10	90 0			9		101	2	14	
.ttiqA	00 E-	- 00	Z 4	*****	40	10.	- :				-	650	: 53	÷
First Quarter,	919	22	13	-	2	54.5		010	120	620	000	35	-=	00 40
March.	10 10 1	410	9-	***	33	0.0	9-		• •	:	*	11	15	54
Гергавту.	400	20.00	10 01	-	: 53	90	9-	-	: 00	90 -		9	13:	:01
January.	1 - au c	C AC	10 01	1:	30	91	- 61		. 9	:	•	15	130	01
CAUSES OF DEATHS.	poplexy	onchitts	incer (all kinds)	nolera Infantum	msumption	nvulsions	bility	" Premature Birth	phtheria	/sentery		ver, Typhold	" Malarial	Adrocephalus.

TABLE XX.—CONTINUED. DEATHS, 1883. Season and Disease.

Whole Year.	88 88 88 88 88 88 88 88 88 88 88 88 88	2,328
Fourth Quarter,	**************************************	544
December.		184
November.	94-59-96-96-96-96-96-96-96-96-96-96-96-96-96	165
October.		105
Third Quarter.	P	600
September		191
August.	201010101010 3010101010101010101010101010	100
July.	245HQ : 02HQ : : : : : : : : : : : : : : : : : : :	250
Second Quarter.	00000 42xx5404	618
Jane.		187
May.	4015-000 : x00081-0101 : 0 :00-00	504
April.		262
First Quarter.	다매교대전에속점점대중대의대-미요4 :	857
Матећ	01-10 C-1-20	183
February.	- :4 :0-00500-GM : : - : : : : : : : : : : : : : : : :	146
January.	01-0000 GG GB - GB - GB - GB	818
CAUSES OF DEATH.	Insanity Intemperatince Kitherys, Diseases of Liver Marismus Maningtis, Cerebro-Spinal Meningtis, Cerebro-Spinal Oil Age Pertronitis, Pretronitis Pretronitis Pretronitis Recumatism Senali Iox Senali Iox Senali Iox Senali Iox Senali Iox Senali Iox Unberculosis Unknown	Totals

NOTES OF THE CAUSES OF DEATH,

DURING THE YEAR 1883.

The following pages are intended as explanations and comments upon the figures in Table XIX and XX; and also contain additional and recapitulatory statistics, in relation to different causes of death, which could not be given in those tables. The percentages given are always reckoned upon the number of deaths from known causes.

The whole number of deaths in Providence, during the year 1883, was 2,328. Of this number, the cause of death was stated in 2,322 cases, and upon this number the percentages are given. The cause was not given in 6 cases, or 0.26 per cent. of the whole number. The number from unknown causes was 4 less than in 1882, and the percentage 0.19 less.

During 28 years, from 1856 to 1883 inclusive, the whole number of deaths in Providence, was 41,294; the number from unknown causes was 581, or 1.41 per cent.; from known causes the number was 40,713. During 44 years, 1840 to 1883 inclusive, there were 54,439 deaths, and of 52,856 the cause was known, and of 1,583 or 2.91 per cent. the cause was unknown.

Table XX. shows the number of deaths in Providence, in each month and quarter of the year 1883, from the most important causes of death. In regard to certain diseases, as pneumonia, cholera infantum, etc., the effect of season is very marked.

Abscesses. There were only 3 deaths reported from abscesses, in 1883, or 0.13 per cent. of all from known causes. The location was given as the leg, in one case; in the other cases the location was not stated.

The statistics of this cause of death are as follows:-

For the year 1883:-

Total deaths from known causes, 2,322.

Deaths from Abscesses, 3; or 0.13 per cent.

Males, 2; female, 1.

American parentage, 2; foreign parentage, 1.

Statistics for 28 years, 1856-1883 inclusive:-

Total deaths from known causes, 40,713.

Deaths from Abscesses, 111; or 0.27 per cent.

Males, 53; females, 58.

American parentage, 51; foreign parentage, 60.

Accidents. There were 66 deaths reported in Providence in 1883. from accidental causes, or 2.84 per cent. of all from known causes. The number was 11 less than in 1882.

There were 9 accidental deaths from burns and scalds; 7 from drowning; 14 from falls; 3 from poisoning; 8 from railroad accidents; and 25 from various other causes. More particulars of some of these accidents are as follows:

- 1. Burns and scalds. There was one each from burning house, hot bandage, clothes taking fire, naphtha stove, playing with matches, hot water, and falling into a wash-boiler; and two not described.
- 2. Falls. Three fell down stairs; two fell from a ladder; and one each fell from a coal cart, and from a coal pocket; and seree not described.
- 3. Poisoning. One each from Mrs. Winslow's Syrup; over dose of opium, and over dose of medicine.
- 4. Railroad accidents. Two each from hit by a bridge, hit by cars, fell from cars and run over, and two not described.
- 5. Various accidents. Fracture of leg, 2; fracture of thigh, 3: fracture of pelvis, 1; fracture of base of skull, 1; run over by hore cars, 2; by low-gear, 1; by carriage, 1; injury to head, 1; to hip, 1; caught in machinery, 1; choked by an orange seed, 1; gangree from frost bite, 1; thrown from wagon, 1; kicked by a horse, 2: gun-shot wounds, 1; hit by a bucket in a well, 1; crushed by a wall, 1; crushed in an elevator, 1; internal injury, 1; asphyxia, 1.

The statistics of accidental deaths, in 1883, were:-

Males, 52; females, 14; total, 66; or 2 84 per cent. of all.

American parentage, 17; foreign parentage, 49.

During 28 years, 1856-1883, decedents from accidents:-

Males, 882; females, 299; total, 1,181; or 2.90 per cent. of all.

American parentage, 447; foreign parentage, 734.

Aneurism. Seven deaths were reported in Providence from aneurism in 1883, or 0.30 per cent. of all; 5 of American and 2 of foreign parentage, all males. The locations of the aneurisms were: aorta, 4; abdomen, 2; cerebral, 1.

The statistics of aneurism for twenty-eight years, from 1856_1883, are:—

Males, 31; females, 5; total, 36; or 0.09 per cent. of all. American parentage, 16; foreign parentage, 20.

Apoplexy. The deaths from this prominent cause have increased very much in the last few years, and especially since 1880. In that year, only 48 deaths were reported from apoplexy, and that was just the average number in each of the five years preceding 1880. But in 1883, there were 84 deaths reported from this disease, and this number was 5 less than in 1882. The statistics of the decedents from apoplexy, in 1883, were as follows:—

Males, 29; females, 55; total, 84; or 3.62 per cent. of all. American parentage, 51; foreign parentage, 33.

For the 28 years, 1856-1883, the statistics were as follows:—

Total deaths from all known causes, 40,713.

From apoplexy, Males, 470; females, 501; total, 971; or 2.38 per cent. of all.

American parentage, 664; foreign parentage, 307.

Other particulars of the deaths from apoplexy, may be found under the head of "Paralysis."

Brain, Diseases of. There were 89 deaths reported under this head, in 1883, of which number, 51 were reported as "inflammation," and 11 "congestion," leaving 27 as "disease" not specified.

The 89 deaths, were 14 more than in 1882, and comprised 3.79 per cent. of all the deaths from known causes in 1883.

Of the decedents from inflammation of the brain, three-fifths were under 5 years of age, and four-fifths were under 10 years.

The decedents, in 1883, were as follows:-

Males, 46; females, 43; total, 89; or 8.79 per cent. of all.
American parentage, 43; foreign parentage, 46.

In twenty-eight years, 1856-1883, the statistics were:-

Males, 838; females, 734; total, 1,572; or 3.86 per cent. of all. American parentage, 815; foreign parentage, 757.

Bronchitis. Among acute diseases of the lungs, bronchitis is always prominent, and numbered 57 victims in 1883, or 3 less than in 1882. Thirty-one of the decedents were under 5 years of age, and 18 were over 60 years, leaving only 8 between 5 and 60 years. The statistics of bronchitis in 1883, were:—

Males, 28; females, 29; total, 57; or 2.46 per cent. of all. American parentage, 24; foreign parentage, 33.

The statistics of the decedents from bronchitis, in 28 years, 1856-1883, were as follows:—

Males, 283; females, 346; total, 629; or 1.54 per cent. of all. American parentage, 279; foreign parentage, 350.

Cancer is an important cause of death, and numbered 84 victims in Providence, in 1883, or 3.62 per cent. of all the deaths from causes that were known. The deaths from cancers were 32 more, and the per cent. 1.02 greater than in 1882. The local manifestations of the 84 cases of cancer were as follows: cancer of the breast, 14; of the stomach, 18; of the uterus, 17; liver, 9; rectum, and abdomen, 3 each; bowels and omentum, 2 each; and neck, spleen, thorax, sub-maxillary gland, chin, chest, ovary, vulva, humerus, prostate gland, bladder, and pancreas, 1 each. This leaves 4 not specified.

The decedents from cancer, in 1883, were:

Males, 26; females, 58; total, 84; or 8.62 per cent. of all.

American parentage, 47; foreign parentage, 37.

In 28 years, 1856-1883, the decedents from cancer were:-

Males, 283; females, 747; total, 1,030; or 2.53 per cent. of all. American parentage, 640; foreign parentage, 390.

Of the 1,030 decedents from cancer, in twenty-eight years, 72.52 in each 100 were females; and 62.13 in each 100 were of American parentage. The percentage of deaths from cancer has increased largely in the last twenty-five years, in Providence.

Child-birth. There were 27 deaths in Providence, in 1883, from child-birth, including 2 from puerperal convulsions, and 16 from puerperal fever. The whole number was the same as in 1882.

The 27 deaths from child-birth in 1883, were 1.17 per cent. of all the deaths from known causes. The decedents were:—

Of American parentage, 10; of foreign parentage, 17.

The decedents in 28 years, 1856-1883, were:-

488 from child-birth; or 1.20 per cent. of all. American parentage, 203; foreign parentage, 285.

The 27 decedents, in 1883, from child-birth, were 0.94 per cent. of the 2.885 mothers who here children in Providence during that year; and the 488 decedents from this cause, in 28 years, were 0.86 per cent. of the 56,866 mothers who hore children during that period.

Cholera Infantum. There were 95 deaths from this cause, in Providence, during the year 1883, or 35 less than in the preceding year.

The decedents were as follows:-

Males, 46; females, 49; total, 95; or 4.09 per cent. of all. American parentage, 37; foreign parentage, 58

The following table shows the sex, and parentage of the decedents rom cholera infantum, in Providence, in each of the 28 years, from 856 to 1883 inclusive:—

	81	EX.	PARENT	AGE.	Whole
	Males.	Females.	American.	Foreign.	Number. Percentage
1856			16	41	57
1857	. 27	23	26	24	50
L-458	. 27	22	13	36	49
l8 5 9	. 14	19	18	20	33
1860	35	33	26	42	08
861	. 29	32	25	36	61 4/
962	. 25	19	17	27	44
863	. 36	80	19	47	66 5.
864	29	29	18	40	58
865	. 20.,	30	13	37	50
866	. 30	17	15	32	47
867	. 24	25	23	26	49 1. 1.
868	. 88	32	22	48	70
869	. 85	80	27	38	65
870	. 48	45	37	56	93
871	. 30	32	26	36	62
872	. 72	79	58	93	151
873	. 42	50	89	53	اي 92ا
874	. 72	62	44	90	134
875	. 62	74	50	86	136
876	. 66	56	48		
877	. 70	52	43	79	122
878	. 85	35	26	44	70
879	. 80	29	2 8	31	59
880	. 42	58	41	59	1004
881	. 58	44	85	67	1024
882	. 66	64	50	80	130
883	. 46	49	87	58	954

Of the 95 decedents, in 1883, from cholera infantum, 75 were under one year of age, and 17 between one and two years.

Of the 2,235 decedents, in 28 years, 830, or 37.14 per cent. were of American parentage, and 1,405, or 62.86 per cent. were of for eign parentage.

Some statistics of cholera infantum, in connection with other diarrhoal diseases, will be found hereafter under the head of "Diarrhoan and Dysentery."

Cholera Morbus. There were 11 deaths from this cause in 1883. or 2 less than in 1882. The annual number of deaths from cholera morbus, has varied, during the last 29 years, from 1 to 17, the annual average being only 5.

The decedents, in 1883, were:-

Males, 5; females, 6; total, 11; or 0.47 per cent. of all. American parentage, 5; foreign parentage, 6. In the last 28 years, 1×56-1883, there were 144 deaths reported from cholera morbus, or 0.35 per cent. of all from known causes.

Consumption. There were 364 deaths reported in Providence from consumption, during the year 1883, or 13 more than in 1882. The percentage from consumption, of all the deaths in 1883, from known causes, was 15.68, or 0.04 less than in 1882.

Estimating the population of Providence at 119,405, for the middle of the year 1883, we find that there was one death from consumption in 328 of the inhabitants. In 1882, the rate was one in 326, or 3.05 in each 1,000; and in 1881, it was one in 312, or 3.20 in each 1,000. Probably, to a considerable extent, on account of more complete and accurate records, there has been a large decrease in the rate of mortality from consumption. During the 15 years, 1840–1854, the average annual rate was one death from consumption in 209.9; during the twenty-eight years, 1856–1883, the average annual rate was one death from consumption in 311.7 of the population of Providence.

The decedents, in 1883, were as follows:-

American parentagenales, 47females, 60total, 10 Foreign parentagemales, 119fem: les. 138total, 22
Totalsfemales. 166females. 196total, 36
The general statistics of consumption, during 28 years, from 1856 to 1883 inclusive, are as follows:—
Total deaths from known causes, 40,713; from consumption, 6,749; or 16.58 percentantal average, 241.
American parentage
Totals
Males, 45.62 per centfemales, 54.38 per cent. American parentage, 44.61 per centforeign parentage, 55.39 per cent

The number and percentage, according to parentage, of the dececlents from consumption, in periods of five years each, since 1856, and for the years 1881, 1882, and 1883, were as follows:—

•	American :	Parentage.	Foreign Pa	rent age .	Deaths from
	Number.	Per cent.	Number.	Per cent.	Consumption
1856 to 1860	525	56.45	405	43.53	93
1861 to 1865	495	50.77	480	49 23	97
1866 to 1870	583	50.57	520	49.43	1,05
1871 to 1875	558	44.78	682	55.21	1 ,23
1876 to 1>80	570	88.05	928	61.95	1.49e
1881	116	81.72	228	66 28	34
1882	118	82.19	238	67.81	35
					3K4
28 years	8.011	44.61	8.788	55.39	6.741

This table shows that, of all the decedents from consumption in Providence, the *percentage* of American parentage has largely decreased since 1856, while the percentage of foreign parentage has correspondingly increased.

The next table shows the number of decedents, from consumption. by parentage, in each year since 1856. We find from that table that the actual number of decedents of American parentage, from consumption, has not increased since 1856, notwithstanding the increase of population. In 1856, there were 104 decedents from consumption, of American parentage; in 1857, there were 120; in 1860, there were 123, and that is the largest number in any year since 1856, and in 1883, there were only 107.

On the contrary, the decedents, from consumption, of foreign parentage, have increased enormously since 1856. In that year, the number was 74 of foreign parentage, and in 1883 the number was 257. The decedents of American parentage, from consumption, in 1883, was only 3 more than in 1856, while those of foreign parentage, from the same disease, were 183 more.

In my report for 1880, it was shown that the ratio of decedents, of American parentage, in 1856, was one death from consumption in 268 2 of the living population of American parentage; while in 1880 the ratio was only one death in 435 2.

On the other hand, the ratio from consumption of the decedents of foreign parentage, was one in 262.6 of the living population of foreign parentage, in 1856, and one in 268.0 in 1880. See City Registrar's Report for 1881, pages 63, 64, and 65.

The following table gives various important particulars in regard to the decedents from consumption, in Providence. in each of the twenty-eight years, from 1856 to 1833 inclusive:—

DEATHS FROM CONSUMPTION IN PROVIDENCE, TWENTY-FIGHT YEARS, 1856-1883.

Nex; parentage; season; age; whole number, and percentage from known causes.

(WHEL MITTHEN	1956, 1857.	1817.	3	1800	1800	1081	1865	1963	1861.	1863.	1866.	1867.	1808	1860.	1870.	871.	1879.	873. 1	1874. 18	1875. 18	1876. 18	877. 18	1878. 1879.	9. 1880.	1881	1. 1882.	_	883
it hole number		1 KT 1 KT 1 KT 1 KT 1 KT 1 KT 1 KT 1 KT	16.97	1 187	118	1 22	1 E 2	1 88.71	10.5	18.40	19.81	80.08	100	210	9.00	188	18.9	8,531	750	73.0	25.	83	1 88	25 35 35 35 35 35 35 35 35 35 35 35 35 35	322 344			3A1 364 5.72 15.68
Ki Malora Permalora		100	28	25	12	28	3.2	23	801	ZE	950	28	82	117	101	8=	12	82	128	910	68	525	82	195	154	167 166	9.9	
Invitan.	Er.	NA AM	22	18	100	82	88	25	102	5.5	58	85	108	861	1111	88	55	55	100	120	819	98	100	25	===	116	213	107
	A*LINIEIIAA	MARKET ELES	お工事の見事力ニュニュ	SSECRESCREE	121211111221	2122252222	111111111111	**********	12811111111	2:22×2::2222	SECSESSESSE	STRITESSTEE	22222222222	*************	822822822822 822822822	787772222778	56885868888	usisseessiau	88888888888	58585555555555555555555555555555555555	81:83882828888	555555555555	HARRAGANNEAN	808882366888	202522333333333333333333333333333333333		22222222222	232453332333

DEATHS FROM CONSUMPTION IN PROVIDENCE. TWENTY-EIGHT YEARS, 1856-1883.—CONTINUED.

Sex; parentage; season; age; whole numler, and percentage from known causes.

Total.	1,752	9,350 11,23 12,238	85 66 66 66 66 66 66 66 66 66 66 66 66 66
883	8528	855	200円のの記録を登録して
1883.1	2258	187 99	**************************************
1881.	8888	252	***********
1880.1	8288	182	+++-#8E66E8c-
.6281	8882	288	-2445886488
1878.	2233	201	-4-4g2256885a
1817.	1135	200	4-44-68-48-
1876.	SZER	300	44-roll 2 4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
1875	ZEZE	588	n un 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
1874.	2523	88.5	00004-25-52 = :
1873.	22.4%	788	-80 -284587
878	2883	818	に4~90点岩色工器 工程をお
1871. 1872.	8448	138	x-1-x 842252
. 1870.	9889	\$85	000-0008C-\$285-
.698	8228	22.5	**********
1868.	\$228	282	- : : : : : : : : : : : : : : : : : : :
1867.	8=45	582	*** **********************************
1868	2849	682	-a-4475252524
. 2982	5248	333	E00040222250000
1864	2232	583	88+84B\$\$\$\$
1803.	4848	152	
1862.	6886	222	******************
1861.	95 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	588	4-0-055525250
1860.	8888	255	0 0 4 1 4 2 5 4 8 2 E 5 u
1859.	25.55	828	84-845842X84
1858.	3233	848	***********
1857.	85448	125	s<252227Ess
1856. 1857. 1858. 1859.	55458	553	o :
CONSUMPTION.	frst quarter econd quarter hird quarter	record 4 months	68: Under 1 year 2 m 10. 10 m 10. 15 m 15. 15 m 15. 16 m 16. 16 m 16. 16 m 16. 17 m 16. 18. 18. 18. 18. 18. 18. 18. 18. 18. 18

Percentage of consumption. The preceding table shows that of all the deaths in 28 years, from known causes, 16.58 per cent. were from consumption; the percentage varying from 13.57, in 1873, to 22.60, in 1860. During the last ten years, the percentage has not exceeded 16.15, in any year.

During the whole period of 15 years, previous to 1855, the percentage from consumption was 22.11 in each 100 of all the deaths from causes that were known.

The decrease in the percentage from consumption, since 1855, has undoubtedly been caused, to some extent, at least, by more accurate diagnosis and records.

Season and consumption. It has not been generally supposed that season had much influence upon the mortality from consumption; but the figures in the preceding table seem to indicate that such influence does exist. In the aggregate of deaths from consumption in 28 years, 1856–1883, the smallest number of deaths in any month, was in August, 517; and the largest number was in January, 611. In fact, the monthly number of deaths increased regularly from August to January, and decreased regularly from January to August, the only exception being the month of February which is two or three days shorter than the other months.

The deaths from consumption in the different quarters of the year, also show the largest numbers in the first and fourth quarters, and much the smallest number in the third quarter. It would seem that cold weather has some influence, in this climate, in increasing the mortality from consumption.

Convulsions. Sixty deaths were reported, in 1883, as caused by convulsions, or 8 more than in 1882. The decedents were:—

Males, 30; females, 30; total, 60; or 2.59 per cent. American parentage, 24; foreign parentage, 36.

During 28 years, from 1856 to 1883 inclusive, the whole number of deaths, from known causes, was 40,713, of which 1,046, or 2.57 per cent., were reported as caused by convulsions. The decedents from convulsions were:—

. Males, 569; females, 477; total, 1,046. American parentage, 486; foreign parentage, 610.

ij

The larger portion of the decedents from convulsions are under 5 years of age, and the term itself is only a symptom of some unknown cause.

Croup. There were 35 deaths reported from croup, in Providence, in 1883, or 3 more than in 1882. The rate was 1.51 per cent. of all the deaths from known causes.

By season, the deaths from croup, in 1883, were as follows:-

1st quarter12	3d quarter 3
2d quarter 6	4th quarter14

The decedents from croup, in 1883, were:

Males, 17; females, 18; total, 35; or 1.51 per cent. of all.

American parentage, 13; foreign parentage, 22.

In 28 years, 1856-1883, the decedents from croup, were:-

Whole number, 883; annual average, 31.

Males, 440; females, 443; total, 883; or 2.17 per cent.

American parentage, 358; foreign parentage, 525.

The next table gives the statistics of croup in Providence, for 28 years, 1856 to 1883 inclusive, showing the sex, color, and parentage of the decedents; the season and locality of the deaths; and the percentage from croup of all the deaths from known causes:—

DEATHS FROM CROUP IN PROVIDENCE, TWENTY-EIGHT YEARS, 1856-1883.

Total.	883 2.17	33	858 525	228823882438833	82 198 198 198 198 198 10	22
1883.	82.	118	5 83	P88-884 : 824-4	B : 04 : - 04 04 05 05	_
88	33	181	78	DH44H : : 1000	00 : H 04 H - 15 4 F- 00	_
1881	8.30	នន	33.1	F-6800000-46F-F-	8 : C 2 6 - C 4 F Q	-
1880	87:	20 25	r- 81	Ø : 27 : : : : : : : : : : : : : : : : :	84 -4-04	
1879.	25.5	16	8122	⊕ 12 30 4 94 94 4 1 − 94	∞ -4 -∞ -∞ -∞	_
	88	8 38	88	4000 : : :NOO	@-8-03147I	9
1877. 1878	2.07	£ 24	78	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	F-04-4 : 04 : 04 00 00 34	_:
1876.	3.5	22 23	20.00	B 95 9: 1:00 5	œ⊣x4≈00€5	_
1875.	88	ន្តន	33		e :6 :4 : :448	_
1874.	1.48	80	12		4 8-46-4	_
1873.	1.83	6.5	15.6	900000000000000000000000000000000000000	4-4	
1872.	1.2	13	213	8668969 9148	4:1-01411180	_
1871.	28.88 88.88	18	85	DONAL RELATE	œ	_
	25.16	16	22	44	O - C 20 : 41 41 40 50	69
1869. 1870.	1.53	5 9	8 81	mp		_:
1867. 1868.	=5.	60	41-		0 0 -20-	
1867.	2.01	62	60	M - M - M - M - M - M - M - M - M - M -	40.00 - 000	
1866.	82.5	F 2	122	# # # # # # # # # # # # # # # # # # #	8 9 6 8 4 4	
1865.	84	18	Ф 8			_
1884.	3.59		83		8 5 4 8 8 8 2 5 · · · · · · · · · · · · · · · · · ·	
1863.	2 28 28	83	=3	::		
1862.	8.5	**	15	:::	************	_ <u>:</u>
1861.	120	21	30			
1860.	1.88	-=	72			
1856. 1857. 1858. 1869.	25.	8,2	22	+++con	01 01 01 01 01 01 01 01 01 01 01 01 01 0	1
1858	2.71	212	138	80-4-80-4-2000 44402314		_
1857.	32.5	22	82			
1856.	39.20	16	~ &	41001-01-03 33014	91-85-64-5	
CROUP.	Whole number	SEX: Males Females	PARENTAGE: American Foreign	Belson: January January Kebruary March. April. April. Angust June July June July June July Lugust Beptember. October. December.	Mard II Ward II III IV. VII VII VII VII VII XX	Colored

A careful study of the figures in the preceding table would furnish the grounds for lengthy comments which we have not time nor space for, at present. A few words must suffice.

It will be noticed that the number of deaths, and the percentages do not show any evidence of an epidemic prevalence of this disease. The number of deaths is greater in some years than in others; and is especially greater in years when diphtheria was epidemic, indicating, perhaps, that in the diagnosis croup was sometimes confounded with diphtheria. From this fact, and others which we do not refer to at this time, we are justified in the opinion that: True croup never prevails as an epidemic. In this respect, it differs very markedly from diphtheria.

Croup depends very much upon season, and especially upon temperature, increasing in the autumn as the weather grows colder. Notice in the table that, in the aggregate for 28 years, there were only 12 deaths in August, while there were 128, 122, and 123 respectively in November, December, and January. The total number in 28 years, in the four quarters of the year, were as follows:—

First quarter...... deaths, 311

Second quarter......deaths, 143

Third quarter.....deaths, 339

Fourth quarter.....deaths, 339

A comparison of the mortality, with respect to season, of the three diseases, croup, diphtheria, and scarlatina, will be found hereafter under the head of "Diphtheria."

Debility. This word gives very slight, if any, information in regard to the real cause of death. In fact it is only a symptom, or a result of the real cause. Nevertheless we find "debility" given as the cause of a considerable number of deaths, every year. In 1883, there were 27 deaths reported from debility, of which 13 were "debility from premature birth." The 27 decedents, in 1883, were as follows:—

Males, 10; females, 17; total, 27; or 1.16 per cent of all. American parentage, 14; foreign parentage, 13.

For the 28 years, 1856-1883, the statistics of debility were:-

Whole number of deaths, 689; annual average, 25.

Males, 363; females, 326; total, 689; or 1.69 per cent. of all.

American parentage, 294; foreign parentage, 395.

Diarrhæa and Dysentry. There were 75 deaths reported from these diseases, in 1883, or 11 more than in 1882; the rate was 3.23 per cent. of all from known causes, or 0.37 per cent. more than in 1882. These and other diseases usually called "diarrhæal," are included in the following table:—

	Males.	Females.	American.	Foreign.	Total.	Per cent
Acute Diarrhœa	19		27	25	52	2.2
Chronic Diarrhœa .	4		4	3	7	0.3
Dysentery	6	10		11	16	0 #
Cholera Infuntum						
Cholera Morbus						
Enteritis						
Enteritis, Gustro						
Enteritis, Colo						
Totals	95	117	92	120	212	9.1

The total decedents from all these causes, were:-

Males, 95; females, 117; total, 212; or 9.13 per cent. of all. American parentage, 92; foreign parentage, 120.

The whole number is 44 less than in 1882, and the rate is 2.32 less; the number of males, 36 less, of females, 8 less; of American parentage, 18 less, of foreign parentage, 26 less.

The decedents from all these diseases, during the 28 years, from 1856 to 1883 inclusive, were as follows:—

Number of decedents, 4,430; annual average, 158.
 Males, 2,204; females, 2,226; total, 4,430; or 10.88 per cent, of all.
 American parentage, 1.811; foreign parentage, 2,619.

The proportions of decedents, according to parentage, from diarrhaal diseases, were as follows:—

	Foreign	. A1	Loticed.
1883			

In the total decedents, in 1883, from all causes, the proportions were 122.3 foreign to each 100 Americans; or 55.03 foreign, and 44.97 American in each 100 decedents.

The following table shows the number of deaths from each diarrhocal disease, and the percentage of the total of the deaths from all these diseases, in different periods of time, from 1840 to 1983 inclusive:—

	Cholera	Cholera					Per cent. of all from
	Infantum.	Morbus.	Diarrhœa.	Dysentery.	Enteritis.	Total.	known causes
1840-1854	560	63	349	796	102	1,870.	16.77
18 5 5–1859	250	14	159	127	23	573.	
1860-1864	297	17	154	189	46	703.	
18 65 –18 6 9	281	35 .	125	187	37	665.	
1870-1874	532	26	185	91	97	981.	12.0
1875-1879.	509	24	169	77	104	888.	9.1
1880	100	4	30	12	19	165.	7.9
							8.5
							9.Ji
44	0.056		1 914	1.500			1210

The deaths from Asiatic cholera are not included in these figures. They were as follows: 158 in 1849; 159 in 1854; and 14 in 1866; total, 331.

Diphtheria. There were 54 deaths reported from diphtheria, in 1883, a smaller number than in any other year since 1875, except 1882, when there were 48 deaths. The percentage in 1883, was 2.33 per cent. of all the deaths from known causes.

The sex and parentage of the decedents from diphtheria, in 1883. were as follows:—

Males, 20; females, 34; total, 54; or 2.33 per cent. of all.
American parentage, 27; foreign parentage, 27.

The first deaths from diphtheria in Providence, in recent times were in 1858. During 26 years, from 1858 to 1883 inclusive, there were 1,372 decedents from diphtheria, as follows:—

Decedents from all known causes in 26 years, 38,785.

From diphtheria, 1,372; annual average, 53.

Males, 680; females. 692; total, 1,372; or 3.39 per cent. of all.

American parentage, 674; foreign parentage, 698.

There seems to be a curious difference in the parentage of the decedents from diphtheria depending upon the fact whether the disease is epidemic or not.

Thus in the period from November, 1876 to 1879, when the disease was epidemic, the decedents from diphtheria were 56.39 of foreign parentage, and 43.61 of American parentage, in each 100.

But from 1858 to October 1876, during which time the disease did not become epidemic, the decedents from diphtheria were 58.68 of American, and 41.32 of foreign parentage in each 100, the figures when the disease was epidemic being almost exactly the reverse of the figures when it was not epidemic.

Since the end of the epidemic, in 1879, the American class is again in excess, thus: in 1880, 57.38 per cent. American, 42.62 foreign; in 1881, 50.86 per cent. American, 49.14 foreign; in 1882, 54.17 American, 45.83 foreign. In 1883, the numbers, American and foreign, were equal.

The following table shows the percentage, season, and locality of the deaths from diphtheria; also the sex and parentage of the decedents, in each of the 26 years, from 1858 to 1883 inclusive:—

	1858-	HERIA. 1888. Sars.	9CARLATINA. 1855-1883. 29 years.		CROUP. 1855–188 8. 29 years.	
Months.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent.	Number of Deaths.	Per cent
January	128	9.88	266	12.53	195	18.71
February	90	6.56	231	10.88	98	10.75
March	106	7.78	198	9.09	98	10.75
lst quarter	824	28.62	690	82.50	321	35.21
April	96	7.00	198	9.09	72	7.89
May	90	6.56	199	9.87	49	5.37
June	82	5.98	166	7.82	81	8.40
2d quarter	268	19.54	558	26.28	152	16 66
July	61	4.45	118	5.82	30	3.29
August	64	4.66	90	4.24	12	1.31
September	119	8.67	95	4.48	50	5.48
3d quarter	244	17.78	298	14.04	92	10.08
October	198	14.48	139	6.55	91	9.98
November	188	18.70	187	3.81	180	14.95
December	150	10.98	251	11.82	126	18.82
4th quarter	586	89.06	877	27.18	847	38.05
Totals	1,872	100.00	2,128	100.00	912	100.00

The preceding table shows the figures for each quarter of the year. If we divide the year into periods of four months, the figures are as follows:—

	DIPHT	HERIA.	SCABLATINA.		CROUP.	
SEASONS.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent
1st 4 months	490	80.62	888	41.59	898	48.10
2d 4 months	297	21.65	568	26.75	123	18.37
3d 4 months	655	47.78	679	81.66	897	48.52

Dividing the year into periods of six months each, the figures are as follows:—-

	DIPHT	HERIA.	SCARL	ATINA.	CROUP.	
SEASONS.	Number.	Per cent.	Number.	Per cent.	Number.	Per cent.
1st 6 months		43.16 56.84	1,248 876	58.78 41.92	473 439	51.87 48.13

The most casual reference to the preceding tables, cannot fail to show very great differences in the three diseases named, with respect to season. Let us notice a few things.

The following statement will contrast the three diseases in regard to season and mortality:—

	DIPHTHERIA.	SCARLATINA.	CROUP.
Percentages.	Per cent.	Per cent.	Per cent.
Largest monthly			
Largest quarterly	Fourth qr89 06	First qr32.50	Fourth qr38.65
Smallest quarterly	Third qr17.78	Third qr14.04	Third qr10.06
Largest 4 months	Third 4 mos47.73	First 4 mos41.59	Third 4 mos43.53
Smallest 4 months	Second 4 mos. 21.65	Second 4 mos.26.75	Second 4 mos.13.37
Largest half yearly	Second 6 mos.56.84	First 6 mos58.78	First 6 mos51.87
Smallest half yearly	First 6 mos. `43.16	Second 6 mos.41.22	Second 6 mos.48.13

An examination of the preceding tables shows marked differences in the mortality of the three diseases given, with respect to season. In the aggregate for the long periods named, we find the greatest mortality to be: of diphtheria, in October; of scarlatina, in January; and of croup, in November.

Diphtheria shows an almost uniform increase from July to October, and then a uniform decrease from October to July.

Scarlatina shows a uniform increase from August to January, and then a uniform decrease from January to August.

Croup shows an increase from August to November, and then a decrease from November to August. But croup seems to depend more upon cold than either of the other diseases. There is but a very slight difference in the percentages from croup, in the three months November, December, and January; and February and March have both a large percentage from this disease.

Diphtheria has 13.68 per cent. more of deaths in the last half of the year than in the first half; while on the contrary, scarlatina has 17.56 per cent. more in the first half than in the last half of the year. The difference in croup is comparatively slight.

Similar and other illustrations of the differences between these diseases, were given in the reports for 1860, 1870, and in those from 1877 to 1880 inclusive. The subject is of very great interest and importance.

Dropsy. This symptom of disease was reported as the sole cause of 4 deaths, in 1883; one male, 3 females; all of American parentage; total, 4, or 0.17 per cent. of all from known causes.

Erysipelas. There were 14 deaths reported from erysipelas, in 1883; the same number as in 1882. The decedents were:—

Males, 7; females. 7; total, 14; or 0.60 per cent. of all. American parentage, 6; foreign parentage, 8.

During 28 years, 1856-1883, the decedents were:-

Males, 105; females, 119; total, 224; or 0 55 per cent. of all. American parentage, 122; foreign parentage, 102.

Fever, Typhoid. The severe epidemic of typhoid fever, in Providence, in the autumn of 1882, was renewed in March and April 1883. There were 15 deaths in January, 6 in February, 11 in March, and 32 in April. There was also the usual increase of typhoid fever in the autumn of 1883, giving 16 deaths in September, and 12 in October. The result was that, in the whole year 1883, there were 128 deaths in the city from this disease, a greater number than was ever before known, in a single year, except in 1882.

The decedents, in 1883, were as follows:-

Males, 72; females, 56; total, 128; or 5.51 per cent. of all. American parentage, 54; foreign parentage, 74. During 28 years, 1856-1883, the decedents were:-

Whole number from typhoid fever, 1,381; annual average, 49. Males. 709; females, 672; total, 1,381; or 3.39 per cent. of all. American parentage, 729; foreign parentage, 652.

In past years, there have been occasionally, deaths reported as caused by "fever," and some as "bilious fever," and about 30 in all as "typhus fever." There was also, in 1869, one death from "intermittent," and one from "malarial" fever; and in 1880, one death from "malarial" fever. These were all included with typhoid fever during the 25 years previous to 1881. Since 1880, with the new and increased development of malaria, in this vicinity, the deaths from malarial, remittent, intermittent fevers and others of this class, have been kept separately. There were three in 1881; 5 in 1882; and 6 in 1883, of what may be termed the malarial class of fevers.

The following table shows the statistics of typhoid fever, in Providence, for each of the 28 years, 1856–1883; including the whole number, percentage, season, and locality of the deaths; and the sex, color, parentage, and ages of the decedents:—

DEATHS FROM TYPHOID FEVER IN PROVIDENCE, TWENTY-EIGHT YEARS, 1856-1883.

Sex; parentage; color; season; age; locality; whole number, and percentage from known causes.

. Total.	1,381	279 56 673		99 1	255 - 25 - 25 - 25 - 25 - 25 - 25 - 25
1883.	128		25		
1882	140	27.08	2.8	60	488468 :49×5
1881.	38	18	201	04	8 8 4 5 5 5 15 15 15 15 15 15 15 15 15 15 15
1880.	2.57	88	23	-	9-6464-92
1879.	1.98	22	14.	-	
1878.	74.88	28	202	60	94 - 0404 4 4 2 2 2 2 2
1877.	55.55	88	8151	28	0 :00 H H W 1-262
1876. 1	1.78	17	15	C1	0100 : :01 :0.000
1875. 1	12.5	88	22	01	m :28
1874. 1	3.06	933	88	12	G : MOINT G251
1873. 1	25.56	83	23 23	9	* was : 4 w + 5 E
1872. 1	4.31	4%	20.00	63	a : more or or 2 = -2
1811.11	28	88	88	63	C+N-4841-003
1870. 1	97.1	98	85	69	u :44-404055
11 889	1 88	3153	85 53	1	
1868.	25	13	92 8	Ì	
1867.1	8.69	138	813	61	
1866.1	1 94	200	128	1+	
1865.13	8.8	28	33	*	
1864. 18	5,55	28	125	-	01-2-032-020-
- 11	17.58	202	14.	0	DE- :01 :01-1-43
1862. 18	2.8	21	7.	1	01- 01 : 01- 01C
1861.18	1 86	40	54	C4	01 :-0101 :
860.18	88	100	25	01	004 : 101-04 :04-4
	1.86.	18	26 11	:	
- 88	24.4	20	55 0		
57. 18	1 34	119	14	÷	
- 18	8.80	22	22	59	
1 18	04				
YPHOID FEVER, 1856, 1857, 1858, 1859	Thole number	KX: Males Females	ARENTAGE: American Foreign	olored	EASON: January January March April June July August September: Roycember: November:

Sex; parentage; color; season; age; locality; whole number, and percentage from known causes.

31, 1882, 1883, Total.	13 10 64 281 11 15 22 300 14 115 42 800	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	4 : 7 4 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8
1880. 1881	225	H : ::::::::::::::::::::::::::::::::::	@#####################################
1879.	983	co.4Ep==40	-84 -2866
1878.	130		e : www.4e54
1677.	1-52	1042000000000000	romen and a
1876.	0.2		31-12-10-10-10-10-10
1875.	6003	unacolitana	ಬರದಬಹೆದಬಹ=
1874.	5 00	HERDE TO SEE	4-5xxxxxx
1873.	339		-621-061-40
1879.	1.23		
1871.	222	- :	81012424000
1870.	02 8 25		10 10 10 10 10 10 10 10 10 10 10 10 10 1
1869.	1-58	:	00 11 C 01 C 01 C 00 C 00 C
1868.	169.6	oi : : : : : : : : : : : : : : : : : : :	P-0103-000401-
1867.	100	. :4+15I	6 46666-0
1866.	× 51.55	.4004040004	5-0-004
1865.	148	- 80809520899	8688562
1864.	555		Seasa Cor- : :
1863.	2122	H=0x22-40	4040000:
1862.	2000		- : caro-so
1861.	222		40,40,004
1860.	1050		W-1201-4-4-1
1869.	1691		*******
1858.	108.6	H : : : : : : : : : : : : : : : : : : :	4 01004-
1837.	401		1-01-000-4-4
1856.	868	- 84 678484	90 94 00 04 40 00 A4
TYPHOID FEVER. [1850, 1857, 1858, 1850.	First 4 months Second 4 months Third 4 months	AGE: Under J year 1 and under 2 2 and under 2 15 and 10 16 and 20 20 and 20 60 and 20	LOCALITY: Ward III IV V V V V V V V V V V

• 17 were in Rhode Island Hospital. † 14 were in Rhode Island Hospital.

The preceding table gives very full particulars of the deaths and the decedents from typhoid fever. The general and comparative results, for the whole period of twenty-eight years, may be expressed plainly and briefly as follows:—

Percentage. In 1876, and in 1881, the deaths from typhoid fever were 1.78 per cent. of all the deaths from causes that were known. In 1882, the percentage was 6.27, and these were the extremes during the whole period. The average percentage, for the whole twenty-eight years, was 3.39.

Sex. In the whole period, from 1856 to 1883 inclusive, there were 709 male, and 672 female decedents from typhoid fever. This is at the rate of 105.5 males to each 100 females; or 51.34 males, and 48.66 females in each 100 decedents from this disease.

Parentage. Of 1,381 decedents, in 28 years, from typhoid fever. 729 were of American, and 652 of foreign parentage; or 111.8 of American to each 100 of foreign parentage; or 52.79 American and 47.21 foreign in each 100 decedents.

Season. Typhoid fever is especially a disease of the autumn, and its annual increase at that season is always expected. Of the whole number, 1.381 deaths from typhoid fever, in 28 years, the number and percentage in each quarter of the year, were as follows:—

	Number.	Per cent.		Number.	Per cent.
1st quarter			_		
3d quarter	348	25.20	4th quarter	617	4.0

The fourth quarter of the year, has many more deaths from typhoid fever, and a much greater percentage, than either two of the other quarters.

If we divide the year into three sections of four months each, the number and proportions of deaths from typhoid fever, were as follows:—

·	Number.	Per cent.
First four months	281	90.36
Second four months	3 00.:	21.72
Third four months	800	57.92

Considerably more than half of all the deaths from typhoid fever are in the last third of the year.

Dividing the year into two parts, we find the number and proportion of the decedents as follows:—

	Number.	Per cent.	Number.	Per cent.
ist 6 months	416	30.12	2d 6 months 965	69.88

These illustrations make the fact very plain, of the great excess of mortality from typhoid fever in the last half of the year, and especially in the last four months.

The following table shows the number and percentage of the decedents from typhoid fever, in each division of age, in the aggregate mortality for the year 1883, and also for the 28 years, from 1856 to 1883 inclusive:—

PERIODS.	Under 1 year.	1 and under 2.	2 and under 5.	5 and under 10.	10 and under 15.	15 and under 20.	20 and under 30.	30 and under 40.	40 and under 50.	50 and under 60.	60 and under 70.	70 and under 80.	80 and under 90.	90 and over.	Total.
1883. Number Per cent	0.78	2.56	3.13	8 6.25	13	23 17.97	43 33.59	19	1.15	8 6.25	3 2.34				128 100.00
1856-1883. Number Per cent	9	-		1	106	224 16.22	399 28.89	193 13.98	1	1	57 4.13	1		0.07	1,381

Of the 128 decedents, in 1883, from typhoid fever, 85, or 66.40 per cent. were between 15 and 40 years of age. Of the 1,381 decedents in 28 years, 816, or 59.08 per cent. were between 15 and 40 years.

The next table shows the number of deaths, and the aggregate and average age of the decedents, in each of the twenty-eight years, from 1856 to 1863 inclusive:—

Decedents from typhoid fever; aggregate and average age.

	Aggregate Age.				Average Age.			
YEARS.	Number o Deaths.	Years.	Mos.	Days.	Years.	Mos.	lwys	
356	80	788	5	3	26	3	11	
357	30	781		13	26		12	
58	24	829	5	23	34	6	21	
359	87	1 252	7	18	33	10	1 1	
360	20	708	6	19	35	3	1 2	
961	23	722	8	7	31 '	4	3	
62	21	540	9	11	25	9		
963	47	1,456	11	28	30	11	#	
864	42	1,360		27	32	3]:	
865	82	2.262	1 4	34	27	7	1	
846	46	1,420	2	17	30	10	1	
867	33	654	4	25	19	9	, 2	
868	24	692	7	8	28	10		
×69	35	1,214	4	13	34		1	
870	52	1,421	1	7	27	3	1	
871	53	1.610	11	15	30	4	1	
372	68	2,161	8	19	31	y	1	
873	61	1,672	6	10	27	5	! .	
574	60	1,553	11	24	25	10	i :	
575	71	1,938	5	19	27 27	3	1	
₹76	83	922	8	17	27	11	j 1	
377	43	1,243	11	23	28	11	1	
٠78	47	1,151	9	27	24	6	ł	
879	40	1.206	9	20	30	2	١.	
880	53	1,505	2	4	23	4	1 1	
×81	38	1,156		13	30	. 5	_	
882	· 140	8.59∺	I .	5	25	8	: 1	
883	128	8.138	11	2	24	6		
8 years	1.381	38,964	4	6	28	2	i i	

The average age, as well as the percentages at different ages shows that the great majority of the decedents from typhoid fever are in the prime of life.

Heart, Diseases of. During the year 1883, there were 172 deaths from diseases of the heart, in Providence, including 3 from "hypertrophy," which are given separately in the table XIX. Among the different kinds of diseases of the heart, named in the returns in 1883. were 8 specified as "fatty degeneration;" 11 "endocarditis;" 2 "pericarditis;" 3 "angina pectoris;" 3 "dilatation;" 2 "endopericarditis;" 1 "dropsy;" 2 "paralysis;" 1 "valvular."

The decedents from diseases of the heart, in 1883, were:-

Males, 93; females, 79; total. 172; of 7.41 per cent. of all. American parentage, 87; foreign parentage, 85.

In the twenty-eight years, from 1856 to 1883 inclusive, there were

2,017 deaths from diseases of the heart, in Providence, or 4.95 per cent. of all the deaths from "known" causes.

The following shows the particulars of the decedents from diseases of the heart, in periods of five years each from 1856 to 1880, and annually since 1880.

	SE	х.	I'AREN	TAGE.	ber.	Percentage.	
PERIODS.	Males.	Females.	American.	Foreign.	Whole Number		
1856-1860	62	. 81	104	39	143	3.01	
1861-1865	116	121	169	68	237	4.29	
1866-1870	131	131	164	98	262	4.73	
1871-1875	218	248	277	184	461	5.50	
1876-1880	239	240	271	208	479	4.87	
1881	57	64	65	56	121	5.68	
1882	59	83	78	64	142	6. 36	
1883	93	79	87	85	172	7.41	
28 Years' Total	975	1,042	1,215	802	2,017	4.95	

In the report for 1882, page 85, these particulars were given for each year separately.

One important fact is shown, beyond all controversy, by the figures of the preceding table, in connection with the table of annual deaths in last year's report. It is this:

There has been a very great actual and relative increase in the mortality from diseases of the heart, in Providence, during the last twenty-eight years.

In 1856, there were 21 deaths from these diseases; in 1883, there were 172 deaths. In 1856, the deaths from diseases of the heart were 2.03 per cent. of all the deaths from known causes; in 1883, they were 7.41 per cent. The smallest number, and the smallest percentage of deaths from diseases of the heart were in 1856; the largest number and largest percentage were in 1883.

Besides this, there has been an increase, between every two periods named in the preceding table, with one exception.

Between 1856 and 1883, the decedents from diseases of the heart increased 719 per cent.; the decedents of American parentage increased 444 per cent.; those of foreign parentage increased 1600 per cent.; and the percentage of deaths from diseases of the heart, increased 265 per cent.

The causes of this great increase of diseases of the heart, present a subject of much importance and interest; but which we are unable to investigate at this time.

Hemorrhage. This was reported as the cause of twelve deaths in 1883; of which six were hemorrhage from the lungs; two were umbilical hemorrhage and the others were not specified. The decedents were:—

Males, 8; females, 4; total, 12; or 0.52 per cent. of all.

American parentage, 7; foreign parentage, 5.

Hernia. There were three deaths from this cause in 1883; 4 less than in 1882, and 0.18 per cent. less.

Males. 1; females, 2; total, 3; or 0 13 per cent. of all.

American parentage, 2; foreign parentage, 1.

Hooping Cough. There were only 4 deaths from hooping cough in 1883; in each of the two preceding years there were 40 deaths from this cause. All the decedents in 1883 were less than two years old. They were:—

Males, 2; females, 2; total, 4; or 0.17 per cent. of all.

American parentage, 4; foreign parentage, 0.

The statistics for the twenty-eight years, 1856-1883, are:-

Whole number, 481; annual average, 17.
Males, 208; females, 273; total, 481; or 1.18 per cent. of all.
American parentage, 213; foreign parentage, 268.

Hydrocephalus was reported as the cause of 32 deaths in 1883; 4 more than in 1882; and 0.13 per cent. more. All but four of the decedents were less than five years old. The sex and parentage were as follows:—

Males, 15; females, 17; total, 32; or 1.38 per cent. of all.
American parentage, 17; foreign parentage, 15

For the twenty-eight years, 1856 to 1883; the decedents from Hydrocephalus, were:—

Whole number, 854; annual average, 30.

Males, 467; females, 387; total, 854; or 2.10 per cent. of all.

American parentage, 439; foreign parentage, 415.

Inanition. There were 7 deaths in 1883, of which the reported cause was inanition; in 1882, there were 4, and that was the first time that it has appeared in the list of causes. In 1882, all the decedents were males, less than a year old; one of American and three of foreign parentage. In 1883, six were under one year, and one was between 60 and 70 years old. There were 5 males and 2 females; 3 of American, and 4 of foreign parentage; total 7, or 0.30 per cent. of all from known causes.

Insanity. The number of deaths from insanity in 1883, was 18; 6 more than in 1882, and 0.24 per cent. more. The decedents were:—

Males, 7; females, 11; total, 18; or 0.78 per cent. of all. American parentage, 11; foreign parentage, 7.

For the twenty-eight years, 1856-1883, the decedents reported from insanity, were:—

Whole number, 302; annual average, 11.

Males, 138; females, 164; total, 302; or 0.74 per cent. of all.

American parentage, 215; foreign parentage, 87.

Of the 302 decedents, 45.70 per cent. were males, and 54.30 per cent., females; 71.19 per cent. of American, and 28.81 per cent. of foreign parentage.

Of the eighteen decedents in 1883, eleven died at the Butler Hospital for the insane; in some years, a much larger proportion of the deaths from insanity occur at this institution, which is within the city limits, but where the patients, of course, are gathered from many different places.

Intemperance. There were 16 deaths reported from intemperance in 1883, including 3 from delirium tremens; the whole number was 2 less than in 1882; the number from delirium tremens, the same. The decedents from intemperance were:—

Males. 9; females, 7; total, 16; or 0.69 per cent. of all. American parentage, 4; foreign parentage, 12.

For the twenty-eight years, 1856-1883, the reported decedents from intemperance, were:—

Whole number, 293; annual average, 10.

Males, 202; females, 91; total, 293; or 0.72 per cent. of all.

American parentage, 82; foreign parentage, 211.

Of the 293 decedents, 68.94 per cent. were males; and 72.01 per cent., of foreign parentage.

Kidneys, Diseases of. The number of deaths from diseases of the kidneys in 1883, was 58, including 34 from "Bright's Disease." The number was 6 more than in 1882, all the increase being among those from Bright's Disease. The sex and parentage of the decedents were as follows:—

Males, 35; females, 23; total, 58; or 2.50 per cent. of all.
American parentage, 30; foreign parentage, 28.

For the twenty-eight years, 1856-1883:-

Whole number, 684; annual average, 24.

Males, 394; females, 290; total, 684; or 1.68 per cent. of all.

American parentage, 370; foreign parentage, 314.

Liver, Diseases of. Seventeen deaths, in 1883, were caused by diseases of the liver; 7 less than in 1882.

Males. 11; female, 6; total, 17; or 0.78 per cent. of all. American parentage, 5; foreign parentage, 12.

For the twenty-eight years, 1856-1883, the decedents from diseases of the liver, were:—

Whole number, 444; annual average, 16.

Males. 232; females. 212; total, 444; or 1 09 per cent. of all.

American parentage, 209; foreign parentage, 235.

Malformations. There were 13 deaths reported from malformation in 1883, which was 4 more than in the year previous. In one case the malformation was not specified; the rest were 6 cases of malformation of the heart; 3 of the spine; 2 of the rectum, and one of the anus. There were:—

Males, 5; females, 8; total. 13; or 0 56 per cent. of all.

American parentage, 9; foreign parentage, 4.

In the twenty-eight years, 1856-1883, there were:-

Males, 141; females, 92; total, 233; or 0.57 per cent. of all.

American parentage, 146; foreign parentage, 87.

Marasmus was assigned as the cause of 48 deaths in 1883; 9 more than in 1882; 41 of the decedents, or 85.42 per cent., were less than one year old.

Males, 34; females, 14; total, 48; or 2.07 per cent. of all. American parentage, 25; foreign parentage, 23.

The statistics for the twenty-eight years, 1856-1883, are:-

Whole number, 774; annual average, 27.

Males, 426; females, 348; total, 774; or 1.90 per cent. of all.

American parentage, 386; foreign parentage, 388.

Measles. There were 8 deaths from this cause reported in 1883; 4 more than in the preceding year.

Males, 6; females, 2; total, 8; or 0 35 per cent. of all. American parentage, 6; foreign parentage, 2.

The following table shows the number of deaths that have occurred from measles, in each year since 1856, and shows how irregular has been its prevalence in Providence.

1866	1864 7	1872 7	18790
1857 2	1865 7	187328	1880 8
185840	1866 9	1874 4	188125
1859 0	1887 0	1875 0	1882 4
1880 0	1868 4	1876 0	1883 8
1861 6	1869 6	1877 2	
1862 1	187016	187850	Total, 28 yrs247
186316	1871 1		• •

Nearly all the deaths reported from measles, are really from a secondary cause which is generally pneumonia.

The following are the particulars of the decedents from measles during the twenty-eight years, 1856-1883:—

Whole number, 247; annual average, 9.

Males, 117; females, 130; total, 247; or 0.61 per cent. of all.

American parentage, 76; foreign parentage, 171.

Of the 152 decedents, 92, or 60.53 per cent. were less than 15 years old.

Old Age. The number of deaths ascribed to old age in 1883, was 84; 5 more than in 1882. Twelve of the decedents were over 90 years old. The 84 deaths were 3.62 per cent. of all deaths from known causes. But this was but a small part of all the aged decedents during the year, as the following figures show. The whole number of decedents over 70 years old, was 332, or 213 of American parentage, and 119 of foreign parentage; or 14.30 per cent. of all decedents from known causes.

The decedents reported from "old age," in 1883, were:-

Males, 27; females, 57; total, 84; or 3.62 per cent. of all. American parentage, 44; foreign parentage, 40.

For the twenty-eight years, 1856-1883:-

Whole number, 1.563; annual average, 56.

Males, 530; females, 1,033; total, 1,563; or 3.84 per cent. of all.

American parentage, 934; foreign parentage, 629.

Of the 1,563 decedents, 66.09 per cent. were females; and 59.76 per cent were of American parentage. The number of decedents, over 70 years of age, during this period, was 4,646; 11.41 per cent. of all from known causes; 3,154 of American, and 1,492 of foreign parentage.

Paralysis. There were 34 deaths from this cause in 1883, including 4 from Locomotor Ataxia.

Males, 16; females, 18; total, 34; or 1.46 per cent. of all. American parentage, 24; foreign parentage, 10.

In the twenty-eight years, 1856-1883, there were :-

Whole number, 527; annual average, 19.

Males, 268; females, 259; total, 527; or 1.29 per cent. of all.

American parentage, 378; foreign parentage, 149.

The following table shows the number of deaths from paralysis, and from apoplexy, in each of the twenty-eight years, since 1856:—

Total	Paralysis.	Apoplexy.	Years.	Total.	Paralysis.	Apoplexy.	Years.
54	24	28	1870	28	2	21	1853
6	28		1871	27	9	18	1857
5	20	81	1872	18	8	15	1858
5	21	86	1878	22	6	16	1859
						17	
		41				14	
						19	
		50				20	
		43				18	
						20	
		48				21	
		74				8 1	
						22	
11		84					
			1008	18	10		1009
	597	971				tal 98 voors	To

The decedents from both causes, for the twenty-eight years were:

Whole number, 1,498; annual average, 58.

Males, 788; females, 760; total, 1,498; or 3.68 per cent. of all.

American parentage, 1,042; foreign parentage, 456.

The decedents were 69.56 per cent. of American, and 30 44 per cent. of foreign parentage.

Peritonitis. The deaths from this cause in 1883, were 22; 5 more than in 1882.

Males, 6; females, 16; total, 22; or 0.95 per cent. of all. American parentage, 18; foreign parentage, 9.

In the twenty-eight years, from 1856 to 1883, the deaths have been as follows:—

Whole number, 288; annual average, 10.

Males, 123; females, 165; total, 288; or 0.71 per cent. of all.

American parentage, 157; foreign parentage, 131.

Preumonia. The number of deaths from pneumonia in 1883, was 188; 18 more than in 1882; and from congestion of the lungs, 16; 10 more than in the previous year. They occurred as follows in the different seasons of the year.

lst quarter	74	8d quarter	15
2d quarter	75	4th quarter	40

Of the 204 decedents, there were: -

Males, 109; females, 95; total, 204; or 8.79 per cent. of all.

American parentage, 89; foreign parentage, 115.

The decedents from preumonia and congestion of the lungs, during the twenty-eight years, 1856-1883, were as follows:—

Whole number, 3,063; annual average, 109.

Males, 1,578; females, 1,485; total, 3,063; or 7.52 per cent. of all.

American parentage, 1,496; foreign parentage, 1,567.

This disease is one of the most important, in this climate, and has destroyed, on an average for 28 years, 7.52 in each 100 of all the decedents in the city. We therefore give very full statistics in relation to it.

The following table gives the statistics of pneumonia, (including congestion of the lungs), showing the whole number, the season, and the percentage from known causes, of the deaths, with the sex, age, and parentage of the decedents from this cause, in Providence, during the twenty-eight years, from 1856 to 1883 inclusive:—

Sex; parentage; season; age; whole number, and percentage from known causes.

Total.	800 925 925 927 74 74 776 930 930 930 930 930 930 930 930 930 930	1,148 897 791	1,092
1883.	8200 :0788882	4658	108 488 488
386	TO DUNCTERSTRA	5555	104
,1881	202-0-14122Co-	2128	284
880	- 第20mmの本質整置数4~	8278	825
1870.	East seet stre	¥828	EH4
1878.	#120-44-51484	2222	5.94
1877.	2550 useduse4-	2222	\$28
1870.	422	4422	228
1875.	485m-4EFFEEEnu		7
1874.	Hardesensia.		
1873.	Famor-decera-	9222	299
879.	Seas deday	28.8	182
1871.	Виденния подда	20112	835
1870,	SSEC CONFESSO	2215	\$ ti z
1860.	2200 : 000211	2000	35.08
.808	Porsa-undializa	asen	988
1867.		85.55	\$0.1
1806.	0.00 - 0.00 - 0.00 0.00	28.2	200
1865.	Boos AutoBoom	2208	2=3
1864.	Maranaguandua.	2511	25.5
.883		82.8	498
1862.	Sunu :	16 91	25 16 20
1981.	gan- triaden		138
.0081	UX 44000000000000	22.2	28
		41.08	225
1828.	86220 4400000	8825	200
.1987.	*************	2502	200
1856, 1857, 1858, 1859.	\$504 (000 prod0)	821.2	253
PNEUMONIA AND CONG. OF LUNGS.	Age: Under 1 year 1 and under 2. 15 2 and under 3. 15 3 0. 10. 40. 40. 60. 60 50 0. 40. 60. 60 50 0. 70. 60. 60 50 0. 60. 60 50 0. 60. 60. 60 50 0. 60. 60. 60 50 0. 60. 60. 60 50 0. 60. 60 50 0. 60. 60 50 0. 60. 60 50 0. 60. 60 50 0. 60. 60 50 0. 60. 60 50 0. 6	First quarter Second quarter Third quarter Fourth quarter	First 4 months

The number of deaths, 204 from pneumonia, in 1883, was larger than in any previous year; but the percentage from pneumonia of all the deaths from known causes had been larger in four previous years, than in 1883.

Percentage. The percentage of the deaths from pneumonia, of all from known causes, has not specially increased during the last twenty-eight years, though it has been larger than the average, during the last five years. The average percentage from pneumonia for twenty-eight years, has been 7.52 in each 100. The percentage has been less than this average in eighteen, and greater in ten of the twenty-eight years.

Sex. The decedents in Providence, from pneumonia, in twenty-eight years, were 1,578 males, and 1,485 females; or 51.51 males, and 48.49 females in each 100 decedents. The percentage of each sex, in the living population, is 47.73 males, and 52.27 females in each 100. It would seem, therefore, that the male sex suffers more than the female, from pneumonia.

Parentage. The living population of Previdence consists of 46.07 of American, and 53.93 of foreign parentage in each 100. During 28 years, 1856-1883, the decedents from pneumonia were 1,496 of American and 1,567 of foreign parentage; or 48.84 American, and 51.16 foreign in each 100. This shows that the population of American parentage suffers more from pneumonia than the population of foreign parentage.

Season. Pneumonia is plainly affected by season, and increases with the increase of cold. But the cold of the early months of the year seems to be more fatal than that of the last months. In the aggregate for twenty-eight years, the greatest mortality is in January; but February, March, and April have, each, a greater mortality than December. So too, the first quarter, January-March, has nearly 400 more deaths than the last quarter, October-December, in the aggregate for twenty-eight years. Even the second quarter of the year, April-June, has a greater mortality from pneumonia than the last quarter.

The following shows the number of deaths in Providence, and the percentage, from pneumonia, in each quarter of the year, in the aggregate of deaths for 28 years, from 1856 to 1883 inclusive:—

	Deaths from	Percentage.	
Season.	Pneumonia.		
First quarter, January-March		87.32	
Second quarter, April-June			
Third quarter, July-September	297	9.70	
Fourth quarter, October-December	761	94.84	
Whole Year	8.068		

Dividing the year into periods of four months each, we have the following deaths, and percentages from pacumonia, in the aggregate mortality for twenty-eight years:—

•	Deaths from	
Season.	Pneumonia.	Percentage.
First four months, January-April	1,582	50.03
Second four months, May-August	652	21.28
Third four months, September-December	679	28.70
	—	
Whole Year		100.00

It would seem that it is not cold weather alone that causes pneumonia; the severe cold of November and December does not produce as much pneumonia as the continued, though not so severe cold, of the season from January to April. Possibly it is the well known depressing effect of continued cold that makes the very young and the very old more liable to pneumonia.

Age. Pneumonia is specially fatal to the very young and the very old. There are more deaths from this disease, in the first year of life, than in any other ten years.

Dividing the years of life into five periods, we have the following numbers and percentages of deaths from pneumonia, in the aggregate for twenty-eight years, 1856–1883:—

	Deaths from		
Periods.	Pneumonia.	Percentage	
One to five — 5 years		20.7	
Five to twenty — 15 years			
Twenty to fifty — 30 years			
Fifty to eighty — 30 years	910	29.7	
Eighty and over — 20 years			
Totals		100.0	

The decedents from pneumonia, under five years, and over fifty years, comprise 71.56 in each 100 of all the decedents from this disease, leaving only 28.44 in each 100, who were in the 45 years of life between 5 and 50 years.

Rheumatism. There were 11 deaths ascribed to rheumatism in 1883; 3 more than in 1882.

Males, 4; females, 7; total, 11; or 0.47 per cent. of all. American parentage, 6; foreign parentage, 5.

Scarlatina. The deaths from this cause in 1883, numbered only 16; 2 less than in 1882, and less than in any year since 1877. In the twenty-nine years, 1855–1883, there have been only four, when the number of deaths from scarlatina was less than in 1883. The decedents, in 1883, were one, under one year old; three between one and two years; seven between two and five; and five between five and ten.

Males, 7; females, 9; total, 16; or 0.69 per cent. of all. American parentage, 6; foreign parentage, 10.

For the twenty-nine years, 1855-1883, the general statistics of scarlatina are as follows:—

Whole number of deaths, 2,123; annual average, 73.

Males, 1,046; females, 1,077; total, 2,123; or 5.21 per cent. of all.

American parentage, 871; foreign parentage, 1,252.

The following table gives the particulars of sex, parentage, season, and locality, and the percentage from known causes, for the aggregate of five years, 1855–1859, and for each of the twenty-four years, 1860–1883:—

Total.	2,123	1,046	R71 1,252	296 193 193 113 113 113 113 113 113 113 113	25 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5
1883.	16	1-0	10	O1	0 :0 ::00
1882.	18.0	130	71	04 60 05 04 : 04 : 00	E : : : : : : : : : : : : : : : : : :
1881.	9.11	18	17	C 0 4 F 0 4 4 - 1 1 1 1 1	800 -400004
1880.	243	25 55	88.55	588888888585858	1,5,25,52
1879.	252 12.49	135	101	250 254 25 8 2 L	2522222
1878.	2.80	63 63	17	annu-netendas	
1877.	0.62	400	95 G3	00 04	100 motes
1876.	35	16	22	400 40 1224	4 ::
1875.	8.6.	53	53	25505004MBF#	2 2 2 1 8 2 4 2 C
1874.	2/18	140	191	82222222	2.8812.282
1873.	132	105	25.	88525555588	2000 00 0 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
1872.	1.20	81	×=	***************************************	E 440
1811.	1.69	200	0.27	201-4-01- : : : : : : : : : : : : : : : : : : :	01 -4 03 03 03 03 00 10
1870.	2.80	15	22	End a set to se	H :H-401-61-
1869.	138	17	86.55	815184410r4x	To Breto and
1868.	4.60	82	35	-401-4001444r4	\$:CH80#0163
1867.	1.06	20.1-	F- 00	-01 : : :- :01	401
1866.	8	H 01	01 FF		; i" ; i" ; i ;
1865.	108	22	20	SHEHEE BOA : OL	885058
. 1981	14.	35	92	-40070000188	1515888
1863.	25.55	13	115	H00400H H 1 10	x → c x x x x x x x x x x x x x x x x x
1862.	1.58	1000	82	- 101-4 11-0	60 04 04 04 00 04 ·
1881	25.	118	77	*	Success
1860. 1	1.1	10	119	8 1 981 1 1	401- 0100
1855-1859.	335	168	155	2845828 x 588	8122218
SCARLATINA. 1	Whole number	Males	PARENTAGE: American Foreign	SEASON: January January March. March. May June July Angust September Scorober November November	Locality: Ward II III IV V V V V V II II II II II II I

12 TO TT. 48.

men a class : : "Therentheis" W entile or reckoned hi an more il serenteen years, : oreign parentage, 8 entire caused by tume - varian : 6 uterine otal. 11: or 0.47 pe in stendage. . foreign parer ... were not tix deaths i -- via via i).26 per cent. wis 4 less, and th PRINTER :: total, 6; or services percentage, 3; for and the second Wante number, 5-A .ues 2 1; Semales, 278; American parenta: CA the field decedents to ern, were of foreign par

1856.
1857.
1858.
1859.
1860.
1861.
1862.
1863.
1864.
1866.
1867.
1867.
1871.
1872.
1873.
1874.
1875.
1876.
1876.
1876.

Evidences of full protection from small-pox are required of all children entering the public schools of Providence.

Spine, Diseases of. There were 4 deaths from diseases of the spine in 1883; 5 less than in 1882, and 0.17 per cent. of all deaths from known causes. The decedents were all females.

American parentage, 1; foreign parentage, 3.

Stomach, Diseases of. There were 21 deaths from diseases of the stomach in 1882, 16 being characterized as "inflammation;" the number was 3 less than in 1882.

Males. 5; females, 16; total, 21; or 0.91 per cent. of all. American parentage, 3; foreign parentage, 18.

Suicide. The number of deaths by suicide in 1883, was 15; 3 more than in 1882. Six were by shooting, and one each by taking arsenic, chloroform, corrosive sublimate, laudanum, morphia, Paris green, and strychnine; one person jumped from a window; and one into a well.

Males, 9; females, 6; total, 15; or 0.65 per cent. of all. American parentage, 8; foreign parentage, 7.

For the twenty-eight years, 1856-1883, the figures are:-

Males, 132; females, 56; total, 188; or 0.46 per cent. of all. American parentage, 105; foreign parentage, 83.

Teething. There were 14 deaths reported in 1883, as caused by teething; 2 less than in 1882.

Males, 8; females, 6; total, 14; or 0.60 per cent. of all. American parentage, 7; foreign parentage, 7.

Tuberculosis was given as the cause of 18 deaths in 1883; 6 more than in 1882.

Males, 8; females, 10; total, 18; or 0.78 per cent. of all.

American parentage, 10; foreign parentage, 8.

There are several diseases which might properly be brought together under this head; but they have never been so classified in

these reports; if it were done, the percentage of such a class would exceed that of any other. The heading of "tuberculosis" was first used separately in 1867; since then the deaths so reckoned have been as follows:—

Whole number from known causes in seventeen years, 29,421.

From tuberculosis, 236; annual average, 14.

Males, 106; females, 130; total, 236; or 0.80 per cent. of all.

American parentage, 152; foreign parentage, 84.

Tumors. There were 11 deaths caused by tumors in 1883; 5 more than in 1882. Three were ovarian; 6 uterine; and 2 were of the brain.

Males, 0; females, 11; total, 11; or 0.47 per cent. of all.

American parentage, 7; foreign parentage, 4.

Unknown. There were but six deaths in 1883, for which no cause was assigned; this was 0.26 per cent. of all the deaths during the year. The number was 4 less, and the per cent., 0.19 less than in 1882.

Males, 5; females, 1; total, 6; or 0 26 per cent. of all.

American parentage, 3; foreign parentage, 3.

The statistics of the "unknown," for the twenty-eight years, 1856-1883, are:—

Whole number, 581; annual average, 21.
Males, 803; females, 278; total. 581; or 1.43 per cent. of all.
American parentage, 199; foreign parentage, 382.

Of the 581 decedents reported from unknown causes, 65.75 per cent. were of foreign parentage.

STILL-BORN.

The following statistics are given concerning the still-born children in Providence in 1883.

Number. There were 162 still-born children in 1883; the same number as in 1882. This number was 5.55 per cent of the 2,916 children born during the year 1883, or one in 18.

Sex. There were 84 male children; 12 less than in 1882; and 78 female children, or 12 more than in the previous year.

Color. White, 153; colored, 9.

Locality. First ward, 23; second, 4; third, 19; fourth, 6; fifth, 19; sixth, 2; seventh, 14; eighth, 17; ninth, 24; tenth, 84.

Season. January, 14; February, 9; March, 10; April, 18; May, 16; June, 18; July, 10; August, 15; September, 24; October, 12; November, 8; December, 8.

The number and percentage in each quarter of the year were:-

	Number.	Per cent.		Number.	Per cent.
1st quarter	83	20.87	8d quarter	49	30.25
2d quarter	52	32.10	4th quarter	28	17 28

Parentage. The following table shows the parentage of the 162 still-born children in 1883; the first letter indicating the birth-place of the father; the second, that of the mother:—

Reckoned according to the father's birth-places, 1,463, or 43.89 per cent., were American; 1,386, or 41.58 per cent. were Irish; and 477, or 14.31 per cent. were children of other foreign fathers; of 7, the parentage was not stated.

The following figures show the whole number of children borne by mothers of different nativities, during the twenty-eight years, 1856–1883 inclusive; also the number of still-born children to mothers of each class; and the percentage of the latter of the whole number of children:—

	Whole number	Still-born	Percentage	Or 1
	Of children.	children.	of still-born.	in
American mothers	27,395	1,461	5.33	18.75
Irish mothers		1,454	6.81	14.69
English, Scotch or Welsh mo	thers 4,447	230	4.95	90.21
German mothers	1,166	38		30.66
British American mothers	•			
Other foreign mothers	1,095	65	5.94	16.85
Total	57 558	3 333	5 79	17.97

BIRTHS AND DEATHS BY PARENTAGE.

The following recapitulatory table shows the number of births and deaths, by parentage, in Providence, in each of the twenty-eight years, from 1856–1883 inclusive. In the three years, 1863, 1864, and 1865, there was an excess of deaths over the births, in the population of American parentage:—

TABLE XXII.

	AMERI	CAN PARE	NTAGE.	Foreign Parentage.		NTAGE.
YEAR.	Deaths.	Births.	Excess of Births	Deaths.	Births.	Excess of Births
1856	541	759	218	524	916	892
1857	528	701	178	402	987	585
1856	512	771	259	505	953	448
1859	479	659	180	420	984	514
1860	512	781	219	489	917	428
1861	538	740	202	513	985	479
1802	460	663	208	454	866	419
1868	597	528	09	617	779	162
1864	619	579	—70	682	765	133
1865	631	546	—85	580	752	172
1866	513	726	218	523	808	888
1867	528	725	197	432	899	467
1868	596	883	286	514	984	470
1869	627	926	299	629	893	964
1870	642	890	248	621	999	878
1871	648	980	837	611	980	869
1872	832	1,035	208	771	1,171	400
1878	863	1,008	141	857	1,125	268
1874	885	1,217	332	1,102	1,563	461
1875	914	1,180	266	1,001	1,484	483
1876	875	1,169	294	990	1.462	479
1877	908	1,238	880	1,080	1,805	275
1878	1689	1,215	826	1,100	1,412	812
1879	941	1,328	2×7	1,085	1.294	209
1880	920	1,252	832	1,160	1,375	215
1881	999	1,365	366	1.146	1,438	292
1882.,	965	1,870	405	1,277	1,418	141
1883	1,047	1,457	410	1,281	1,459	178
28 years	20,028	26,535	6 507	21,966	81,021	9,755



Vital Statistics, Providence, R. L. 1883.

CITY REGISTRARS

TWENTY-NINTH ANNUAL REPORT

.

. • • i

. .

1 . . • •

.

.

.

.

.

•

.

.

. .

